### (19) World Intellectual Property Organization International Bureau



# 

### (43) International Publication Date 18 January 2001 (18.01.2001)

### PCT

### (10) International Publication Number WO 01/04270 A1

- (51) International Patent Classification7: C12N 5/10, 15/12, 15/63, 15/64, C07K 14/47, 14/705, 14/715, G01N 33/53, 33/567
- (21) International Application Number: PCT/US00/19094
- (22) International Filing Date:

13 July 2000 (13.07.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/143,435

13 July 1999 (13.07.1999) US

- (71) Applicant (for all designated States except US): GENAIS-SANCE PHARMACEUTICALS, INC. [US/US]; Five Science Park, New Haven, CT 06511 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CHEW, Anne [US/US]; 837 Orange Street, Apartment 2, New Haven, CT 06511 (US). DENTON, R., Rex [US/US]; 129 Hunters Trail, Madison, CT 06443 (US). DUDA, Amy [US/US]; 93 Marvel Road, New Haven, CT 06515 (US), NAND-ABALAN, Krishnan [IN/US]; 228 Village Pond Road, Guilford, CT 06437 (US). STEPHENS, Joel, Claiborne [US/US]; 46 Crabapple Lane, Guilford, CT 06437 (US). WINDEMUTH, Andreas, K. [US/US]; 91 Center Road, Woodbridge, CT 06524 (US).

- (74) Agent: HENDERSON, Melodie, W.; Genaissance Pharmaceuticals, Inc., Five Science Park, New Haven, CT 06511 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, S1, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

### DRUG TARGET ISOGENES:

# POLYMORPHISMS IN THE INTERLEUKIN 4 RECEPTOR ALPHA GENE

### RELATED APPLICATIONS

5

10

15

20

25

30

35

This application is a continuation-in-part of, and claims priority to, U.S. Provisional Application Serial No. 60/143,435 filed July 13, 1999.

### FIELD OF THE INVENTION

This invention relates to variation in genes that encode pharmaceutically important proteins. In particular, this invention provides genetic variants of the human Interleukin 4 Receptor Alpha (II.4Ra) gene and methods for identifying which variant(s) of this gene is/are possessed by an individual.

### BACKGROUND OF THE INVENTION

Current methods for identifying pharmaceuticals to treat disease often start by identifying, cloning, and expressing an important target protein related to the disease. A determination of whether an agonist or antagonist is needed to produce an effect that may benefit a patient with the disease is then made. Then, vast numbers of compounds are screened against the target protein to find new potential drugs. The desired outcome of this process is a drug that is specific for the target, thereby reducing the incidence of the undesired side effects usually caused by a compound's activity at non-intended targets.

What this approach fails to consider, however, is that natural variability exists in any and every population with respect to a particular protein. A target protein currently used to screen drugs typically is expressed by a gene cloned from an individual who was arbitrarily selected. However, the nucleotide sequence of a particular gene may vary tremendously among individuals. Subtle alteration(s) in the primary nucleotide sequence of a gene encoding a target protein may be manifested as significant variation in expression of or in the structure and/or function of the protein. Such alterations may explain the relatively high degree of uncertainty inherent in treatment of individuals with drugs whose design is based upon a single representative example of the target. For example, it is well-established that some classes of drugs frequently have lower efficacy in some individuals than others, which means such individuals and their physicians must weigh the possible benefit of a larger dosage against a greater risk of side effects. In addition, variable information on the biological function or effects of a particular protein may be due to different scientists unknowingly studying different isoforms of the gene encoding the protein. Thus, information on the type and frequency of genomic variation that exists for pharmaceutically important proteins would be useful.

The organization of single nucleotide variations (polymorphisms) in the primary sequence of a gene into one of the limited number of combinations that exist as units of inheritance is termed a haplotype. Each haplotype therefore contains significantly more information than individual unorganized polymorphisms. Haplotypes provide an accurate measurement of the genomic variation in the two chromosomes of an individual.

It is well-established that many diseases are associated with specific variations in gene sequences. However while there are examples in which individual polymorphisms act as genetic markers for a particular phenotype, in other cases an individual polymorphism may be found in a variety of genomic backgrounds and therefore shows no definitive coupling between the polymorphism and the causative site for the phenotype (Clark AG et al. 1998 Am J Hum Genet 63:595-612; Ulbrecht M et al. 2000 Am J Respir Crit Care Med 161: 469-74). In addition, the marker may be predictive in some populations, but not in other populations (Clark AG et al. 1998 supra). In these instances, a haplotype will provide a superior genetic marker for the phenotype (Clark AG et al. 1998 supra; Ulbrecht M et al. 2000, supra; Ruaño G & Stephens JC Gen Eng News 19 (21), December 1999).

5

10

15

20

25

30

Analysis of the association between each observed haplotype and a particular phenotype permits ranking of each haplotype by its statistical power of prediction for the phenotype. Haplotypes found to be strongly associated with the phenotype can then have that positive association confirmed by alternative methods to minimize false positives. For a gene suspected to be associated with a particular phenotype, if no observed haplotypes for that gene show association with the phenotype of interest, then it may be inferred that variation in the gene has little, if any, involvement with that phenotype (Ruaño & Stephens 1999, *supra*). Thus, information on the observed haplotypes and their frequency of occurrence in various population groups will be useful in a variety of research and clinical applications.

One possible drug target for the treatment of allergies, asthma, and other immune responses is the Interleukin 4 Receptor Alpha (IL4Rα) gene or its encoded product. IL4R is a transmembrane complex composed of two different protein subunits, a 140-kDa high affinity binding subunit named interleukin-4 receptor α (IL-4Rα; also known as CD124 antigen) and either a gamma-c subunit, which is present in several cytokine receptors, or an interleukin-13 receptor 1 (IL-13R1) subunit. Both subunits of the IL-4R are required to bind interleukin-4 (IL-4) and to mediate its transcription-activating effects through the tyrosine kinases, Jak1 and Jak3. Upon binding of IL-4 to the IL-4R, Jak 1 and Jak 3 phosphorylate the IL-4Rα subunit, creating binding sites in the cytoplasmic domain for many other proteins, including SOS, Stat-6, c-fes, and *src* homology phosphatase 1 (SHP-1). Activated Jak proteins also phosphorylate Stat-proteins, which travel into the nucleus and function as transcription factors. Other IL-4 signal pathways exist, but are less well characterized.

IL-4, one of the most important cytokines involved in the allergic response, is produced when cells from the immune system, in particular T cells, are activated in response to an allergen. Regulation of the immune response involves Helper T-cells that differentiate into two subtypes, Th1 and Th2. Th1 cells express interferon-gamma and interluckin-2 (IL-2), and mediate a cell-based immunity, where macrophages and neutrophils are prominently involved. Th1 cells also direct the IgE-producing B cells, as well as mast cells, basophils and eosinophils. Th2 cells produce IL-4, IL-5, IL-6, IL-10 and IL-13. Each Th cell subtype represses the other, so the immune system is forced into differentiation into either a Th1 or Th2 response against an external allergic challenge. In many instances an aberration of this response can render a pathological state such as a Th2 response against ragweed.

IL-4 induces in B cells the synthesis of IgE type antibodies that recognize specific allergens. IgE binds to receptors on mast cells and basophils and mediates the early humoral (sub-chronic) response on the B-side of the immune system. If an allergen binds mast cell-attached IgE, the mast cell releases mediators like histamine, and the eicosanoid leukotrienes and prostaglandins products, some of which cause the familiar symptoms of an acute allergic reaction: swelling, itching, mucous, and reddening of the skin. Later in this process eosinophils and other inflammatory cells migrate to the site of inflammation. This later phase is important in asthma, because the eosinophils may instigate a more chronic inflammation which can adversely scar lung tissue. IL-4 is at least partly responsible for recruiting eosinophils, because it induces synthesis of specific adhesion molecules on the capillary endothelium, and stimulates expression of IL-5 and eotaxin. IL-5 leads to the development of a large number of eosinophils from precursor cells in the bone marrow, and eotaxin stimulates their migration into the lung tissue.

It has been proposed that inhibition of IL-4 activity would disarm the Th2 component of the immune system. This would then allow the immune system to develop a natural tolerance towards common allergens without the full acute response to the challenge. In this way, tolerance may be induced in many patients, similar to what is sometimes achieved with hyposensitization shots for allergy patients. Thus, substances that inhibit IL-4 production and/or its binding to the IL-4 receptor, may improve the therapy of allergies and asthma.

The gene for IL-4Rα has eleven exons encoding an 825 amino acid protein and spans over 24 kb of the short arm of chromosome 16 (16p12.1) (Pritchard et al., *Genomics* 10:801, 1991; GenBank Accession No. AC004525). A reference sequence for IL4R-α gene, which corresponds to the reverse complement of nucleotides 100020-71331 in the GenBank Accession No. AC004525, is shown in Fig. 1 (SEQ ID NO:1). Reference sequences for IL-4Rα mRNA (GenBank Accession No. NM\_000418) and the encoded IL4Rα precursor protein (GenBank Accession No. P24394) are shown in Figs. 2 and 3, respectively (SEQ ID NOS:2 and 3). Significant features reported for the IL-4Rα precursor include: a signal peptide located between a.a. 1 and 25; an extracellular domain between a.a. 26 and 232; disulfide bonds between a.a. 34 and 44 and between a.a. 74 and 86; glycosylation sites at amino acids 53, 98, 128, 134, 176, and 209; a transmembrane region between a.a. 233 and 256; and a cytoplasmic domain between a.a. 257 and 825.

Recently, several studies have suggested that genetic polymorphisms in the IL4-R $\alpha$  gene are associated with genetic predisposition to atopy and/or elevated serum IgE. Mitsuyasu et al., reported that polymorphisms at codons 75 and 576 affect IL-4R function (*Nat. Genet.* 19:119-120, 1998). The IL-4R $\alpha$  allele with isoleucine at amino acid position 75 (Ile75) in the extracellular domain is more responsive to IL-4 than the allele with valine at that position (Val75) and is associated with atopic asthma but not with non-atopic asthma (Mitsuyasu et al., *supra*). Also, the allele with arginine at position 576 (Arg576) in the cytoplasmic domain exhibits higher receptor activity than the glutamine allele (Glu576) due to reduced

binding by the Arg576 allele of a negative regulatory molecule, src homology phosphatase 1 (Imani et al., J. Biol Chem. 272:7927-7931, 1997). The Arg576 allele has a higher frequency in patients with allergic inflammatory disorders, including atopy (Khurana Hershey et al., New Eng. J. Med. 337:1720-1725. 1997). In a recent study, Hershey et al.; (WO 00/34789) reported that the Arg576 allele is significantly associated with asthma. Studies showed that patients who were homozygous for this allele had about a 9fold higher risk towards asthma and that two copies of Arg576 are associated in an increase in asthma prevalence and severity. Kanemitsu et al. recently reported that the presence of either the Ile75 or Arg576 variant is significantly associated with susceptibility for developing systemic lupus erythematosus (SLE), a Th2-dominant systemic autoimmune disorder (Arthritis Rheum. 42:1298-1300). Another variant IL-4Rα allele that is reportedly associated with atopy susceptibility has proline rather than serine at position 503 (Kruse et al., Immunol. 96:365-371, 1999). Other IL-4Ra gene polymorphisms leading to amino acid changes in the cytoplasmic domain of the protein product have been identified at codons 400 (E400A). 431 (C431R) and 786 (S786P) (Kruse et al., supra; Deichmann et al., Biochem. Biophys. Res. Commun. 231:696-697, 1997). A polymorphism of guanine or adenine at a position corresponding to nucleotide 55328 in Figure 1 has also been reported as well as a polymorphism of cytosine or thymine at a position corresponding to nucleotide 55430 (Buetow et al., 1999, Nat Genet. 21:323-5).

Because of the potential for polymorphisms in the ILAR $\alpha$  gene to affect the expression and function of the encoded protein, it would be useful to determine whether additional polymorphisms exist in the ILAR $\alpha$  gene, as well as how such polymorphisms are combined in different copies of the gene. Such information would be useful for studying the biological function of ILAR $\alpha$  as well as in identifying drugs targeting this protein for the treatment of disorders related to its abnormal expression or function.

## SUMMARY OF THE INVENTION

10

15

20

25

30

35

Accordingly, the inventors herein have discovered 38 novel polymorphic sites in the IL4Ra gene. These polymorphic sites (PS) correspond to the following nucleotide positions in the reverse complement of the indicated GenBank Accession Number: 32884 (PS1), 32903 (PS2), 32961 (PS3), 33135 (PS4), 35763 (PS6), 35770 (PS7), 35817 (PS8), 35905 (PS9), 35944 (PS10), 35958 (PS11), 37330 (PS12), 37473 (PS13), 37586 (PS14), 37591 (PS15), 37604 (PS16), 37644 (PS17), 37678 (PS18), 43446 (PS19), 43703 (PS20), 53008 (PS21), 53099 (PS22), 53153 (PS23), 53456 (PS25), 53507 (PS27), 53513 (PS28), 53915 (PS30), 53949 (PS32), 54237 (PS33), 54468 (PS34), 54611 (PS35), 54698 (PS36), 54700 (PS37), 54741 (PS38), 54780 (PS39), 55083 (PS40), 55142 (PS41), 55539 (PS44) and 55758 (PS45) in AC004525. The polymorphisms at these sites are adenine or guanine at PS1, cytosine or thymine at PS2, guanine or thymine at PS3, guanine or cytosine at PS4, cytosine or thymine at PS6, guanine or adenine at PS7, thymine or cytosine at PS8, cytosine or thymine at PS9, cytosine or thymine at PS10, guanine or adenine at PS11, guanine or adenine at PS12, cytosine or thymine at PS13, cytosine or thymine at PS14, guanine or adenine at PS15, adenine or thymine at PS16, cytosine or adenine at PS17, cytosine or

thymine at PS18, guanine or adenine at PS19, thymine or cytosine at PS20, adenine or cytosine at PS21, cytosine or thymine at PS22, thymine or cytosin at PS23, guanine or thymine at PS25, cytosine or thymine at PS27, thymine or cytosine at PS28, cytosine or thymine at PS30, guanine or adenine at PS32, cytosine or thymine at PS33, thymine or guanine at PS34, thymine or cytosine at PS35, thymine or cytosine at PS36, thymine or cytosine at PS37, cytosine or thymine at PS38, cytosine or guanine at PS39, adenine or guanine at PS40, guanine or adenine at PS41, cytosine or thymine at PS44 and guanine or adenine at PS45. In addition, the inventors have determined the identity of the alternative nucleotides present at these sites, as well as at the previously identified sites at nucleotides 35749 (PS5), 53413 (PS24), 53505 (PS26), 53721 (PS29), 53941 (PS31), 55328 (PS42), and 55430 (PS43). It is believed that IL4Rα-encoding polynucleotides containing one or more of the novel polymorphic sites reported herein will be useful in studying the expression and biological function of ILARa, as well as in developing drugs targeting this protein. In addition, information on the combinations of polymorphisms in the IL4Ra gene may have diagnostic and forensic applications.

10

20

25

30

35

Thus, in one embodiment, the invention provides an isolated polynucleotide comprising a 15 nucleotide sequence which is a polymorphic variant of a reference sequence for the ILARα gene or a fragment thereof. The reference sequence comprises SEQ ID NO:1 and the polymorphic variant comprises at least one polymorphism selected from the group consisting of guanine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, cytosine at PS8, thymine at PS9, thymine at PS10, adenine at PS11, adenine at PS12, thymine at PS13, thymine at PS14, adenine at PS15, thymine at PS16, adenine at PS17, thymine at PS18, adenine at PS19, cytosine at PS20, cytosine at PS21, thymine at PS22, cytosine at PS23, thymine at PS25, thymine at PS27, cytosine at PS28, thymine at PS30, adenine at PS32, thymine at PS33, guanine at PS34, cytosine at PS35, cytosine at PS36, cytosine at PS37, thymine at PS38, guanine at PS39, guanine at PS40, adenine at PS41, thymine at PS44, and adenine at PS45. In a preferred embodiment, the polymorphic variant comprises one or more additional polymorphisms selected from the group consisting of guanine at PS5, cytosine at PS24, cytosine at PS26, cytosine at PS29, guanine at PS31, adenine at PS42, and thymine at PS43. A particularly preferred polymorphic variant is a naturally-occurring isoform (also referred to herein as an "isogene") of the IL4Rα gene. An IL4Rα isogene of the invention comprises adenine or guanine at PS1, cytosine or thymine at PS2, guanine or thymine at PS3, guanine or cytosine at PS4, cytosine or thymine at PS6, guanine or adenine at PS7, thymine or cytosine at PS8, cytosine or thymine at PS9, cytosine or thymine at PS10, guanine or adenine at PS11, guanine or adenine at PS12, cytosine or thymine at PS13, cytosine or thymine at PS14, guanine or adenine at PS15, adenine or thymine at PS16, cytosine or adenine at PS17, cytosine or thymine at PS18, guanine or adenine at PS19, thymine or cytosine at PS20, adenine or cytosine at PS21, cytosine or thymine at PS22, thymine or cytosine at PS23, guanine or thymine at PS25, cytosine or thymine at PS27, thymine or cytosine at PS28, cytosine or thymine at PS30, guanine or adenine at PS32, cytosine or thymine at PS33, thymine or guanine at PS34, thymine or cytosine at PS35, thymine or cytosine at PS36, thymine or cytosine at PS37, cytosine or thymine at PS38, cytosine or

guanine at PS39, adenine or guanine at PS40, guanine or adenine at PS41, cytosine or thymine at PS44 and guanine or adenine at PS45. The invention also provides a collection of IL4R $\alpha$  isogenes, referred to herein as an IL4R $\alpha$  genome anthology.

An ILAR $\alpha$  isogene may be defined by the combination and order of these polymorphisms in the isogene, which is referred to herein as an ILAR $\alpha$  haplotype. Thus, the invention also provides data on the number of different ILAR $\alpha$  haplotypes found in the above four population groups. This haplotype data is useful in methods for deriving an ILAR $\alpha$  haplotype from an individual's genotype for the ILAR $\alpha$  gene and for determining an association between an ILAR $\alpha$  haplotype and a particular trait.

5

10

15

20

25

30

35

In another embodiment, the invention provides a polynucleotide comprising a polymorphic variant of a reference sequence for an ILAR $\alpha$  cDNA or a fragment thereof. The reference sequence comprises SEQ ID NO:2 (Fig. 2) and the polymorphic cDNA comprises at least one polymorphism selected from the group consisting of thymine at a position corresponding to nucleotide 237, adenine at a position corresponding to nucleotide 244, cytosine at a position corresponding to nucleotide 291, thymine at a position corresponding to nucleotide 501, adenine at a position corresponding to nucleotide 554, cytosine at a position corresponding to nucleotide 939, thymine at a position corresponding to nucleotide 1242, thymine at a position corresponding to nucleotide 1293, cytosine at a position corresponding to nucleotide 1299, thymine at a position corresponding to nucleotide 1701, adenine at a position corresponding to nucleotide 2023, guanine at a position corresponding to nucleotide 2254 and cytosine at a position corresponding to nucleotide 2397. In a preferred embodiment, the polymorphic variant comprises one or more additional polymorphisms selected from the group consisting of guanine at a position corresponding to 223, cytosine at a position corresponding to nucleotide 1199, cytosine at a position corresponding to 1291, cytosine at a position corresponding to nucleotide 1507 and guanine at a position corresponding to 1737.

Polynucleotides complementary to these IL4R $\alpha$  genomic and cDNA variants are also provided by the invention.

In other embodiments, the invention provides a recombinant expression vector comprising one of the polymorphic genomic variants operably linked to expression regulatory elements as well as a recombinant host cell transformed or transfected with the expression vector. The recombinant vector and host cell may be used to express ILAR $\alpha$  for protein structure analysis and drug binding studies.

In yet another embodiment, the invention provides a polypeptide comprising a polymorphic variant of a reference amino acid sequence for the ILAR $\alpha$  protein. The reference amino acid sequence comprises SEQ ID NO:3 (Fig. 3) and the polymorphic variant comprises at least one variant amino acid selected from the group consisting of threonine at a position corresponding to amino acid 82, histidine at a position corresponding to amino acid 185, isoleucine at a position corresponding to amino acid 579, serine at a position corresponding to amino acid 675, and alanine at a position corresponding to amino acid 752. In some embodiments, the polymorphic variant also comprises at least one variant amino acid

selected from the group consisting of valine at a position corresponding to amino acid 75, alanine at a position corresponding to amino acid 400, arginine at a position corresponding to amino acid 431, proline at a position corresponding to amino acid 503, and arginine at a position corresponding to amino acid 576. A polymorphic variant of ILAR $\alpha$  is useful in studying the effect of the variation on the biological activity of ILAR $\alpha$  as well as studying the binding affinity of candidate drugs targeting ILAR $\alpha$  for the treatment of allergies, asthma, and other immune responses.

5

10

15

20

25

30

35

The present invention also provides antibodies that recognize and bind to the above polymorphic IL4R $\alpha$  protein variant. Such antibodies can be utilized in a variety of diagnostic and prognostic formats and therapeutic methods.

In other embodiments, the invention provides methods, compositions, and kits for haplotyping and/or genotyping the IL4Rα gene in an individual. The methods involve identifying the nucleotide or nucleotide pair present at one or more polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45 in one or both copies of the IL4Rα gene from the individual. The compositions contain oligonucleotide probes and primers designed to specifically hybridize to one or more target regions containing, or that are adjacent to, a polymorphic site. The methods and compositions for establishing the genotype or haplotype of an individual at the novel polymorphic sites described herein are useful for studying the effect of the polymorphisms in the etiology of diseases affected by the expression and function of the IL4Rα protein, studying the efficacy of drugs targeting IL4Rα, predicting individual susceptibility to diseases affected by the expression and function of the IL4Rα protein and predicting individual responsiveness to drugs targeting IL4Rα.

In yet another embodiment, the invention provides a method for identifying an association between a genotype or haplotype and a trait. In preferred embodiments, the trait is susceptibility to a disease, severity of a disease, the staging of a disease or response to a drug. Such methods have applicability in developing diagnostic tests and therapeutic treatments for allergies, asthma, and other immune responses.

The present invention also provides transgenic animals comprising one of the ILAR $\alpha$  genomic polymorphic variants described herein and methods for producing such animals. The transgenic animals are useful for studying expression of the ILAR $\alpha$  isogenes in vivo, for in vivo screening and testing of drugs targeted against ILAR $\alpha$  protein, and for testing the efficacy of therapeutic agents and compounds for allergies, asthma, and other immune responses in a biological system.

The present invention also provides a computer system for storing and displaying polymorphism data determined for the ILAR $\alpha$  gene. The computer system comprises a computer processing unit; a display; and a database containing the polymorphism data. The polymorphism data includes the polymorphisms, the genotypes and the haplotypes identified for the ILAR $\alpha$  gene in a reference

population. In a preferred embodiment, the computer system is capable of producing a display showing IL4Rα haplotypes organized according to their evolutionary relationships.

### BRIEF DESCRIPTION OF THE DRAWINGS

5

10

15

20

25

30

Figure 1 illustrates a reference sequence for the IL4R $\alpha$  gene that is the reverse complement of part of Genbank Accession Number AC004525.1; contiguous lines; SEQ ID NO:1), with the underlines indicating the start and stop codons, shading indicating the reference coding sequence, and the bold nucleotides indicating the polymorphic sites and polymorphisms identified by Applicants in a reference population.

Figure 2 illustrates a reference sequence for the IL4Ra coding sequence (GenBank Accession Number X52425; contiguous lines; SEQ ID NO:2), with the underlines indicating the start and stop codons, and the bold nucleotides indicating the polymorphic sites and polymorphisms identified by Applicants in a reference population.

Figure 3 illustrates a reference sequence for the IL4Ra protein (GenBank Accession Number CAA36672; contiguous lines; SEQ ID NO:3), with the bold amino acids indicating the amino acid variations caused by the polymorphisms of Fig. 2.

### DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is based on the discovery of novel variants of the ILAR $\alpha$  gene. As described in more detail below, the inventors herein discovered 38 novel polymorphic sites by characterizing the ILAR $\alpha$  gene found in genomic DNAs isolated from Index Repository IA that contains immortalized cell lines from one chimpanzee and 93 human individuals and Index Repository IB that contains 70 human individuals. Theses two repositories contain 51 individuals in common.

The human individuals in Index Repository IA included a reference population of 79 unrelated individuals self-identified as belonging to one of four major population groups: Caucasian (22 individuals), African descent (20 individuals) Asian (20 individuals) Hispanic/Latino (17 individuals). To the extent possible, the members of this reference population were organized into population subgroups by the self-identified ethnogeographic origin of their four grandparents as shown in Table 1 below. In addition, Index Repository IA contains three unrelated indigenous American Indians (one from each of North, Central, and South America), one three-generation Caucasian family (From the CEPH Utah cohort) and one two-generation African-American family.

Table 1. Population Groups in Index Repository IA

Population Group	Population Subgroup	No. of Individuals
African descent		. 20
	Sierra Leone	1
Asian		. 20
•	Burma	1
	China	3
	Japan	6 .
	. Korea	1
	Philippines	5
	Vietnam	4
Caucasian		22
	British Isles	3
	British Isles/Central	4
	British Isles/Eastern	1
	Central/Eastern.	1
	Eastern	3
	Central/Mediterranean	1
	Mediterranean	2
	Scandinavian	2
Hispanic/Latino		17
	Caribbean	7
	Caribbean (Spanish Descent)	2
	Central American (Spanish Descent)	1
	Mexican American	4
	South American (Spanish Descent)	3

Index Repository IB contains a reference population of 70 human individuals comprised of 4 three-generation families (from the CEPH Utah cohort) as well as unrelated African-American, Asian, and Caucasian individuals. A total of 38 individuals in this reference population are unrelated.

Using the ILAR $\alpha$  genotypes identified in the Index Repositories and the methodology described in the Examples below, the inventors herein also determined the haplotypes found on each chromosome for most human members of this repository. The ILA $\alpha$  genotypes and haplotypes found in the Index Repositories include those shown in Tables 4 and 5, respectively. The polymorphism and haplotype data disclosed herein are useful for studying population diversity, anthropological lineage, the significance of diversity and lineage at the phenotypic level, paternity testing, forensic applications, and for identifying associations between the ILAR $\alpha$  genetic variation and a trait such as level of drug response or susceptibility to disease.

10

15

In the context of this disclosure, the following terms shall be defined as follows unless otherwise indicated:

Allele - A particular form of a genetic locus, distinguished from other forms by its particular nucleotide sequence.

Candidate Gene - A gene which is hypothesized to be responsible for a disease, condition, or the

response to a treatment, or to be correlated with one of these.

10

15

20

25

30

35

Gene - A segment of DNA that contains all the information for the regulated biosynthesis of an RNA product, including promoters, exons, introns, and other untranslated regions that control expression.

Genotype – An unphased 5' to 3' sequence of nucleotide pair(s) found at one or more polymorphic sites in a locus on a pair of homologous chromosomes in an individual. As used herein, genotype includes a full-genotype and/or a sub-genotype as described below.

Full-genotype – The unphased 5' to 3' sequence of nucleotide pairs found at all known polymorphic sites in a locus on a pair of homologous chromosomes in a single individual.

Sub-genotype – The unphased 5' to 3' sequence of nucleotides seen at a subset of the known polymorphic sites in a locus on a pair of homologous chromosomes in a single individual.

Genotyping - A process for determining a genotype of an individual.

Haplotype – A 5' to 3' sequence of nucleotides found at one or more polymorphic sites in a locus on a single chromosome from a single individual. As used herein, haplotype includes a full-haplotype and/or a sub-haplotype as described below.

Full-haplotype – The 5' to 3' sequence of nucleotides found at all known polymorphic sites in a locus on a single chromosome from a single individual.

Sub-haplotype – The 5' to 3' sequence of nucleotides seen at a subset of the known polymorphic sites in a locus on a single chromosome from a single individual.

Haplotype pair - The two haplotypes found for a locus in a single individual.

**Haplotyping** – A process for determining one or more haplotypes in an individual and includes use of family pedigrees, molecular techniques and/or statistical inference.

Haplotype data - Information concerning one or more of the following for a specific gene: a listing of the haplotype pairs in each individual in a population; a listing of the different haplotypes in a population; frequency of each haplotype in that or other populations, and any known associations between one or more haplotypes and a trait.

Isoform – A particular form of a gene, mRNA, cDNA or the protein encoded thereby, distinguished from other forms by its particular sequence and/or structure.

Isogene – One of the isoforms of a gene found in a population. An isogene contains all of the polymorphisms present in the particular isoform of the gene.

Isolated — As applied to a biological molecule such as RNA, DNA, oligonucleotide, or protein, isolated means the molecule is substantially free of other biological molecules such as nucleic acids, proteins, lipids, carbohydrates, or other material such as cellular debris and growth media. Generally, the term "isolated" is not intended to refer to a complete absence of such material or to absence of water, buffers, or salts, unless they are present in amounts that substantially interfere with the methods of the present invention.

Locus - A location on a chromosome or DNA molecule corresponding to a gene or a physical or phenotypic feature.

Naturally- ccurring – A term used to designate that the object it is applied to, e.g., naturally-occurring polynucleotide or polypeptide, can be isolated from a source in nature and which has not been intentionally modified by man.

Nucleotide pair - The nucleotides found at a polymorphic site on the two copies of a chromosome from an individual.

5

10

15

· 20

25

30

35

Phased — As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, phased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is known.

Polymorphic site (PS) - A position within a locus at which at least two alternative sequences are found in a population, the most frequent of which has a frequency of no more than 99%.

Polymorphic variant – A gene, mRNA, cDNA, polypeptide or peptide whose nucleotide or amino acid sequence varies from a reference sequence due to the presence of a polymorphism in the gene.

Polymorphism – The sequence variation observed in an individual at a polymorphic site.

Polymorphisms include nucleotide substitutions, insertions, deletions and microsatellites and may, but need not, result in detectable differences in gene expression or protein function.

Polymorphism data — Information concerning one or more of the following for a specific gene: location of polymorphic sites; sequence variation at those sites; frequency of polymorphisms in one or more populations; the different genotypes and/or haplotypes determined for the gene; frequency of one or more of these genotypes and/or haplotypes in one or more populations; any known association(s) between a trait and a genotype or a haplotype for the gene.

**Polymorphism Database** – A collection of polymorphism data arranged in a systematic or methodical way and capable of being individually accessed by electronic or other means.

**Polynucleotide** — A nucleic acid molecule comprised of single-stranded RNA or DNA or comprised of complementary, double-stranded DNA.

Population Group - A group of individuals sharing a common ethnogeographic origin.

Reference Population – A group of subjects or individuals who are predicted to be representative of the genetic variation found in the general population. Typically, the reference population represents the genetic variation in the population at a certainty level of at least 85%, preferably at least 90%, more preferably at least 95% and even more preferably at least 99%.

Single Nucleotide Polymorphism (SNP) – Typically, the specific pair of nucleotides observed at a single polymorphic site. In rare cases, three or four nucleotides may be found.

Subject – A human individual whose genotypes or haplotypes or response to treatment or disease state are to be determined.

Treatment - A stimulus administered internally or externally to a subject.

Unphased — As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, unphased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is not known.

The inventors herein have discovered 38 novel polymorphic sites, and confirmed the existence of 7 other sites, in the ILAR $\alpha$  gene. The polymorphic sites identified by the inventors are referred to as PS1-45 to designate the order in which they are located in the gene (see Table 3 below), with the novel polymorphic site referred to as PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45 and the previously reported polymorphic sites referred to as PS5, PS24, PS26, PS29, PS31, PS42, and PS43.

Thus, in one embodiment, the invention provides an isolated polynucleotide comprising a polymorphic variant of the ILARα gene or a fragment of the gene which contains at least one of the novel polymorphic sites described herein. The nucleotide sequence of a variant ILARα gene is identical to the reference genomic sequence for those portions of the gene examined, as described in the Examples below, except that it comprises a different nucleotide at one or more of the novel polymorphic sites PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45, and may also comprise one or more additional polymorphisms selected from the group consisting of PS5, PS24, PS26, PS29, PS31, PS42, and PS43. Similarly, the nucleotide sequence of a variant fragment of the ILARα gene is identical to the corresponding portion of the reference sequence except for having a different nucleotide at one or more of the novel polymorphic sites described herein. Thus, the invention specifically does not include polynucleotides comprising a nucleotide sequence identical to the reference sequence (or other reported ILARα sequences) or to portions of the reference sequence (or other reported ILARα sequences), except for genotyping oligonucleotides as described below.

The location of a polymorphism in a variant gene or fragment is identified by aligning its sequence against SEQ ID NO:1. The polymorphism is selected from the group consisting of guanine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, cytosine at PS8, thymine at PS9, thymine at PS10, adenine at PS11, adenine at PS12, thymine at PS13, thymine at PS14, adenine at PS15, thymine at PS16, adenine at PS17, thymine at PS18, adenine at PS19, cytosine at PS20, cytosine at PS21, thymine at PS22, cytosine at PS23, thymine at PS25, thymine at PS27, cytosine at PS28, thymine at PS30, adenine at PS32, thymine at PS34, cytosine at PS35, cytosine at PS36, cytosine at PS37, thymine at PS38, guanine at PS39, guanine at PS40, adenine at PS41, thymine at PS44, and adenine at PS45. In a preferred embodiment, the polymorphic variant comprises a naturally-occurring isogene of the ILARα gene which is defined by any one of haplotypes 1-53 shown in Table 5 below.

Polymorphic variants of the invention may be prepared by isolating a clone containing the ILAR $\alpha$  gene from a human genomic library. The clone may be sequenced to determine the identity of the nucleotides at the polymorphic sites described herein. Any particular variant claimed herein could be

prepared from this clone by performing in vitro mutagenesis using procedures well-known in the art.

ILARα isogenes may be isolated using any method that allows separation of the two "copies" of the ILARα gene present in an individual, which, as readily understood by the skilled artisan, may be the same allele or different alleles. Separation methods include targeted *in vivo* cloning (TIVC) in yeast as described in WO 98/01573, U.S. Patent No. 5,866,404, and copending U.S. application Serial No. 08/987,966. Another method, which is described in copending U.S. Application Serial No. 08/987,966, uses an allele specific oligonucleotide in combination with primer extension and exonuclease degradation to generate hemizygous DNA targets. Yet other methods are single molecule dilution (SMD) as described in Ruaño et al., Proc. Natl. Acad. Sci. 87:6296-6300, 1990; and allele specific PCR (Ruaño et al., 17 Nucleic Acids. Res. 8392, 1989; Ruaño et al., 19 Nucleic Acids Res. 6877-6882, 1991; Michalatos-Beloin et al., 24 Nucleic Acids Res. 4841-4843, 1996).

The invention also provides ILARα genome anthologies, which are collections of ILARα isogenes found in a given population. The population may be any group of at least two individuals, including but not limited to a reference population, a population group, a family population, a clinical population, and a same sex population. An ILARα genome anthology may comprise individual ILARα isogenes stored in separate containers such as microtest tubes, separate wells of a microtitre plate and the like. Alternatively, two or more groups of the ILARα isogenes in the anthology may be stored in separate containers. Individual isogenes or groups of isogenes in a genome anthology may be stored in any convenient and stable form, including but not limited to in buffered solutions, as DNA precipitates, freeze-dried preparations and the like. A preferred ILARα genome anthology of the invention comprises a set of isogenes defined by the haplotypes shown in Table 5 below.

An isolated polynucleotide containing a polymorphic variant nucleotide sequence of the invention may be operably linked to one or more expression regulatory elements in a recombinant expression vector capable of being propagated and expressing the encoded ILARα protein in a prokaryotic or a eukaryotic host cell. Examples of expression regulatory elements which may be used include, but are not limited to, the lac system, operator and promoter regions of phage lambda, yeast promoters, and promoters derived from vaccinia virus, adenovirus, retroviruses, or SV40. Other regulatory elements include, but are not limited to, appropriate leader sequences, termination codons, polyadenylation signals, and other sequences required for the appropriate transcription and subsequent translation of the nucleic acid sequence in a given host cell. Of course, the correct combinations of expression regulatory elements will depend on the host system used. In addition, it is understood that the expression vector contains any additional elements necessary for its transfer to and subsequent replication in the host cell. Examples of such elements include, but are not limited to, origins of replication and selectable markers. Such expression vectors are commercially available or are readily constructed using methods known to those in the art (e.g., F. Ausubel et al., 1987, in "Current Protocols in Molecular Biology", John Wiley and Sons, New York, New York). Host cells which may be used to express the variant ILARα sequences of the

inv ntion include, but are not limited to, eukaryotic and mammalian cells, such as animal, plant, insect and yeast cells, and prokaryotic cells, such as E. coli, or algal cells as known in the art. The recombinant expression vector may be introduced into the host cell using any method known to those in the art including, but not limited to, microinjection, electroporation, particle bombardment, transduction, and transfection using DEAE-dextran, lipofection, or calcium phosphate (see e.g., Sambrook et al. (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York). In a preferred aspect, eukaryotic expression vectors that function in eukaryotic cells, and preferably mammalian cells, are used. Non-limiting examples of such vectors include vaccinia virus vectors, adenovirus vectors, herpes virus vectors, and baculovirus transfer vectors. Preferred eukaryotic cell lines include COS cells, CHO cells, HeLa cells, NIH/3T3 cells, and embryonic stem cells (Thomson, J. A. et al., 1998 Science 282:1145-1147). Particularly preferred host cells are mammalian cells.

5

10

15

20

25 ·

30

35

As will be readily recognized by the skilled artisan, expression of polymorphic variants of the IL4Ra gene will produce IL4Ra mRNAs varying from each other at any polymorphic site retained in the spliced and processed mRNA molecules. These mRNAs can be used for the preparation of an IL4Ra cDNA comprising a nucleotide sequence which is a polymorphic variant of the ILAR reference coding sequence shown in Figure 2. Thus, the invention also provides ILARa mRNAs and corresponding cDNAs which comprise a nucleotide sequence that is identical to SEO ID NO:2 (Fig. 2), or its corresponding RNA sequence, except for having one or more polymorphisms selected from the group consisting of thymine at a position corresponding to nucleotide 237, adenine at a position corresponding to nucleotide 244, cytosine at a position corresponding to nucleotide 291, thymine at a position corresponding to nucleotide 501, adenine at a position corresponding to nucleotide 554, cytosine at a position corresponding to nucleotide 939, thymine at a position corresponding to nucleotide 1242, thymine at a position corresponding to nucleotide 1293, cytosine at a position corresponding to nucleotide 1299, thymine at a position corresponding to nucleotide 1701, adenine at a position corresponding to nucleotide 1735, thymine at a position corresponding to nucleotide 2023, guanine at a position corresponding to nucleotide 2254 and cytosine at a position corresponding to nucleotide 2397, and may also comprise one or more additional polymorphisms selected from the group consisting of guanine at a position corresponding to 223, cytosine at a position corresponding to nucleotide 1199, cytosine at a position corresponding to 1291, cytosine at a position corresponding to nucleotide 1507 and guanine at a position corresponding to 1737. Fragments of these variant mRNAs and cDNAs are included in the scope of the invention, provided they contain the novel polymorphisms described herein. The invention specifically excludes polynucleotides identical to previously identified and characterized IL4Ra cDNAs and fragments thereof. Polynucleotides comprising a variant RNA or DNA sequence may be isolated from a biological sample using well-known molecular biological procedures or may be chemically synthesized.

Genomic and cDNA fragments of the invention comprise at least one novel polymorphic site identified herein and have a length of at least 10 nucleotides and may range up to the full length of the

gene. Preferably, a fragment according to the present invention is between 100 and 3000 nucleotides in length, and more preferably between 200 and 2000 nucleotides in length, and most preferably between 500 and 1000 nucleotides in length.

5

10

15

20

25

30

35

In describing the polymorphic sites identified herein, reference is made to the sense strand of the gene for convenience. However, as recognized by the skilled artisan, nucleic acid molecules containing the ILAR $\alpha$  gene may be complementary double stranded molecules and thus reference to a particular site on the sense strand refers as well to the corresponding site on the complementary antisense strand. Thus, reference may be made to the same polymorphic site on either strand and an oligonucleotide may be designed to hybridize specifically to either strand at a target region containing the polymorphic site. Thus, the invention also includes single-stranded polynucleotides which are complementary to the sense strand of the ILAR $\alpha$  genomic variants described herein.

Polynucleotides comprising a polymorphic gene variant or fragment may be useful for therapeutic purposes. For example, where a patient could benefit from expression, or increased expression, of a particular ILAR $\alpha$  protein isoform, an expression vector encoding the isoform may be administered to the patient. The patient may be one who lacks the ILAR $\alpha$  isogene encoding that isoform or may already have at least one copy of that isogene.

In other situations, it may be desirable to decrease or block expression of a particular ILARα isogene. Expression of an ILARα isogene may be turned off by transforming a targeted organ, tissue or cell population with an expression vector that expresses high levels of untranslatable mRNA for the isogene. Alternatively, oligonucleotides directed against the regulatory regions (e.g., promoter, introns, enhancers, 3' untranslated region) of the isogene may block transcription. Oligonucleotides targeting the transcription initiation site, e.g., between positions –10 and +10 from the start site are preferred. Similarly, inhibition of transcription can be achieved using oligonucleotides that base-pair with region(s) of the isogene DNA to form triplex DNA (see e.g., Gee et al. in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, N.Y., 1994). Antisense oligonucleotides may also be designed to block translation of IL4Rα mRNA transcribed from a particular isogene. It is also contemplated that ribozymes may be designed that can catalyze the specific cleavage of IL4Rα mRNA transcribed from a particular isogene.

The oligonucleotides may be delivered to a target cell or tissue by expression from a vector introduced into the cell or tissue *in vivo* or *ex vivo*. Alternatively, the oligonucleotides may be formulated as a pharmaceutical composition for administration to the patient. Oligoribonucleotides and/or oligodeoxynucleotides intended for use as antisense oligonucleotides may be modified to increase stability and half-life. Possible modifications include, but are not limited to phosphorothioate or 2' Omethyl linkages, and the inclusion of nontraditional bases such as inosine and queosine, as well as acetyl, methyl-, thio-, and similarly modified forms of adenine, cytosine, guanine, thymine, and uracil which are not as easily recognized by endogenous nucleases.

The invention also provides an isolated polypeptide comprising a polymorphic variant of the reference IL4Rα amino acid sequence shown in 3. The location of a variant amino acid in an IL4Rα polypeptide or fragment of the invention is identified by aligning its sequence against Fig. 3. An ILARa protein variant of the invention comprises an amino acid sequence identical to SEQ ID NO: 3 except for having one or more variant amino acids selected from the group consisting of threonine at a position corresponding to amino acid 82, histidine at a position corresponding to amino acid 185, isoleucine at a position corresponding to amino acid 579, serine at a position corresponding to amino acid 675, and alanine at a position corresponding to amino acid 752, and may also comprise one or more additional variant amino acids selected from the group consisting of valine at a position corresponding to amino acid 75, alanine at a position corresponding to amino acid 400, arginine at a position corresponding to amino acid 431, proline at a position corresponding to amino acid 503, and arginine at a position corresponding to amino acid 576. The invention specifically excludes amino acid sequences identical to those previously identified for ILARα, including SEQ ID NO: 3, and previously described fragments thereof. ILARα protein variants included within the invention comprise all amino acid sequences based on SEQ ID NO: 3 and having the combination of amino acid variations described in Table 2 below. In preferred embodiments, an ILARa protein variant of the invention is encoded by an isogene defined by one of the observed haplotypes shown in Table 5.

20

15

5

10

25

30

35

B. (		Table	2. Nove	l Polymo	rphic Va	riant of IL	.4Rα			
Polymorphic Variant				Amina /	Naid Dani	tion and	ld stitle	_		
Number	75	82	185	400	431	tion and	576	579	675	752
1 1	13	A	R R	E	C -	S S	0 Q	V	P P	732 A
2		A	R	E	<del>  č</del>	S	a	V	S	S
3	1		R	Ē	c	S	a	V	S	
		A								<u>A</u>
4		Α	R.	E	C	S	Q		Р	S
5		Α	R	E	C	S	Q	!	Р	<u>A</u>
6	!	<u>A</u>	R	E	C	S	Q	<u> </u>	S	S
7	l l	Α	R	E	С	S	Q		S	<u>A</u>
8		Α_	R	E	С	S	R	>	P	A
. 9	1	Α	R	E	С	·S	R	>	S	S
10	<u> </u>	A	R	E	С	S	R	<b>&gt;</b>	S	A
. 11	. 1 .	Α	R	E	С	S	R	1 .	Р	S
12	-	Α	R	E	С	S	R	1	P	Α
13	l	Α	R	E	С	S	R		S	S
14		Α	R	E.	С	Ş.	R.	_	S	Α
15	1	Α	R	E	С	Р	Q	>	Р	Α
16		Α	R	E ·	С	Р	Q	V	S	S
17	1	Α	R	E	С	Р	Q	V ·	·s	Α
18	i	Α	R	E	С	Р	Q		Р	S
19	ī	Α	R	Е	С	Р	Q	ı	Р	Α
20		Α	R	Е	С	Р	· Q	1	S	S
21	. ]	A	R	E	С	Р	Q		S	Α
22	$\overline{}$	Α	R	E	C	P	R	V	Р	Α
23	i	A	R	E	Ċ	Р	R	V	S	S
24	i	A	R	E	Č	Р	R	V	S	A
25 .	1	A	R	Ē	č	P	R	1	P	S
26	i	A	R	Ē	C	P	R	i	Р	A
27		·A	R	Ē	· Č	P	R	<del>i i</del>	S	S
28	1	A	R	E	Č	P	R	i	S	Ā
29	<u>i</u>	Ä	R	Ē	R	S	Q -	V	P	Â
30	<u> </u>	Â	R	Ē	R	S	Q	v	S	S
31	<del></del>	A	R	Ē	R	S	Q	V	S	A
			R	E	R	S	à		P	S
32		A								
33		A	R	E	R	S	Q		P (	<u>A</u>
34	1	Α	R	E	R	S	Q	!	S	S
35	<u> </u>	A	R	E	R	S	Q	1	S	· A
36		Α	R	E	R	S	R	<b>V</b>	Р	A
37	!	Α	R	E	R	S	R	V	S	S
38	<u></u>	Α	R	E	R	S	R	V	S	A
39		Α	R	E	R	S	R		Р	S
40		Α	R	E	R	·S	· R	<u> </u>	Р	Α
41	1	Α	R	E	R	S	R		S	S
42	1	Α	R	E	R	S	R		S	Α
43		Α	R	E	R.	Р	Q	V	Р	Α
44		Α	R	Е	R	Þ	q	V	S	S
45	l l	Α	R	E	R	P	Q	٧	S	Α
46	I	Α	R	E	R	Р	Q	J	Р	S
47	ı	Α	R	E	R	Р	Q		Р	Α
48	1	Α	R	E	R	Р	Q	T	S	S
49	1	A	R	Ē	R	P	Q		S	Ā
50 .	i	Ā	R	Ē	R	P	R	V	P	A
51	<del></del> i	A	R	. E	R	P	R	V	S	S
52	<del>-                                    </del>	A	R	E	R	P	R	V	S	Ā
53		A	<u> </u>	Ē	R	P				

	7	Table2. N	lovel Po	lymorphi	c Variant	of IL4R	α (cont'd	)		
Polymorphic	<u> </u>							<u></u>		
Variant				Amino A	Acid Posi	tion and	Identities	3		
Number	75	82	185	400	431	503	576	579	675	752
54	·	Α	R	E	R	Р	R	I	Р	Α
55 ·	_	. A	R	E	R	Р	R		S	S
- 56	1.	Α	R	E	R	Р	R	_ 1	S	Α
57	ı	Α	R	Α	С	S	Q	V	Р	Α
58	ı ı	Α	R	Α	С	S	Q	V	S	S
59	I	Α	R	Α	С	S	Q	٧.	S	Α
60	Ī	Α	R	Α	С	S	Q	ı	Р	S
61		Ä	R	Α	С	S	Q		Р	Α
62		Α	R	Α	С	S	Q		S	S
63	_	Α	R	Α	С	S	Q	1.	S	·A
64	· 1	Α	R	Α	С	S	R	V	Р	Α
65	ı	· , A	R	Α	С	S	R	V	S	S
66	ı	Α	R	Α .	С	S	R.	V	S	Α
67	ı	Α	R	Α	С	S	R	1	Р	S
68	i	Α	R	Α	С	S	R		P	Α
69	ı	Α	R	Α	С	S	R	1	S	S
70	1	Α	R	Α	C	S	R		S	Α
71	I	Α	R	Α	С	Р	Q	>	Р	Α
72 ·		Α	R	. A	С	Р	Q.	V	S	S
73	1	Α	R	Α·	С	Р	Q	V	S	Α
74	ı	Α	٠R	Α	С	Р	Q	1	Р	S
75		Α	R	Α	С	Р	Q		Р	Α
76	ı	Α	R ·	Α	С	Р	Q	ı	S	S
. 77	ı	Α	R	Α	С	Р	Q		S	Α
78	T T	Α.	R	Α	С	Р	R	V.	Р	Α
79	ı	Α	R	Α	С	Р	R	٧	S	S
80		Α	R	Α	С	Р	R	V	S	Α
81	1	Α	R	Α	С	Р	R	ł	Р	S
82		Α	R	Α	С	. Р	R	1	Р	A
83	I	Α	R	Α	С	P	R	1 "	S	S
84	I	Α	R	Α	С	Р	R	1	S	Α
85	1	Α	R	A .	R	S-	Q	V	Р	Α
86		Α	R	Α	R	· S	Q	٧	S	S
87	1	Α	R	Α	R	S	Q	V	S	Α
88	l	Α	R	Α	R	S	Q		Р	S
89	ı ı	Α	R	Α	R	S	Q	ı	Р	Α
90		Α	R	Α	R	S	a	ı	S	S
91	1	Α	R	Α	R	S	Q	ı	S	Α
92		Α	R	Α	R	S	R	V	Р	Α
93		Α	R	Α	R	S	R	V	S	S
94	1	Α	R	Α	R	S	R	V	S	Α
95		Α	R	Α	R	S	R	1	Р	S
96		- A	R	Α	R	S	R	1	Р	Α
97	11	Α	R	Α	R	S	R		S	\$
98		Α	R	Α	R	S	R		S	Α
99		Α	R	Α	R	Р	Q ·	. V	Р	Α
100	ŀ	Α	R	Α	R	Р	α	_ V	S	S
101		Α	R	Α	R	Р	Q	٧	S	Α
102		Α	R	Α	R	P.	Q		P	S
103		Α	R	Α	R	P	Q	T	P	Α
104		Α	R	Α	R	Р	Q	1	S	S
105		Α	R	Α	R	Р	Q	1	S	Α
106		A	R	A	R	Р	R	V	P	Α
							·			

Polymorphio		Table2. N	lovel Po	ymorphi	c Varian	of IL4R	α (cont'd	)		
Polymorphic				A: /	haid Daai	:	Idontitio			
Variant	75	- 02	105	400		ition and			675	750
Number	75	82	185		431	503 P	576	579 V	675	752 S
107	<u> </u>	A	R	A	R		R	<del>                                     </del>	S	
108	1	A	R	A	R	Р	R			A
109	!	A	R	A	R	P	R	<del>-                                    </del>	P	S
.110	!	A	R	A	R	Р	R	<u> </u>	Р	A
111	<u> </u>	A	R	Α	R	Р	R	<u> </u>	S	S
112	1.	Α	R	A	R	Р	R	<del> ;</del>	S	A
113	<u> </u>	Α	Н	E	С	S	Q	<u> </u>	Р	S
114	1	Α	Н	E	С	S	Q	V	Р	Α
115		Α	Н	E	C	S	Q	V	S	S
116		Α	Н	E	С	S	Q	V	S	Α
117	<u> </u>	Α	Н	E	С	S	Q		Р	S
118	<u> </u>	Α	I	E	С	S	Q		Р	Α
119		Α	H	E	С	S	Q		S	S
120	1	Α	Н	E	С	S	Q		S	Α
. 121	1	Α	H	E	С	S	R	V	Р	S
122		Α	H	E	С	S	.R	V	Р	Α
123	Ī	Α	Н	Е	С	S	R	٧	S	S
124	ı	Α	Н	E	С	S	R	٧	S	Α
125	ı I	Α	H	E	С	S	R		Р	S
126	ı	Α	Н	E	С	S	R	ı	·P	Α
127		Α	H	E	С	S.	R		S	S
128		Α	Н	E	С	S	R		S	Α
129	ı	Α	Н	E	С	Р	Q	٧	Р	S
130	ı	Α	Н	E	С	Р	Q	V	Р	Α
131	T	Ä	Н	E	С	Р	Q	V	S	S
132		Α	Н	E	C ·	P.	Q	V	S	Α
133		Α	Н	E	С	Р	Q	i	Р	S
134		A	Н	E	С	Р	Q		Р	A
135	<u> </u>	A	H	E	c	P	Q		S	S
136		A	Н	E	Ċ	Р	ä		S	A
137	<u> </u>	A	Н	E	Ċ	Р	R	V	P	S
138	<u> </u>	A	Н	E	Č	Р	R	V	Р	A
139	<u> </u>	A.	Н	E	č	P	R	V	S	S
140	<u> </u>	A	H	Ē	Č	P	R	V	Š	· A
141	<del></del>	Ā	H	Ē	č	Р	R	i	P	S
142	<del>l i ·</del>	A	Н	Ē	Č	P	R	· i .	P	Ā
143	i	Â	H	Ē	č	P	R	<del>- i -</del>	S	ŝ
144	<del></del>	Â	H	E	C	P	R	<del>i i</del>	S	Ā
145	<del>                                     </del>	Ä	- <del>''</del>	F	R	s	Q	V	P	s
146	<del>-                                    </del>	A	Н	E	R	S	Q	v	P	A
147	<del></del>	Â	H	E	Ŗ	S	q	V	S	S
148	<del>i i</del>	Â	H	E	R	S	à	V	S	·A
149	<del> </del>	Ā	H	E	R	S	Q	<u> </u>	P	- <del>S</del>
150	<del></del>		H	E	R	S			P	
	<del></del>	A				S	Q		S	A
151		A	H	E	R		Q	1		S
152	<del>                                     </del>	A	H	E	R	S	Q	1	S	A
153	- !	A	H	E	R	S	R	<b>&gt;</b>	P	S
154	<u> </u>	Α	Н	E	R	S	R	٧	Р	A
155		A	Н	E	R	S	R	V	S	S
156		A	Н	E	R	S	R	V	S	Α
157		Α	H	E ·	R	S	R		Р	S
158		Α	Н	E	R	S	R		Р	Α
159		Α	Н	ш	R	S	R		S	S

	7	able2. N	lovel Pol	lymorphi	c Variant	of IL4R	x (cont'd)	)		
Polymorphic										
Variant				Amino A		tion and	Identities			
Number	75	82	185	400	431	503	576	579	675	752
_160		Α	Н	E	R	S	R	1	S	Α
161	ı	Α	Н	E	R	Р	Q	V	Р	S
162		Α	H	E	R	Р	a	٧	Р	Α
163		Α	H.	E	R	P	Q	٧	S	S
164	I	Α.	Н	E	R	Р	Q	V	S	Α
165	Ī	Α	Н	E	R	Р	Q	1	P	S
166		Α	.H	E	R	Р	Q		Р	Α
167	l	Α	Н	E	R	Р	Q	1	S	S
168	Ī	Α	H	E	R	Р	Q	· 1	S	Α
169		Α	H	E	R	Р	R	V	Р	S
170	ı	Α	Н	E	R	Р	R	V	Р	Α
171.	1	Α	Н	E	R	Р	R	V	S	S
172	7.0	Α	Н	E	R	Р	R	V	S	Α
173		Α	Н	E	R	Р	R	ı	Р	S
174	1	Α	Н	E	R	Р	R	• 1	Р	Α
175		Α	Н	E	R	Р	R	T I	S	S
176		Α	Н	E	R	Р	R		Š	A
177		Α	H	A	С	S	Q	·V	P	S
178	i	Α	Н	Α	Č	S	Q	v	P	Ā
179	<del>_</del>	A	H	A	Č	s	Q	v	s	S
180	i	A	H	Ā	č	s	Q	V	Š	Ā
181	<u> </u>	A	H	A	c	S	Q	Ť	P	S
182		A	H .	A	C	S	Q	<del></del>	P	Ā
183		A	Н	A	Č	S	Q	j	S	S
184		A	H	Ā	C	S	·Q	<del>i</del>	S	Ā
185		A	H	Â	c	S	R	<del>\</del>	P	S
186	<del>-</del>	Â	H	Â	c	S	R	v	·Р	Ā
187	<del>- i -</del>	A	H	A	c	S	R	V	S	s
188	-	A	H	A	c	S	R	V	Š	Ā
189		A	Н	A	C	S	R	<del>i</del>	P	S
190	<u> </u>	Â	H	A	C	S	R	- i	P.	Ā
191	1	A	H	A	č	S	R		S	S
192		A	H	A	<del>  č</del>	. S	R	1.	· S	A
193	-	A.	H	Â	c	P	Q	V	P	S
194		A	H	Â	c	Р	à	· v	P	A
195	<u> </u>	A	H	A	c	P	à	v	s	s
196		A	H	A	<del>C</del>	P	ď	V	S	A
197		A	H	Â	c	Р	Q	<del></del>	P	S
198	1	A	H	A	c	P	Q		P	A
199	<del></del>	A	H	A	c	P	a	<del></del>	S	S
200	<del>'</del> -	A	H	A	c	P	Q	<del>  </del>	S	A
200		A	<del>                                     </del>		c	P	·R	<del>- '</del>	P	S
201		A	H	A	<del>  c</del>	P	R	V	P	A
				A	C.	P		- <del>v</del> -	S	S
203		A	Н	A			·R	V		
204	<u> </u>	A	H	A	C	Р	R	<del></del>	S	A
205		A	H_	Α	C	Р	R		P	_ S
206	!_	Α	Н	· A	C	Р	·R			A
207	- !	Α	Н	A	C	.P	R		S	S
208	!	Α	Н	Α	C	Р	R	!	S	A
209	!	Α	Н	A	R	S	Q	V	Р	S
210		Α	Н	Α	R	S	Q	V	Р	A
211		Α	Н	Α	R	S	Q	V	S	S
212		Α	. Н	Α	R	S	Q	V	S	Α

	7	Table2. N	lovel Pol	ymorphi	c Variant	of IL4R	x (cont'd	)		-
Polymorphic				A i /		4:d		_		_
Variant Number	75	82	185	400	431	tion and 503	576	579	675	752
. 213		A	Н	400 A	R R	S S	Q Q	1	P	/32   S
214		A	H		R	S		-	F	A
215	<del></del>	A	H	A	R	S	Q	<del>-                                    </del>	S	S
216		A	H	Â	R	S	l ä	- ;	S	Ā
217	<del></del>	A	H	A	R	S	R	l 😾	P.	S
218		A	H	A	R	S	R	l v	P	A
219	<u>'</u>	A	Н	Â	R	S	R	V	ร	ŝ
220		Â	H	A	R	S	R	V	S	A
221	i	Â	H	Â	R	S	R	Ť	P	S
222	1	Â	H	Â	R	S	R.	<del>l i</del>	P	Ā
223	<u>i</u>	Â	H	Â	R	S	R	<del> </del>	s	ŝ
224	-	Â	H	Â	R	S	R	<del>                                     </del>	S	Ä
225	1	A	H	Â	R	P	à	V	P	ŝ
226 .	<del>- i-</del>	A	Н	A	R	P	à	V	P	A
227	- i	A	Н	A	R	P	<del>\</del>	V	S	S
228	i	A	Н	Â	R	P	ă	V	·s	A
229	<u> </u>	Â	Н	Â	R	P	ă	l i	P	ŝ
230	<u> </u>	Â	Н	Ä	R	P	ă	<del> </del>	P	A
231		A	Н	A	R	P	à	<del>l i</del>	S	S
232	-i-	Ā	H	A	R	P	à	<del></del>	s	A
233	Ť	Ā	Н	A	R	Р	Ř	Ι <del>ν</del>	P	S
234	i	A	H	Ā	R	Р	R	V	P	A
235		A	Н	A	R	Р	R	V	S	s
236	i	A	Н	A	R	Р	R	V	s	Ā
237		Α	Н	A	R	P	R	T	P	S
238	1	Α	Н	Α	R	Р	R	1	Р	Α
239		Α	Н	Α	R	Р	R	1	S	S
240	· I	Α	Н	Α	R	P·	R		S	A
241		Т	R	Е	С	S	Q	·V	Р	S
242	ı	T	R	E	С	S	Q	V	·P	Α
243	ı	T	R	E	С	·S	Q	V ·	S	S
244	ı	T	R	Е	С	S	Q	V	S	Α
245	1	T	R	E	С	S	Q		Р	S
246		T	R	E	С	S	Q	1	Р	Α
247	1	T	R	Е	C	S	a	_	S	S
248		T	R	E	С	S	Ø	_	S	Α
249	1	T	R	E	С	. S	R	<b>V</b>	Р	S
250	1	T	R .	E	С	S	R	<b>V</b>	_ P	Α
251	1	T	R	E	С	S	R	>	S	S
252		T	R	E	С	S	R	<b>&gt;</b>	S	Α
253		T	R	E	С	S	R		Р	S
254		T	R	E	C	S	R		Р	Α
255	1	Ţ	R	E	С	S	R		S	S
256		T	R	E	C	S	R		S	A
257	ı		R	E	C	P	Q	V	Р	S
258		<u>T</u>	R	<u>E</u>	<u>C</u> .	Р	Q	V	Р	A
259	!	T	R	E	ပ	Р	Q	V	S	S
260	<u> </u>	Ţ	R	E	ပ	Р	· Q	V	S	A
261		Ţ	R	E	C	Р	Q	· !	Р	S
262		<u></u>	R	<u> </u>	C	Р	Q	!	Р	A
263		Ţ	R	E	C	P	Q	l	S	S
264	1	. T	R	Ę_	C	P	Q	1	S	A
265	l	T	R	Е	С	Р	R	V	Р	S

·		Γable2. N	lovel Pol	lymorphi	c Variant	of IL4R	x (cont'd	)		
Polymorphic								•		
Variant	·					tion and				r-==-
Number	75 .	82	185	400	431	503	576	579	675	752
266	<u> </u>	T	R	E	C	Р	R	·V	Р	A
267	!	T	R	E	C	P	R	V	S	S
268		Τ.	R	E	С	Р	R	V	S	A
269	<u> </u>	<u></u>	R	E	С	Р	R	!	Р	S
270	<u> </u>	T	R	E	C	Р	R	!	Р	<u>A</u>
271	!	Ţ	R	E	C	P	R		S	S
272		Ţ	R	E	C	P	R	1	S	A
273		T	R	E	R	S	Q	V	Р	S
274		T	R	E	R	S	Q	V	Р	A
275	- !	T	R	E	R_	S	Q	V	S	S
276		T	R	E	R	S	Q	V	S	Α
277	- !	<u>T</u>	R.	E	R	S	α		Р	S
278		Ţ	R	E	R	S	Q		P	A
279	1	Ţ	R	E	R	S	α	1	S	S
280	1	T	R	E	R	S	Q	ı	S	A
281		T	R	E	R	S	R	> >	P	S
282	1	T	R	E	R	S	R	<b>&gt;</b>	Р	A
283	-!	T	R	E	R	S	R	V	S	S
. 284	_ !	T.	R	E	R	S	R	<b>&gt;</b>	S	A
285	<u> </u>	Ţ	R	E	R	S	R		Р	s
286 -		T	R	E	R	S	R.		P	A
287	. !	T	R	E	R	S	R <sub>z</sub>	!	S	S
288	_	T	R	E	R	S	R	1	S	A·
289		T	R	E	R	Р	Q	<u> </u>	Р	S
290		T	R	E	R	Р	Q	<b>&gt;</b>	Р	A
291		T	R	E	R	P	Q	<b>&gt;</b>	S	S
292	1	T	R	E	R	P	Q	· V	S	· A
293		Ţ	R	E	R		Q	- !	Р	S
294	1	T	R	E	R	Р	Q	1.	Р	A S
230			R	E	·R	P.	Q	_	S	
296 297		T	R	E	R	P	Q R	V	S	A
		<u></u>			R	. Р		<b>V</b>	P	<u> </u>
298 299	1		· R	E	R	. P	R	V		A S
300		T	R	듵		<u>.</u> Р	R	- <del>V</del>	S	
301		<del>'</del>	R	Ē	R	P	R	<u> </u>	P	S
302	-	Ť	R	<u> </u>	R	P	R		P	A
303		<del></del>	R	<u> </u>	R	P	R		S	S
303		<del></del>	·R	Ē	R	P	R	<del>  </del>	S	A
305	<del>                                     </del>	Ť	R	_ <del>_</del> _	C	S	Q	V	P	S
315	<del></del>	Ť	R	A	č	S	R	V	S	S
316		Ť	R	Â	l č	S	R	V	S	A
317		<del>'</del>	R	A	c	S	R	1	P	S
318	1	<del></del>	R	A	<del>  c</del>	S ·	R		P	A
319	<del></del>	<del></del>	R	A	C	S	R	1	S	S
320	<del>-                                    </del>	<del>'</del>	R		c	S	R		S.	
321	1	Ť	R	A	c	P	Q	V	<u>э.</u> Р	· A
		+-			C			V		S
322		1	R	A		Р	σc		Р	A
323	1		R	A	C	P	α		S	S
324	1	Ī	R	A	C	P	Q	-	S	A_
325		T	R	A	C	Р	Q		P	S
326	!	T	R	A	C	P	Q	<u>.</u>	P	A
327		T	R	Α	С	Р	α		S	S

Polymorphic	1	Table2. N	lovel Po	lymorphi	c Variant	of IL4Re	x (cont'd	)	<del></del>	
Variant				Amino A	aid Dasi	ition and	Idontitio	•		
Number	75	82	185	400	431	503	576	579	675	752
328	- 75	T	183 R	A	C	P	0 Q	3/9	S S	
306								$-\frac{1}{V}$	P	A A
307		T	R	A	C	S	Q	l v		A
		Ţ	R:	A	C	S	Q		S	S
308		T	R	A	C	S	Q.	V	S	Α
309		T·	R	Α	С	S	Q		Р	S
310		T	R	A	С	S	Q		Р	Α
311		T	R	Α	С	S	Q	ı	S	S
312	1	T	R	Α	С	S	Q	l	S	Α
313		T	R	Α	С	S	R	٧	Р	S
314	ı	T	R	Α	С	S	R	V	P	·A
329		T	R	Α	С	Р	R	V	Р	S
330		T	R	Α	C	Р	R	V	Р	Ā
331	i	T	R	A	Č	P	R	Ÿ	s	S
332	i	Ť	R	Â	c	Р	R	V	S	A
333	<del>-</del>	Ť	R	Â	c	P	R	<del></del>	P	S
334		<del></del>	R	A	<del>c</del>	P	R	<del>- '</del>	P	A
335		T	R	A	C	Р	R		S	S
336	!	T	R	A	C	Р	R		S	<u>A</u>
337		T	R	Α	R	S	Q	V	Р	S
338	١. ا	Т	R	Α ·	R	S	Q	V	Р	Α
339	. 1	T	R ·	Α	R.	S	Q	V	S	S
340	1	Т	R	A	R	S	Q	V	S	Α
341		T	R	Α	R	S	Q	1	Ρ	S
342	1.	Т	R	Α	R	S	Q		Р	Α
343	1.	Т	R	Α	R	S	Q	1	S	S
344		T	R	Α	R.	S	Q	i	S	Ā
345	i	T	R	A	R	S	R	V	P.	S
346	i	Ť	R	Â	R	Š	R	v	P	Ā
347	i	Ť	R	A	R	S	R	v	S	S
348		Ť	R	A	R	S	R	v	S	A
349		+	R	A		S		<u> </u>	P	
					R		R			S
. 350	!	<u>T</u>	R	A	R	S	R		Р	A
351		T	R	A	R	S	R		S	S
352	·	T	R	Α	R	S	R	l ,	S	A
353	ı	T	R	A	R	P	Q	V	Ρ	S
354	1	T	R	Α	R	Р	Q	V	· P	Α
355	1	T	R	Α	R	Р	Q	V	S	S
356	ı	Т	R	Α	R	Р	Q	٧	S	Α
357	T	T	R	Α	R	P.	Q		Р	S
- 358		T	R	Α	R	Р	Q		Р	Ā
359	i	T	R	Α	R	P	ā	i	S	S
360	i	<del></del>	R	Ā	R	P	a	i	S	A
361	<del>-                                    </del>	Ť	·R	Ä	R	P	R.	V	P	S
362	<u> </u>	Ť	R	Â	R	P	· R	V	P	A
363		T	R	A	R	P	R	V	S	S
	- ! -									
364		T	R	Α	R	P	R	<b>\</b>	S	A
365	ı	T	R	Α	R	Р	R		Р	S
366		T	R	Α	R	Р	R	1	Р	Α
367	1	T	R	Α	R	Р	R		S_	S
368	1	T	R	Α	R	P	R		S	A
369	1	T	Н	E	С	S	Q	V	· P	S
370	• ]	T	Н	E	C	S	Q	V	Р	A
371	<del>-</del>	T	H	Ē	Č	s	à	v	S	S

		Table2. N	lovel Po	lymorphi	c Variant	of IL4R	x (cont'd	) .		
Polymorphic Variant				Amino A	Acid Posi	tion and	Identities	8		
Number	75	82	185	400	431	503	576	579	675	752
372	1	T	Н	E	С	S	Q	V	S	Α
373	T	T	Н	E	С	S	Q		Р	S
374		T	Н	E	С	S	Q	i i	Р	Α
375	1	Ť	Н	E	С	S .	Q	1	s	S
376		T	Н	E	С	S	. Q	-1	S	A
377		T	Н	E	C	S	R	V	P	S
378		T	Н	E	C	S	R	V	·Р	A
379		T	Н	E	С	S	R	V	S	S
380		T	Н	E	C	S	R	V	S	A
381		T	Н	E	C	S	R		P	S
382	<del></del>	Ť	H	E	c	Ŝ.	R	i	Р	Ā
383	<del>- i -</del>	Ť	Н	E	č	s	R	i	S	S
384	i	Ť	H	Ē	Ċ	S	R	i	Š	Ā
385	—i	<del>-                                    </del>	H	Ē	c	P	à	v	Р	S
386	<del></del>	<del></del>	H	Ē	Č	P.	à	V	P	Ä
387	·i	Ť	H	Ē	C	P	a	V	S	S
388	<del>- i-</del>	Ť	H	Ē	C	P	a	V	S	A
389	— i	Ť	Н	Ē	Č	P	Q	Ì	P	S
390	i	Ť	H	Ē	Č	P	Q	<del></del>	P	l ă
391	<del></del>	Ť	H	E	C	P	• Q	<del>- i -</del>	S	S
. 392	<del>i</del>	Ť	H	Ē	C	P	Q	<del></del>	s	Ā
393	<u>'</u>	<del>'</del>	H	Ē	c	P	R	V	P	S
394	· · · · ·	Ť	H	E	C	P	R	V	· P	Ä
395	i	Ť	H	Ē	C	P	R	V	S	S
396	i	÷	H	E	C	P	R	V	S	A
397	1	Ť	H	Ë	č	P	R	1	P	ŝ
398		Ť	H	Ē	c	P	R		P	A
399	-	Ť	·H	Ē	č	P	R	i	S	S
400		Ť	H	Ē	C	P	R	<del>-                                    </del>	S	A
401	- ; -	Ť	· H	E	Ř	S	Q.	V	P	S
402		+	H	E	R	S	a	V	P	A
403	<u>'</u>	Ť	H	Ē	R	S	a	V	S	S
404		Ť	H	늗	R	S	à	V	S	A
405		Ť	Н	E	R	S	a	V	P	S
406		+	H	E	R	8	Q	1 .	P	
407		T	H	E	R	S				A
408		<del>-</del>	Н	E	R	S	Q		S	S
409	<del> </del>	Ť	Н	E	R	S	R	V	P	S
410	<del>-                                    </del>	T	Н	E	R	<u> </u>	R	V	P	A
411		+	Н	E	·R			V		
411		<del></del>	Н	E	R	S	R	- <del>V</del> -	S	S
412	<del></del>	<del>- </del>	Н			S		<u>v</u>	P	A
				E	R		R			S
414	<u> </u>	Ţ	Н	E	R	S	R	1	Р	A
415	<del></del>	Ţ	Н	Ш	R	8	R		S	S
416		T	H	E	R	S	R	1	S	A
417			Н	ш	R	P	a	<b>V</b>	Р	S
418	1	T	Н	E	R	. P	Q	V	Р	A
419			Н	E	R	Р	Q	V	S	S
420		T	Н	E	R	Р	Q	V	S	Α
421	!	Ī	Н	E	R	P	Q		Р	S
422		T	I	E	R	Р	Q	1	Р	Α
423		ा	I	E	R	Р	Q	1	S	S
424		T	I	E	R	Р	Q	1	S	Α

1	٦	Table2. N	lovel Po	lymorphi	c Variant	of IL4R	α (cont'd	)		
Polymorphic				,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,						
Variant				Amino A	Acid Posi	ition and	Identities	8		À
Number	75	82	185	400	431	503	576	579	675	752
425		T	H =	E	R	Р	R	V	Р	S
426		<u> </u>	H	E	R	Р	R ·	V	Р	Α
427		Ţ	Н	E	R	Р	R	V	S	S
428	!	T	Н	E	R	Р	R	. V .	S	Α
429		T	Н	Ε	R	Р	R		Р	S
430		T	Н	E	R	Р	R	<u> </u>	Р	A
431 ·		T	Н	E	R,	Р	R	<u> </u>	S	S
432		T	H	E	R	Р	. R	1	S	Α
433		T.	H	A	C	S	Q	V	P	S
434 435		. T .	Н	A	C	S	Ğ	V	Р	A
			Н	A	C	S	Q	V	S	S
436			H	A	C	S	Q	V	S	A
437	<u> </u>	T	H	A	C	S	Q		P	S
438 · 439		T	H	A	C	S	Q		P	A
440	1	+	H	A	C	S			S	S
440	1	<del>-</del>	- <del>H</del> -		C	S	Q	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	P ·	A
442	- 1	÷	H	A	C	S	R R	V	P	S
443	-	÷	H	A	c	S	R	V	S	S.
- 444	-i-	Ť	H	Â	c	S	R	V	S	A
445	- i	Ť	H	A	· C	S	R		P	ŝ
446	<del>- i -</del>	Ť	H	A	C	S	R		P	A
447	<del></del> -	Ť	H	A	C	S	R	<del></del>	S	ŝ
448	<del>- i</del>	÷	Н	Ä	C	S	R	<u> </u>	S	Ā
449	<del>- i.</del>	Ť	H	A	Č	P	Q	·	P	S
450	i	Ť	H	A	C	P	Q	v	P	Ā
451	ì	Ť	Н	A	c	P	à	V	· S	S
452	i	Ť	Н	A	č	P	Q	V	S	A
453		T	Н	A	Ċ	Р	Q	i	P	S
454	i	T	Н	Α	C	Р	à	<u> </u>	P	Ā
455	i	Т	Н	Α	С	Р	Q		S	S
456	ī	T	Н	Α	С	Р	Q		S	A
457	T I	Т	Н	Α	С	Р	R	V	Р	S
458	ı	T	Н	Α	С	Р	R	V	Р	Α
459	T T	T	Н	·A	C	Р	R	V	S	S
460		T	Н	Α	С	Р	R	V	S	Α
461	-	Τ	Н	. A	С	Р	R	ı	P	S
462	ı	Τ	Н	Α	С	Р	R	1	P	Α
463		T	H	Α	С	Р	R		. S	S
464	_	Т	Н	Α	С	Р	R	I	S	Α
465 .	1	T	I	Α	R	S	Q	V	Р	S
466		T	Н	Α	R	S	Q	٧	Р	. A
467	1	T	Н	Α	R	S	Q	V	S	S
468		T	H	Α	R	S	Q	V	S	Α
469		<u> </u>	Н	Α	R	S	Q		Р	S
470			Н	Α	R	S	Q		Р	Α.
471		T	Н	<u>A</u>	R	S	Q		S	S
472		Ţ	H	Α	R	S	Q		S	Α
473	!	T	H	Α	R	S	R	V	Р	S
474.	!	T	Н	Α	R	S	R	V	Ρ.	Α.
475	!	T	Н	A	R	S	R	V	S	S
476	!	T	Н	A	R	S	R	V	S	A
477		Т	Н	Α	R	S	R		P	S

olymorphic		able2. N	lovel Po	lymorphi	c Variant	of IL4R	z (cont'd	)		
Variant				Amino A	laid Dani	tion and	Identities			
	75	82	185	400			576	579	675	752
Number	75	OZ T			431	503		3/9		
478	!		H	A	R	S.	R	<u> </u>	P	A
479	!	<u>T</u>	Н	Α	R	S	R		S	S
480		T	Н	Α	R	S	R		S	Α
481		T	H	Α	R	Р	Q	>	Р	· S
482	ı	T	Н	Α	R	Р	Q	V	Р	Α
483		T	Н	Α	R	Р	Q	V	S	S
484		T	Н	Α	R	Р	Q	V	S	A
485	1	T	Н	Α	R	Р	Q	1	Р	S
486		T	H	Α	R	Р	Q		Р	A
. 487	1	Ť	H	A	R	P	- Q	<del></del>	s	·s
488	·	Ť	H	Ā	R	P	Q	<del></del>	S	A
489		Ť	H	Â	R	P	R	v	P	S
		+						V	<del></del>	
490	!		H	Α	R	P	R			A
491	!	Ţ	Н	Α.	R	Р	R	V	S	S
492	1	T	H	Α	R	Р	R	V	S	A
493	1	T	Н	Α	R	Р	R		Р	·S
494	1	T	Н	٠A	R	Р	R	.	P	Α
495	1	T	H	Α	R	Р	R		S	S
496	ı	T	Н	Α	R	P	R		S	Α
497	V	Α	R	E	С	S	Q	٧	Р	Α
498	V	Α	R	E	С	\$	· Q	V	S	S
499	V	Α	R	E	С	S	Q	V	S	A
500	V	Α	R	E	Ċ	Š	·Q		P	S
501	v	A	R	Ē	C	S	Q	<del></del>	· P	Ä
502	v	A	R	Ē	C	S	à	<del></del>	s	S
503	V	A	R	Ē	C	S	à		s	A
	V					S		V	P	
504		Α_	R	E	O		R			A
505	V	Α	R	E	С	S	R	, V	S	S
506	V	Α	R	Ε.	С	S	R	V	S	Α
507	V	Α	R	E	ပ	S	R		Р	S
508	٧	Α	R	E	C	S	R		Р	Α
509	ν	Α	R	E	C	S	R	- 1	S	S
510	V	·A	R	E	С	S	R	]	S	Α
511	٧	Α	R	Ε	C	Р	Q	٧	Р	Α
512	٧	Α	R	E	С	Р	Q	٧	S	S
513	V	Α	R	E	С	Р	Q	V	S	A
514	V	Α	R	E	Č	Р	Q	1	Р	S
515	v	A	R	Ē	Č	P	à	i	P	A
516	v	A	R	Ē	c	P	G	— <u> </u>	s	$\frac{\Lambda}{S}$
517	V	A	R	Ē	C	P	Q	i i	S	A
518	V	A	R	E	Č	P	R	Ÿ	P	A
519	V	A	R	E	<del>c</del>	P	R	V	S	S
	V									
520		A	R	E	C	Р	R	>	S	A
521 ·	> ;	A	R	E	C	P	R		Р	S
522	V	Α	R	E	C	P	R		Р	A
523	<b>\</b>	Α	R	E	С	Р	R		S	S
524	٧	Α	R	E	С	Р	R		S	Α
525·	V	Α	R	E	R	S	Q	V	Р	Α
526	V	Α	R	E	R	S	Q	V	S	S
527	V	A.	R	E	R	S	Q	V	S	A
528	V	A	R	Ē	R	S	ã	i i	P	S
529	v	A	R	Ē	R	S	ď	i	P	A
530	V	A	R	늗늗	R	S	ď	<del></del>	S	- <del>- 2</del>

		Table2. N	lovel Po	ymorphi	c Variant	of IL4R	α (cont'd	)		
Polymorphic							4.4			
Variant Number	75	02	105	400	Acid Posi 431	503			675	750
531	75 ·	82 A	185 R	E	R R	S S	576 Q	579	S S	752
532	V	A	R	E	R	S	R	<del>- '-</del>	P	A
533	V	A	R	듵	R	S	R	V	S	S
534		A	R	E	R	S	R.	V	S	
535		A	R	<del>-</del> -	R	S	R	<del></del>	P	A S
536	<u>v</u>		R		R	S.			P	
537	V	A	R	E	R	S	R		S	· A
538	V	A	R	듵	R	S	R		8	A
539	V	A	R	E	R	P	Q	\ \frac{1}{V}	P	A
540	V	A	·R	Ē	R	P.	a	<del>V</del>	S	S
541	V	A	R	E	R	P	a	V	s	·A
542	V	A	R	Ē	R	P	a	<u> </u>	P	S
543	V	A	R	E	R	P	l ä		F	A
544	V		R	E.	R	P	<del></del>		ร	
545		A	R	Ē	R	F	Q	1	S	· A
546	<u>v</u>	·A	R	E	R	P	R	V	P	A
547	V	A	R	E	R	P	R	V ·	S	S
548	V	A	R	E	R	P	R	V	S	
549		A	R	E	R	P	R	<del></del>	P	S
550	V	A	·R	Ē	. R	P	R	<del> </del>	P	A
551	Ť	A	R	Ē	R	P	R	<del>-                                    </del>	S	S
552	V	A	R	E	R	P	R		S	A
553	V	Â	R	Ā	<del>  'c  </del>	S	Q	V	P	A
554	v	A	R	A	<del>  č</del>	S	à	V	s	S
555	Ť	A	R	A	c	S	ă	V	S	A
556	v	A	R	A	Č	S	à	Ť	P	ŝ
557.	<del>-v</del>	A	R	Â	C	S	Q	<del></del>	P	Ā
558	v	Ā	R	A	C	S	-Q	<del>- i -</del>	S	S
559	v	·A·	R	A	c	S	, Q	<del></del>	S	Ā
560	v	A	R	Â	c	S	R	·	P	A
561	v	A	R	Â	C	S	R	V	s	S
562	v	A	R	A	č	S.	R	Ÿ	S	A
563	V	A	R	Ā	č	S	R	<del>- i</del>	P	s
564	V	·A	R	Â	Č	S	R	<u>.</u>	P	Ā
565	V	A	R	A	Č	S	R	i i	S	S
- 566	V	A	R	A	c	Š	R	i	S	Ā
567	V	A	R	A	č	P	à	V	P	Ā
568	V	A	R	A	Č	P	. Q	V	S	S
569	V	A	R	A	Č	P	Q	V	. S	A
570	V	A	R	Α	Ĉ	Р	Q		P	S
571	V	Α	R	A	Ċ	Р	Q	<u> </u>	P	Ā
572	V	Α	R	Α	C	Р	Q	ī	· S	S
573	V	Α	R	Α	c	Р	Q	i i	s	A
574	V	Α	R	A	Ċ	Р	R	V	P	Α
575	V	A	R	A	Ċ	P	R	V	s	s
576	V	A	R	Α	. c	Р	R	V	S	A
577	V	Α	R	A	c	P	R		P	S
578	V	A	R	Â	č	Р	R	i	P	Ā
579	V	A	R	Â	Č	P	. R	i	s	S
580	V	A	R	A	Č	P	R	<del>                                     </del>	S	A
581	v	A	R	Ā	R	S	à	v	P	A
		A	R		R	S		Ť	S	s
582	<b>\</b>	A	I • 1×	ΙA	ıĸ		Q			

		Γable2. N	Nov I Po	lymorphi	c Varian	of IL4R	α (cont'd	)		
Polymorphic										
Variant						tion and				
Number	75	82	185	400	431	503	576	579	675	752
584	V	Α_	R	Α	R	S	Q		Р	S
585	٧	Α	R	A	R	S	Q		P	Α
586	V	Α	R	A	R	S	Q		S	S
587	V	, A	R	Α	R	S	Q.		S	Α
588	V	Α	R	Α	R	S	R	V	Р	Α
589	V	A	R	Α	R	S	R	V	S	S
590	٧	Α	R	Α	R.	S	R	V	S	Α
591	V	Α	R	Α	R	S	R	1	Р	S.
592	V.	Α	R	Α.	R	S	R	l l	Р	A
593	V	Α	R	Α	R	S	R		S	S
594	V	Α	R	Α	R	S	R		S	A.
595	V	Α	R	Α	R	Р	Q	V	Р	Α
596 .	V	Α	R	Α	R	Р	Q	V	S	S
597	V	Α	R	Α	R	Р	Q	V	S	A
598	V	A	R	A	R	Р	Q	i i	P	S
599	V	A	·R	Α	R	P	Q	<u> </u>	P	Ā
600	V	A	R	A	R	P	Q	i i	s	S
601	Ÿ	A	R	Α	R	P	à	<del>-</del>	S	Ā
602	V	A	R	Â	R	P	R	Ÿ	P	A
603	v	A	R	Â	R	P	R	v	S	s
604	v	A	R	A.	R	Р	R	v	S	Ā
605	V	Ä	R	A	R	P	·R	<del>-</del>	P	s
606	V	A	R	· A	R	P	R	<del></del>	P	Ā
607	v	A	R	A	R	P	R		S	s
608	v	A	R	Â	R	P	R	<u> </u>	S	A
609	V	A	<del>-                                    </del>	Ê	C	S	Q	· V	P	S
610	V ·	A	H	Ē	<del>- 6-</del>	S		V	P	A
611	V	A	H		c	8	Q	V	S	S
	V			E			Q	V		
612 613	V	Α	H	E	C	S	Q ·		S	A
	V	Α .	Н			S	α	<u>-</u>	Р	S
614		A	H	E	C		Q	<u> !</u>	Р	A
615	V	Α	Н	E	C	S	Q	!	S	S
616	V	A	H	<u> </u>	С	S	Q	1.	S	Α
617	V	A	Н	E	С	S	R	V	Р	S
. 618	V	Α	Н	E	С	S	R	V	Р	Α
619	V	Α	Н	E	C	S	R	. V	S	S
620	V	A	Н	Е	С	S	R	V	S	A
621.	V	Α	. H	E	С	S	R	ı	Р	S
622	·V	Α	Н	Е	C	S	R		Р	Α
623	V	Α	H	E	С	S	R		S	S
624	V	Α	Н	E	C	S	R		S	Α .
625	V	Α	Н	E	С	Ρ.	Q	<b>V</b>	Р	S
626	V	Α	Н	E	С	Р	Q.	V	Р	A
627	V	Α	Н	E	C	Р	Q	V	S	S
628	V	Α	Н	E	С	Р	Q	V	S	Α
629	٧	Α	Н	· È	С	Р	Q	1	Р	S
630	V	Α	Н	E	С	Р	Q	1	Р	Α
631	V	Α	Н	Е	C	P.	Q		S	S
632	V	Α	Н	E	Ċ	Р	Q		s	Ā
633	V.	Α	H	E	C	Р	R	V	P	S
634	V	A	H	Ē	Č	Р	R	V	P	Ā
635	V	A	H	Ē	<del>- c</del>	P	R	V	S	S
636	Ť	A	Н	Ē	Ċ	P	R	V	S	Ā
030			<u> </u>		<u> </u>		_ r<		<u> </u>	_^_

· · · · · · · · · · · · · · · · · · ·		able2. N	lovel Po	lymorphi	c Variant	of IL4R	α (cont'd	)		
Polymorphic Variant				A mai m	aid Dasi	tion and	ldantitia	_		•
Number	75	82	185	400	431	tion and 503	576	579	675	752
637	V	A	H	E	<del>- 231</del>	P	R	1 1	P	8
638	v	A	H	E	c	P	R	<del>                                     </del>	P	A
639	v	A	H.	E	c	P	R	<del>                                     </del>	S	S
640	- <del>v</del>	A	Н.	Ē	C	P	R	<del>                                     </del>	s	A
641	v	A	H	E	R	S	Q	<del>  v</del>	P	S
642 ·	Ÿ	A	H	E	R	Š	à	l v	P	Ā
643	v	A	H	Ē	R	Š	a	ľ	s	S
644	Ÿ	A	H	Ē	R	s	ā	Ì	ŝ	Ā
645	V	A	H	E	R	Š	Q	l i	P	S
646	v	Α	Н	Ē	R	S	Q	<del>                                     </del>	P	Ā
647	V.	Α	Н	E	R	s	à	<del>                                     </del>	S	S
648	V	A	H	E	R ·	s	ā	<u> </u>	s	A
649	V	Α	Н	E	R	S	R	V	P	S
650	V	Α	H	E	R	s	R	V	P	A
651	V	Α	Н	E	R	s	R	V	s	S
652	V	Α	Н	E	R	S	R	V	S	Α
653	V	Α	Н	E	R	S	R		Р	S
654	ν.	. A	Н	E	R	. S	R	. 1	Р	Α
655	V	Α	Н	E	R	S	R	ı	S	S
656	V	Α	Н	E	R	S	R	ı	S	Α
657	V	Ά	Н	E	R	Р	Q	V	Р	S
658	V	Α	Н	E	R ·	Р	Q	· V	Р	Α
659	V	Α	Н	E	R	Р	Q	V	S	S
660	V	Α	Η	E	R	Ρ	a	V	S	Α
661	V	. А	Н	E	R	Р	Q	-	Р	S
662	V	Α	Н	E.	R	Р	Q		Р	Α
663	٧	Α	H	Е	R	Р	Q	l	S	S
664		Α	Н	Е	·R	Р	· Q	.	S	Α
665	V	A	H	E ·	R	Р	R	٧	Р	S
666	٧	A	Н	E	R	Р	R	V	Р	Α
667	٧	Α	Н	E	R	P	R.	V	S	S
668	V	Α	Н	E	R:	P	R	V	S	A
669	V	. A	Н	E	R	Р	R		Р	S
670	V.	A	Н	ш	R	Р	R		Р	A
671	V	Α	Н	E	R	Р.	R	!	S	S
672	V	A	Н	E	R	P C	R		S	A
673	V	A	H	Α	C	S	Q	V	Р	S
674	V	A	H	A	C	S	Q	V	P	A
675 676	V	Α	H	A	ပပ	S	Q	<b>&gt;</b>	S	S
677	V	A .	Н	A	C	S	Q	V	S	- A
678	V	A ·	Н	A	C	S	a	1	P	S
679	V	A	Н	A	C	S	Q			<u>A</u>
680		A	Н	A	C	S		1	S	S
681	- V	Ä	Н	A	C	S	Q R	V	P	S
682	V	A	Н	A	C	S	R	V	P	
683	V	A	Н	A	C	<u> </u>	R	V		<u>A</u>
684	<del>- v</del> -	A	H	A	C	<u>s</u>		V	S	S
685	V	A	H		C		R	<u> </u>	P	A
	V	A	H	A	C	S	R	<del>   </del>	P	S
				A			R			S
686	\/	_ ^								
687 688	V V	A	H	A	O O	S	R R	<u> </u>	S	A

Dahasashia	7	Fable2. N	lovel Po	lymorphi	c Variani	t of IL4Re	α (cont'd	)		
Polymorphic Variant		•	-	Amino 4	Acid Posi	ition and	Identities	2		
Number	75	82	185	400	431	503	576	579	675	752
690 .	V	Ā	Н	A	C	P	Q	V	P	A
691	V	Α	Н	A	Č	P	à	V	S	S
692	V	Α	Н	Α	c	P	Q	V	S	Ā
693	V	Α	Н	A	C	P	Q		P	S
694	V	Α	Н	Α	C	P	Q		P	- Ā
695	V	· A	Н	Α	С	P	Q		S	S
696	V	Α	Н	A.	С	P	Q	T	S	Α
697	V	A	Н	Α	C	Р	R	V	P	S
698	V	Α	H	Α	С	. Р	R	V	Р	Α
699	V	Α	Н	Α	С	Р	R	V	S	·S
700	· V	Α	н	Α	С	Р	R	V	S	Α
· 701	٠٧.	Α	Н	Α	· C	P	Ŕ		Р	S
702	V	Α	Н	Α	С	P	R		P	Α
703	٧	Α	Н	Α	С	Р	R	ı	S	S
· 704	٧	Α	Н	. A	С	Р	R	1	S	Α
705	<b>V</b>	Α	Н	Α	R	S	Q	<b>V</b>	Р	S
706	<b>V</b>	Α	Н	Α	R	S	Q	٧	Р	A
707	٧	Α	.H	Α	R	S	Q	V	S	S
708	V	Α	H	Α	R	S	Q	V	S	Α
709	V	Α	Н	Α	R.	S	Q	1	Р	S
710	V	Α	Н	Α	R	S	Q	ĺ	Р	Α
711	V	Α	Н	Α	R	S	a		S	S
712	<b>V</b>	Α	· H	Α	R	S	Q		S	Α
713	· V	Α	H	Α	R	S	R	٧	Р	S
714	V	Α	Н	Α	R	S	R	V	Р	Α
715	V	Α.	Н	Α	R	S	R	<b>&gt;</b>	S	S
716	V	Α	Н	Α	Ŗ	S.	R	V	S	Α
717	<b>V</b>	Α	Н	Α	R	S	R	1 .	Р	S
718	<b>V</b>	Α	Н	Α	R	S	R		Р	Α_
719	V	Α	Н	Α	R	S	R		S	S
720	V	Α	Н	. <b>A</b>	R	S	R		S	Α
721 .	<b>&gt;</b>	Α	. Н	Α	R	Р	Q		Ρ.	S
722	V	Α	Н	Α	R	Р	Q	V	Р	A
723	V	Α	H	Α	R	. P	Q	V	S	S
724	V	A	Н	Α	R	Р	Q	V	S	Α
725	٧	A	Н	A.	R	Р	Q		Р	S
726	V	Α	H	Α	R	Р	. Q		P	Α
727	<u> </u>	A	H	Α	R	P	Q	<u> </u>	S	S
728	V	A	H	A	R	Р	αı		S	A
729	V	A	H	A	R	P	R	> >	Р	S
730	> ;	A	H	A	R	P	R	V	Р	<u>A</u>
731	V	A	Н	Α	.R	Р	R	<b>&gt;</b>	S	S
732	<b>V</b>	A	H	A	R	. P	R	<b>&gt;</b>	S	A
733	> ;	A	Н	Α	R	Р	R	1	P	S
734	V	A	H	A	R	. <u>b</u>	R	!	Р	A
735 ·	<b>&gt;</b>	A .	H	Α	R	Р	R	· I	S	S.
736	V	A	H	A	R	. Р.	R		. S	A
737	V	Τ.	R	E	C	S	Q	٧	Р	S
738	<b>V</b>	T	R	E	С	S	·Q	V	Р	Α
739	V	T	R	E	С	S	Q	V	S	S
740	V	T	R	E	· C	S	Q	V	S	A
741	V	T	R	E	С	S	Q		Р	S
742	V	T	R	E	С	S	Q		Р	Α

	7	Fable2. N	lovel Po	lymorphi	c Variant	of IL4R	α (cont'd	)		· ·
Polymorphic Variant				Amino 4	Acid Posi	ition and	Identities	2		•
Number	75	82	185	400	431	503	576	579	675	752
743	V	T	R	E	C	S	Q	1	S	S
744	V	Ť	R	Ē	č	S	à	i	S	Ā
745 <sup>-</sup>	V	Ť	R	E	Č	S	R	v	P	S
746	V	Ť	R	. E	c	S	R	V	P	Ā
747	V	Ť	R	E	Ċ	S	R	v	S	s
748	V	T	R	E	Ċ	S	R	V	S	Ā
749	V	T	R	E	C	S	R		P	S
750	V	Т	R	E	С	S	R		P	A
751	V	T	R	E	С	S	R		S	S
752	V	·T	R	E	Ç	S	R	1 .	S	A
753	V	T	R	Ε	Ċ	Р	Q	V	Р	S
754	V	Т	R	E	С	Р	Q	V	·P	A
755	V	Т	R	E	С	Р	Q	V	S	S
756	V	T	R	E	С	Р	Q	V	S	A
757	V	T	R	E	С	Р	Q	ı	Р	S
758	V	T	R	E	С	Р	Q	ı	Р	Α
759	V	T	R	E	С	Р	Q	T	S	S
760	V	T	R	E	. C	Р	Q		S	Α
761	٧	Т	R	E	С	Р	R	V	Р	S
762	V	T	R	· E	С	Р	R	· V	Р	Α
763	<b>V</b>	Т	R	E	С	Р	R.	· >	·S	S
764	٧	T	R	E	С	.P	R	V	S	Α
· 765	V	T	R	E	C ·	Р	R		Р	S
766	V	T	R	E	С	Р	R		Р	Α
767	V	T	R	E	С	Р	R		S	S
768	V	T	R	E	С	Р	R		S	Α
769	V	T	R	E	R	S	Q	>	Р	S
770	V	<u>T</u>	R	<u> </u>	R	· s	Q	V	P.	Α
771	V	T	R	E	R	S	Q	V	S	S
772	V	- T	R	E	R	S	Q ·	V	S	Α
773	V	Ţ	R	E	·R	S	Q		P	S
774	V ·	Ţ	R	E	R	S	Q		Р	A
775	V	<u></u>	R	E	R	_ S	Q		S	S
776	<b>&gt;</b>	T	R	E	R	S	Q	1	S	A
777	V	Ţ	R	E	R	S	R	>>	P	S
778 779	> >	T	R	E	R	S	R R	V	P S	S
780	V	+	R	E	R	S	R	V	S	
781	V	<del></del>	R	E	R	S	R	1	P	S
782	V	<del>-                                    </del>	R	E	R	S	R	-	P	A
783	·V	<del></del>	R	Ē	R	S	R	<del>   </del>	S	S
784	V	Ť	R	E	R	S	R	<del>                                     </del>	S	A
785	V	Ť	R	E	R	P	Q	V	P	S
786	V	Ť	R	Ē	R	P.	Q	V	P	A
787	V	Ť	R	E	R	P	Q	V	S	S
788 ·	v	<del></del> -	R	E	R	P	Q	V	S	A
789	V.	Ť	R	Ē	R	P	Q	- 1	P	ŝ
790	V.	<del>-                                    </del>	R	E	R	P	Q		P	A
791	V.	<del>-                                    </del>	R	Ē	R	P	Q		S	S
792	V	T	R	듣	R	P	Q	+	S	A
793	V	Ť	R	퉅	R	P	R	V	P	S
794	V	Ť	R	E	R	P	R	V	P	A
795	V	<del></del>	R	È	R	P	R	- v	S	S
1 20								٧	_ ح	

Polymorphia	1	Table2. N	lovel Po	ymorphi	c Variant	of IL4R	α (cont'd	)				
Polymorphic Variant	Amino Acid Position and Identities											
	75	00	405			075						
Number	75	82	185	400	431	503	576	579	675	752		
796		Ī	R	E	R	Р	R	V	S	A		
797		Ţ	R	E	Ŗ	Р	R	- !	P	S		
798	V	Ţ	R	E	R	Р	R	<u> </u>	P	Α		
799	\ \	T	R	E	R	Р	R		S	S		
800		T	R	E	R	Р	R		S	Α		
801	V	T	R	· A	C	S	Q	V.	Р	S		
802	V	T	R	A	С	S	Q	V	p.	Α		
803	٧	Τ :	R	Α	С	S	Q	V	S	·S		
804	٧	T	R	Α	С	S	Q	V	S	·A		
805	V	T	R	Α	С	S	Q	ı	. P	S		
806	V	T	R	Α	С	S	Q	1	P.	Α		
807	·V	T	R	Α	С	S	Q		S	S		
808	V	Т	R	A	С	S	Q		S	A		
809	V	ī	R	A	C	S	R	V	P	S		
810	v	T	R	A	Č	S	R	V	P	Ä		
811	v	Ť	R	A	Č	S	·R	V	S	S		
812	v	Ť	R	A	č	S	R	v	S	Ä		
813	v	Ť	R	A	Č	· S	R	i	·P	S		
814	v	Ť	R	A	C	·S	. R	- 1	P	A		
815	v	Ť	R	Ä	Č	S	R		s	S		
816	Ť	Ť	R	A	C	S	R	<del>- :</del>	. S	A		
817	v	Ť	·R	A	C	P	à	Ÿ	P	ŝ		
818	v	Ť	R	A	<del>  c</del>	P	à	V	P	A		
819	V	Ť	R	A	C	P		V				
820	V	+	R		č	P.	Q	V	S	S		
821	$\overline{}$	Ť	R	- A	C	P	Q	V	S	<u>A</u>		
822	v	<del></del>		A			Q			S		
			R.	Α	C	P P	· Q	<u> </u>	Р	A		
823	<u> </u>		R	A	C	P	Q	!	S	S		
824	V	T	R	A	С	P	Q	-1	S	A		
825	V	T	. R	A	С	P	R ·	V	Р	S		
826	V	<u>. T</u>	R	Α	C	Р	R	۷ .	Р	· A		
827	V	<u>T</u>	R	Α	С	Р	R	V	S	S		
828	V	T	R	Α	C	P	R	V	S	Α		
829	V	Т	R	Α	С	Р	R	1 .	Р	S		
830	V	T	R	Α	С	P_	R		Р	Α		
831 .	V	T	.R	Α	С	P ·	R		S	S		
832	V	T	R	Α	С	Р	R		S	Α		
833	V	T	·R	Α	R	S	Q	V	Р	S		
834	V	T	R	Α	R	· S	Q	V	Р	Α		
835	V	T	R	Α	R	S	Q	V	S	S		
836	V	T	R	Α	R	S	Q	V	S	_ A		
837	V	T	R	Α	R	S	Q	T I	Р	S		
838	V	T	R	Α	R	S	Q		Р	Α		
839	V	Т	R	Α	R	S	Q	1	S	S		
.840	v	Т	R	Α	R	S	Q	i	S	A		
841	V	Ť	R	A	R	S	Ř	V	P	S		
842	Ÿ	Ť	R	A	R	S	R	Ÿ	P	Ā		
843	v	Ť	R	A	R	S	R	V	S	S		
844	<del>v</del> l	T	R	Â	R	S	R	V	S	<u>S</u>		
845	V.	+	R	A	R	S	R	V	P	S		
846	V	+	·R		R				P			
	<del>-v</del> -	+	R	A	R	S	R			A S		
847									S			

	7	rable2. N	lovel Po	ymorphi	c Variant	of IL4R	z (cont'd	)		
Polymorphic							1.1	_		
Variant	75	00	405		_	tion and			1 056	T 750
Number	75 V	82	185	400	431	503	576	579	675	752
849	V V	T	R	A	R	P	Q_	V	Р	S
850	V		R.	A	R		Q		Р	A
851		T	R	A	R	Р	Q	V	S	S
852	\ \	<u>T</u>	R	A	R	Р	Q.	V	S	A ·
853	\ \	Ţ	R	A	R	P	<u>Q</u>	<u> </u>	Р	S
854	V	T	R	Α	R	Р	Q	·	Р	A
855	V	T	R	A	R	Р	Q	1	S	S
856	V	Ţ	R	A	R	Р	Q	1	S	<u>, Α</u>
857	\ \	T	R	A	R	Р	R	V	P	S
858		T	R	· A	R	Р	, R	V	Р	A
859	V	T	R	A	R	Р	R	V	S	S
860	V	T	R	Α	R.	Р	R	V	S	Α
861	\ \ \	Ţ	R	Α	R	Р	R	<u> </u>	Р	S
862		<u> </u>	R	Α	R	Р	, R	<u> </u>	P	A
863		T	R	Α	R	Р	R	· 1	S	S
864	V	<u>T</u>	R	_ <u>A</u>	R	Р	R		S	A
865	V	. T	H	·E	C	S	Q	V	Р	S
866	V	T	H	<u>E</u>	С	S	Q	V	Р	Α
867	V	T	Н	E	С	S	Q	>	s	S
868	V	Т	Н	· E	С	S	Q	V	·S	A
869		T	Н	E	С	S	Q	ı	Р	S
870	V .	Т	Н	E	С	S	Q	1	Р	Α
871		T	Н	E	C	S	Q		S	S
872		T	Н	E	С	S	Q	1	S	Α
873	ν.	T	Τ	Е	С	S	R	٧	_ P	S
874	V	T	H	E	С	S	R.	٧	Р	Α
875	V	Т	H	E	Ç	S	R	>	S	S
876		Т	H	E	С	S	R	٧	·S	Α
877	٧	T	H	E	С	S	R		Р	S
878	V	Ţ	H	E	С	S	Ŗ	-	Р	Α
879	V.	T	Н	Ш	С	S	R		S	S
880	٧	Τ.	H	E	C .	S	R	1	S	Α
881	٧	T	Н	ш	С	Р	Q	V	Р	S
882	. V	Т	Н	E	С	Ρ.	Q	V	Р	A
883	V	T	Н	E	С	Р	Q	V	S.	S
884	V	_ T	Н	E	С	Р	Q	<b>V</b>	S	Α
885	V	T	Н	E	· C	Р	Q	ı	Р	S
886	V	Ť	· H	Е	C	Р	Q	1	Р	Α
887		T ·	Н	E	င	Р	Q	1	S	S
888	· V	T	Н	E	С	Р	Q	ı	S	Α
889	V	. T	Н	E	С	P	R	<b>V</b>	·P	. S
890	V	T	Н	E	С	Р	R	V	Р	Α
891	V	T	Н	E	С	Р	R	٧	S	S
892	V	T	Н	E	С	Р	R	V	S	Α
893	V	T	Н	E	С	Р	R	i	Р	S
894	V	T	Н	E	С	P	R		Р	Ā
895	V	T	Н	E	Ċ	P	R∙	i	S	S
896	Ÿ	Ť	Н	Ē	Č	P	R	<del>-i</del>	S	Ā
897	v	Ť	H	Ē	·R	s	Q	Ÿ	P	$\frac{\alpha}{s}$
898	v	<del>- i</del> -	H	Ē	R	S	Q	v	P	$\frac{3}{A}$
899	v	Ť	H	E	R	S.	a	v	S	$-\hat{s}$
900	Ť	÷	H	E	R	S	à	V	S	$\frac{3}{A}$
901	Ť	<del>-                                    </del>	H	E	R	S	à		P	<del>- ^</del>

	1	Tabl 2. N	lovel Po	ymorphi	c Variant	of IL4R	α (cont'd	)	<u>-</u> -	
Polymorphic										
Variant					cid Posi					
Number	75	82	185	400	431	503	576	579	675	752
902	V	T	Н	E	R	S	Q		Р	Α
903	V	T	Н	E	R	S	Q	<u> </u>	S	S
904	V	1	Н	E	R	S	Q		S	Α
905	V	1	Н	E	R	S	R	٧	Р	S
906	V	T	Н	Е	R	S	R	V	Р	Α
907	V	T	Н	Е	R	S	R	V	S	S
908	V	T	Н	E	R	S	R	V	S	Α
909	V	T	Н	E	R	S	R	ı	Р	S
910	V	T	Н	E	R	S	R	T	Р	·A
911	V	T·	Н	E	R	S	R		S	S
912	V	T	Н	E	R	S	R		S	A
913	v	T	H	E	R	P	Q	v	P	S
914	v	Ť	H	Ē	R	Р	Q	Ÿ	P	Ā
915	v	Ť	H	Ē	R	P	Q	v	s	ŝ
916	V	- <del>;</del>	H	Ē	R	P	à	V	S	A
917	Ť	÷	H	E	R	P	Q	Ť	P	S
918	·	+	Н	E	R	P.	ă	<del></del>	P	A
919	v	÷	Н.	Ē	R	· P	à	<del></del>	S	- <del>S</del>
920	Ť	Ť	Н	Ē	R	P	Q	<del>-                                    </del>	S	A
921	V	Ť.	Н	E	R	P	R	V.	P	S
921	- <del></del>	<del></del> -	Н			P		V	P	
	V	+		E	R		R	V		· A
923			Н	<u>E</u> .	R	P	R		S	S
924	V	T	H	E	R	P	R	V	S.	A
925	V	T	H	E	R	Р	R	!	Р	S
926	ν.	T	Н	E	R	Р	R		Р	Α
927		T	H	E	R	Р	R		S	S
928		Ţ	H	E	R	Р	R		S	Α
929	٧	T	Н	A	С	S	Q	V	Р	S
930	V	<u>T</u>	Н	Α	C ·	S	Q	V	Р	Α
931	V	<u>T</u>	H	Α	С	S	Q	V	S	S
932	V	T	Н	Α	С	S	Q	٧	S	Α
933	<b>V</b>	T	. Н	Α	C ·	S	Q		Р	S
934	V	T	Н	Α	С	S	Q	<u> </u>	.P	Α
935 .	V	T	Н	Α	С	S	Q	1.	S	S
936	V	T	H	Α	С	S	Q		S	Α
937	V	T	Н	Α	С	S	R	· V	Р	S
938	V	T	Н	Α	С	S	R	V	Р	Α
939	V	T	H	Α	С	S	R	٧	S	S
940	V	T	Н	Α	С	S	R	٧	S	Α
941	V	T	Н	Α	С	S	R	1	Р	S
942 ·	V	T	Η.	Α	С	S	R	ı	Ρ.	Α
943	V	T	Н	Α	С	S	R	ı	S	S
944	V	T	Н	Α	С	S	R	ī	S	Α
945	V	Ť	Н	Α	С	Р	Q	·V	Р	S
946	V	T	Н	A	С	Р	Q	V	P	A
947	V	Ť	Н	Α	Ċ	Р	Q	V	S	S
948	V	Ť	H	A	Č	P	Q	Ý	S	Ā
	v	Ť	H	A	c	P	à	<u> </u>	P	S
949	Ť	+	H	A	C	P	Q	. ;	P	A ·
949 950		1								
950		Ť		Δ		D	C.	1	C	C
950 951	V	T	Н	A	C	P	άc	1	S	S
950		T		A A A	CCC	P P	Q Q R	>	S S P	S A S

	•	Table2. N	lovel Po	lymorphi	c Variant	of IL4Re	x (cont'd	)		•			
Polymorphic									•				
Variant		Amino Acid Position and Identities											
Number	75	82	185	400	431	503	576	579	675	752			
955	V	T	Н	Α	C	Р	R	V	S	S			
956	V	T	Н	Α	С	P	R	V	S	A			
957	V	T	Н	A	С	P.	R	1	Р	S			
958	V	T	Н	Α	С	Р	R	T	. P	Α			
959	V	T	Н	Α	С	Р	R	1	S	S			
960	V	T	Н	A	С	Р	R	1	S	A			
961	V	T	Н	Α	R	S	Q	· V	Р	S			
962	V	T	Н	Α	R	S	Q	V	Р	A			
963 .	V	T	Н	A	R	S	· Q	V	S	S			
964	V	Т	H	Α	R	S	Q	V	S	Α			
965	V	. T	Н	A	R	S	Q		Р	S			
966	V	T	Н	Α	R	S	Q		Р	A			
967	V	Т	Н	Α	R	S	Q	1 7	S	s			
968	V	T	H	A	R	S	Q	T	S	A			
969	· V	T	Н	A	R	S	R	V	P	S			
970	V	T	. H	A	·R	·S	R	V	Р	A			
971	V	T	Н	A	·R	.S	R	V	S	S			
972 -	V	Ŧ	Н	. A	R	S	R	V	S	A			
973	V	T	Н	Α	R.	S	. R	ı	Р	S			
974	V	T	. н	A	R	S	R		Р	Α			
975	V	T	Н	A	·R	S	R	T	S	S			
976	V	Т	Н	A	R	S	R	1	S	A			
977	ν.	Т	Н	Α	R	Р	Q	V	Р	S			
978	<b>V</b> .	T	Н	Α	R	Р	Q	V	Р	Α			
979	V	T	Н	Α	R	Р	Q	V	S	S			
980	V	T	Н	A.	·R	Р	Q	V	S	A			
981	V	T	Н	Α	R	·P	Q	l "	Р	S			
982	V	Т	. H	Α	R	Р	Q	1	Р	Α			
983	V	Т	Н	Α	R	Р	Q	I	S	S			
984	V	Т	Н	Α	·R	Р	Q	T.	S	Α			
985	V	Т	Н	Α	R	P	R	V	Р	s			
986	V	T	Н	Α	R	Р	R	V	Р	Α			
987	V	Т	Н	Α	R	Р	R	V	·S	S			
988	V	T	Н	Α	R	Р	R	V	S	A			
989	V	Т	Н	A	R	· P	R		Ρ.	S			
990.	V	T	Н	Α	R	Р	R		Р	A			
991	V	Ť	Н	Α	R	Р	R	1	S	S			
992	V	T	Н	A	R	P	R	T	S	Ā			

The invention also includes IIARα peptide variants, which are any fragments of an IIARα protein variant that contains one or more of the amino acid variations shown in Table 2. An IIARα peptide variant is at least 6 amino acids in length and is preferably any number between 6 and 30 amino acids long, more preferably between 10 and 25, and most preferably between 15 and 20 amino acids long. Such IIARα peptide variants may be useful as antigens to generate antibodies specific for one of the above IIARα isoforms. In addition, the IIARα peptide variants may be useful in drug screening assays.

An ILAR $\alpha$  variant protein or peptide of the invention may be prepared by chemical synthesis or by expressing one of the variant ILAR $\alpha$  genomic and cDNA sequences as described above. Alternatively, the ILAR $\alpha$  protein variant may be isolated from a biological sample of an individual having an ILAR $\alpha$ 

10

isogene which encodes the variant protein. Where the sample contains two different ILAR $\alpha$  isoforms (i.e., the individual has different ILAR $\alpha$  isogenes), a particular ILAR $\alpha$  isoform of the invention can be isolated by immunoaffinity chromatography using an antibody which specifically binds to that particular ILAR $\alpha$  isoform but does not bind to the other ILAR $\alpha$  isoform.

5

10

15

20

. 25

30

35

The expressed or isolated IL4R $\alpha$  protein may be detected by methods known in the art, including Coomassie blue staining, silver staining, and Western blot analysis using antibodies specific for the isoform of the IL4R $\alpha$  protein as discussed further below. IL4R $\alpha$  variant proteins can be purified by standard protein purification procedures known in the art, including differential precipitation, molecular sieve chromatography, ion-exchange chromatography, isoelectric focusing, gel electrophoresis, affinity and immunoaffinity chromatography and the like. (Ausubel et. al., 1987, In Current Protocols in Molecular Biology John Wiley and Sons, New York, New York). In the case of immunoaffinity chromatography, antibodies specific for a particular polymorphic variant may be used.

A polymorphic variant ILAR $\alpha$  gene of the invention may also be fused in frame with a heterologous sequence to encode a chimeric ILAR $\alpha$  protein. The non-ILAR $\alpha$  portion of the chimeric protein may be recognized by a commercially available antibody. In addition, the chimeric protein may also be engineered to contain a cleavage site located between the ILAR $\alpha$  and non-ILAR $\alpha$  portions so that the ILAR $\alpha$  protein may be cleaved and purified away from the non-ILAR $\alpha$  portion.

An additional embodiment of the invention relates to using a novel IIARα protein isoform in any of a variety of drug screening assays. Such screening assays may be performed to identify agents that bind specifically to all known IIARα protein isoforms or to only a subset of one or more of these isoforms. The agents may be from chemical compound libraries, peptide libraries and the like. The IIARα protein or peptide variant may be free in solution or affixed to a solid support. In one embodiment, high throughput screening of compounds for binding to an IIARα variant may be accomplished using the method described in PCT application WO84/03565, in which large numbers of test compounds are synthesized on a solid substrate, such as plastic pins or some other surface, contacted with the IIARα protein(s) of interest and then washed. Bound IIARα protein(s) are then detected using methods well-known in the art.

In another embodiment, a novel ILAR $\alpha$  protein isoform may be used in assays to measure the binding affinities of one or more candidate drugs targeting the ILAR $\alpha$  protein.

In another embodiment, the invention provides antibodies specific for and immunoreactive with one or more of the novel ILAR $\alpha$  variant proteins described herein. The antibodies may be either monoclonal or polyclonal in origin. The ILAR $\alpha$  protein or peptide variant used to generate the antibodies may be from natural or recombinant sources or produced by chemical synthesis using synthesis techniques known in the art. If the ILAR $\alpha$  protein variant is of insufficient size to be antigenic, it may be conjugated, complexed, or otherwise covalently linked to a carrier molecule to enhance the antigenicity of the peptide. Examples of carrier molecules, include, but are not limited to, albumins (e.p. human.

bovine, fish, ovine), and keyhole limpet hemocyanin (Basic and Clinical Immunology, 1991, Eds. D.P. Stites, and A.I. Terr, Appleton and Lange, Norwalk Connecticut, San Mateo, California).

In one embodiment, an antibody specifically immunoreactive with one of the novel IL4R $\alpha$  protein isoforms described herein is administered to an individual to neutralize activity of the IL4R $\alpha$  isoform expressed by that individual. The antibody may be formulated as a pharmaceutical composition which includes a pharmaceutically acceptable carrier.

Antibodies specific for and immunoreactive with one of the novel ILAR $\alpha$  protein isoform described herein may be used to immunoprecipitate the ILAR $\alpha$  protein variant from solution as well as react with ILAR $\alpha$  protein isoforms on Western or immunoblots of polyacrylamide gels on membrane supports or substrates. In another preferred embodiment, the antibodies will detect ILAR $\alpha$  protein isoforms in paraffin or frozen tissue sections, or in cells which have been fixed or unfixed and prepared on slides, coverslips, or the like, for use in immunocytochemical, immunohistochemical, and immunofluorescence techniques.

10

15

20

25

30

35

In another embodiment, an antibody specifically immunoreactive with one of the novel IL4Ra protein variants described herein is used in immunoassays to detect this variant in biological samples. In this method, an antibody of the present invention is contacted with a biological sample and the formation of a complex between the IL4R\alpha protein variant and the antibody is detected. As described, suitable immunoassays include radioimmunoassay, Western blot assay, immunofluorescent assay, enzyme linked immunoassay (ELISA), chemiluminescent assay, immunohistochemical assay, immunocytochemical assay, and the like (see, e.g., Principles and Practice of Immunoassay, 1991, Eds. Christopher P. Price and David J. Neoman, Stockton Press, New York, New York; Current Protocols in Molecular Biology, 1987, Eds. Ausubel et al., John Wiley and Sons, New York, New York). Standard techniques known in the art for ELISA are described in Methods in Immunodiagnosis, 2nd Ed., Eds. Rose and Bigazzi, John Wiley and Sons, New York 1980; and Campbell et al., 1984, Methods in Immunology, W.A. Benjamin, Inc.). Such assays may be direct, indirect, competitive, or noncompetitive as described in the art (see, e.g., Principles and Practice of Immunoassay, 1991, Eds. Christopher P. Price and David J. Neoman, Stockton Pres, NY, NY; and Oellirich, M., 1984, J. Clin. Chem. Clin. Biochem., 22:895-904). Proteins may be isolated from test specimens and biological samples by conventional methods, as described in Current Protocols in Molecular Biology, supra.

Exemplary antibody molecules for use in the detection and therapy methods of the present invention are intact immunoglobulin molecules, substantially intact immunoglobulin molecules, or those portions of immunoglobulin molecules that contain the antigen binding site. Polyclonal or monoclonal antibodies may be produced by methods conventionally known in the art (e.g., Kohler and Milstein, 1975, Nature, 256:495-497; Campbell Monoclonal Antibody Technology, the Production and Characterization of Rodent and Human Hybridomas, 1985, In: Laboratory Techniques in Biochemistry and Molecular Biology, Eds. Burdon et al., Volume 13, Elsevier Science Publishers, Amsterdam). The

antibodies or antigen binding fragments thereof may also be produced by genetic engineering. The technology for expression of both heavy and light chain genes in E. coli is the subject of PCT patent applications, publication number WO 901443, WO 901443 and WO 9014424 and in Huse et al., 1989, Science, 246:1275-1281. The antibodies may also be humanized (e.g., Queen, C. et al. 1989 Proc. Natl. Acad. Sci. 86:10029).

Effect(s) of the polymorphisms identified herein on expression of ILARα may be investigated by preparing recombinant cells and/or organisms, preferably recombinant animals, containing a polymorphic variant of the ILARα gene. As used herein, "expression" includes but is not limited to one or more of the following: transcription of the gene into precursor mRNA; splicing and other processing of the precursor mRNA to produce mature mRNA; mRNA stability; translation of the mature mRNA into ILARα protein (including codon usage and tRNA availability); and glycosylation and/or other modifications of the translation product, if required for proper expression and function.

To prepare a recombinant cell of the invention, the desired ILARα isogene may be introduced into the cell in a vector such that the isogene remains extrachromosomal. In such a situation, the gene will be expressed by the cell from the extrachromosomal location. In a preferred embodiment, the ILARα isogene is introduced into a cell in such a way that it recombines with the endogenous ILARα gene present in the cell. Such recombination requires the occurrence of a double recombination event, thereby resulting in the desired ILARα gene polymorphism. Vectors for the introduction of genes both for recombination and for extrachromosomal maintenance are known in the art, and any suitable vector or vector construct may be used in the invention. Methods such as electroporation, particle bombardment, calcium phosphate co-precipitation and viral transduction for introducing DNA into cells are known in the art; therefore, the choice of method may lie with the competence and preference of the skilled practitioner. Examples of cells into which the ILARα isogene may be introduced include, but are not limited to, continuous culture cells, such as COS, NIH/3T3, and primary or culture cells of the relevant tissue type, i.e., they express the ILARα isogene. Such recombinant cells can be used to compare the biological activities of the different protein variants.

Recombinant organisms, i.e., transgenic animals, expressing a variant II.4R $\alpha$  gene are prepared using standard procedures known in the art. Preferably, a construct comprising the variant gene is introduced into a nonhuman animal or an ancestor of the animal at an embryonic stage, i.e., the one-cell stage, or generally not later than about the eight-cell stage. Transgenic animals carrying the constructs of the invention can be made by several methods known to those having skill in the art. One method involves transfecting into the embryo a retrovirus constructed to contain one or more insulator elements, a gene or genes of interest, and other components known to those skilled in the art to provide a complete shuttle vector harboring the insulated gene(s) as a transgene, see e.g., U.S. Patent No. 5,610,053. Another method involves directly injecting a transgene into the embryo. A third method involves the use of embryonic stem cells. Examples of animals into which the II.4R $\alpha$  isogenes may be introduced include,

but are not limited to, mice, rats, other rodents, and nonhuman primates (see "The Introduction of Foreign Genes into Mice" and the cited references therein, In: Recombinant DNA, Eds. J.D. Watson, M. Gilman, J. Witkowski, and M. Zoller; W.H. Freeman and Company, New York, pages 254-272). Transgenic animals stably expressing a human ILARα isogene and producing human ILARα protein can be used as biological models for studying diseases related to abnormal ILARα expression and/or activity, and for screening and assaying various candidate drugs, compounds, and treatment regimens to reduce the symptoms or effects of these diseases.

5

10

15

20

25

30

35

An additional embodiment of the invention relates to pharmaceutical compositions for treating disorders affected by expression or function of a novel ILARa isogene described herein. The pharmaceutical composition may comprise any of the following active ingredients: a polynucleotide comprising one of these novel ILARa isogenes; an antisense oligonucleotide directed against one of the novel ILARa isogenes, a polynucleotide encoding such an antisense oligonucleotide, or another compound which inhibits expression of a novel ILAR isogene described herein. Preferably, the composition contains the active ingredient in a therapeutically effective amount. By therapeutically effective amount is meant that one or more of the symptoms relating to disorders affected by expression or function of a novel ILARα isogene is reduced and/or eliminated. The composition also comprises a pharmaceutically acceptable carrier, examples of which include, but are not limited to, saline, buffered saline, dextrose, and water. Those skilled in the art may employ a formulation most suitable for the active ingredient, whether it is a polynucleotide, oligonucleotide, protein, peptide or small molecule antagonist. The pharmaceutical composition may be administered alone or in combination with at least one other agent, such as a stabilizing compound. Administration of the pharmaceutical composition may be by any number of routes including, but not limited to oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, intradermal, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

For any composition, determination of the therapeutically effective dose of active ingredient and/or the appropriate route of administration is well within the capability of those skilled in the art. For example, the dose can be estimated initially either in cell culture assays or in animal models. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. The exact dosage will be determined by the practitioner, in light of factors relating to the patient requiring treatment, including but not limited to severity of the disease state, general health, age, weight and gender of the patient, diet, time and frequency of administration, other drugs being taken by the patient, and tolerance/response to the treatment.

Information on the identity of genotypes and haplotypes for the ILARa gene of any particular

individual as well as the frequency of such genotypes and haplotypes in any particular population of individuals is expected to be useful for a variety of basic research and clinical applications. Thus, the invention also provides compositions and methods for detecting the novel ILAR $\alpha$  polymorphisms identified herein.

5

10

15

20

25

30 ·

35

The compositions comprise at least one ILARa genotyping oligonucleotide. In one embodiment, an IL4Ra genotyping oligonucleotide is a probe or primer capable of hybridizing to a target region that is located close to, or that contains, one of the novel polymorphic sites described herein. As used herein, the term "oligonucleotide" refers to a polynucleotide molecule having less than about 100 nucleotides. A preferred oligonucleotide of the invention is 10 to 35 nucleotides long. More preferably, the oligonucleotide is between 15 and 30, and most preferably, between 20 and 25 nucleotides in length. The oligonucleotide may be comprised of any phosphorylation state of ribonucleotides, deoxyribonucleotides, and acyclic nucleotide derivatives, and other functionally equivalent derivatives. Alternatively, oligonucleotides may have a phosphate-free backbone, which may be comprised of linkages such as carboxymethyl, acetamidate, carbamate, polyamide (peptide nucleic acid (PNA)) and the like (Varma, R. in Molecular Biology and Biotechnology, A Comprehensive Desk Reference, Ed. R. Meyers, VCH Publishers, Inc. (1995), pages 617-620). Oligonucleotides of the invention may be prepared by chemical synthesis using any suitable methodology known in the art, or may be derived from a biological sample, for example, by restriction digestion. The oligonucleotides may be labeled, according to any technique known in the art, including use of radiolabels, fluorescent labels, enzymatic labels, proteins, haptens, antibodies, sequence tags and the like.

Genotyping oligonucleotides of the invention must be capable of specifically hybridizing to a target region of an ILARα polynucleotide, i.e., an ILARα isogene. As used herein, specific hybridization means the oligonucleotide forms an anti-parallel double-stranded structure with the target region under certain hybridizing conditions, while failing to form such a structure when incubated with a non-target region or a non-ILARα polynucleotide under the same hybridizing conditions. Preferably, the oligonucleotide specifically hybridizes to the target region under conventional high stringency conditions. The skilled artisan can readily design and test oligonucleotide probes and primers suitable for detecting polymorphisms in the ILARα gene using the polymorphism information provided herein in conjunction with the known sequence information for the ILARα gene and routine techniques.

A nucleic acid molecule such as an oligonucleotide or polynucleotide is said to be a "perfect" or "complete" complement of another nucleic acid molecule if every nucleotide of one of the molecules is complementary to the nucleotide at the corresponding position of the other molecule. A nucleic acid molecule is "substantially complementary" to another molecule if it hybridizes to that molecule with sufficient stability to remain in a duplex form under conventional low-stringency conditions.

Conventional hybridization conditions are described, for example, by Sambrook J. et al., in Molecular

Cloning, A Laboratory Manual, 2<sup>nd</sup> Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989)

and by Haymes, B.D. et al. in Nucleic Acid Hybridization, A Practical Appr ach, IRL Press, Washington, D.C. (1985). While perfectly complementary oligonucleotides are preferred for detecting polymorphisms, departures from complete complementarity are contemplated where such departures do not prevent the molecule from specifically hybridizing to the target region. For example, an oligonucleotide primer may have a non-complementary fragment at its 5' end, with the remainder of the primer being complementary to the target region. Alternatively, non-complementary nucleotides may be interspersed into the oligonucleotide probe or primer as long as the resulting probe or primer is still capable of specifically hybridizing to the target region.

Preferred genotyping oligonucleotides of the invention are allele-specific oligonucleotides. As used herein, the term allele-specific oligonucleotide (ASO) means an oligonucleotide that is able, under sufficiently stringent conditions, to hybridize specifically to one allele of a gene, or other locus, at a target region containing a polymorphic site while not hybridizing to the corresponding region in another allele(s). As understood by the skilled artisan, allele-specificity will depend upon a variety of readily optimized stringency conditions, including salt and formamide concentrations, as well as temperatures for both the hybridization and washing steps. Examples of hybridization and washing conditions typically used for ASO probes are found in Kogan et al., "Genetic Prediction of Hemophilia A" in PCR Protocols, A Guide to Methods and Applications, Academic Press, 1990 and Ruano et al., 87 Proc. Natl. Acad. Sci. USA 6296-6300, 1990. Typically, an allele-specific oligonucleotide will be perfectly complementary to one allele while containing a single mismatch for another allele.

10

20

25

Allele-specific oligonucleotide probes which usually provide good discrimination between different alleles are those in which a central position of the oligonucleotide probe aligns with the polymorphic site in the target region (e.g., approximately the 7<sup>th</sup> or 8<sup>th</sup> position in a 15 mer, the 8<sup>th</sup> or 9<sup>th</sup> position in a 16mer, the 10<sup>th</sup> or 11<sup>th</sup> position in a 20 mer). A preferred ASO probe for detecting IL4R $\alpha$  gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

		•					
	TTGCACCACTGCACT	(SEQ	ID	NO:4)	and	its	complement,
	TTGCACCGCTGCACT	(SEQ	ID	NO:5)	and	its	complement,
	TTTTGTGCTATTCCC	(SEO	ID	NO: 6)	and	its	complement,
30	TTTTGTGTTATTCCC			-			complement,
	CTGGGCCGCTCAGGC	(SEO	TD	NO:8)	and	its	complement,
	CTGGGCCTCTCAGGC			•			complement,
					-		•
35	TAAGCCTGCGCTGGA	(SEQ	ID	NO:10)	and	its	complement,
	TAAGCCTCCGCTGGA	(SEQ	ID	NO:11)	and	its	complement,
	AGAACAACGGAGGCG	(SEQ	ID	NO:12)	and	its	complement,
	AGAACAATGGAGGCG	(SEQ	ID	NO:13)	and	its	complement,
40							•
	CGGAGGCGCGGGTG	(SEQ	ID	NO:14)	and	its	complement,
• 0	CGGAGGCACGGGGTG	(SEQ	ID	NO:15)	and	its	complement,

	GTGCGGATAACTATA GTGCGGACAACTATA		NO:16) NO:17)		complement, complement,
5	CGGAGTGCGGCAGGG CGGAGTGTGGCAGGG		NO:18) NO:19)		complement, complement,
·	GCCTGGGCTGAGGGT GCCTGGGTTGAGGGT		NO:20) NO:21)		complement, complement,
10 .	TGGGGTGGGCAGGGG TGGGGTGAGCAGGGG		NO:22) NO:23)		complement,
15	TTCTCCCGCAGTGAA TTCTCCCACAGTGAA	•	NO:24) NO:25)		complement,
	GTGAAAACGACCCGG GTGAAAATGACCCGG		NO:26) NO:27)		complement, complement,
20	GGCAAGCCCTGGGGC GGCAAGCTCTGGGGC		NO:28) NO:29)		complement, complement,
	GCCCTGGGGCTGGAT GCCCTGGAGCTGGAT				complement, complement,
25	ATAGCAAATCCCAGG ATAGCAATTCCCAGG		NO: 32) NO: 33)		complement, complement,
30	GCTCTGCCCTAGGCA GCTCTGCACTAGGCA				complement,
	CCCCCACCCCTCACA CCCCCACTCCTCACA				complement,
35	TCCCTCCGCATCGCA TCCCTCCACATCGCA		NO:38)		complement, complement,
· .	CACCTGCTGTGGTGT CACCTGCCGTGGTGT		NO:40) NO:41)		complement, complement,
40	ATGTCTGAAGTAGAC ATGTCTGCAGTAGAC				complement, complement,
45	TGACCAACCTTTGCT TGACCAACTTTTGCT				complement,
.5	CCTGTTTTCTGGAGC CCTGTTTCCTGGAGC				complement, complement,
50	TGGACCTGCTCGGAG TGGACCTTCTCGGAG				complement, complement,
	AGTCATGCCTTCTTC AGTCATGTCTTCTTC				complement, complement,
55	GCCTTCTTCCACCTT GCCTTCTCCCACCTT				complement, complement,

```
CAGCCCCCGTCTCGG (SEQ ID NO:54) and its complement,
          CAGCCCCTGTCTCGG (SEQ ID NO:55) and its complement,
          GGAGTTTGTACATGC (SEQ ID NO: 56) and its complement,
 5
          GGAGTTTATACATGC (SEQ ID NO:57) and its complement,
          CAGCTCCCCAGAGCA (SEQ ID NO:58) and its complement,
          CAGCTCCTCAGAGCA (SEQ ID NO:59) and its complement,
          AGACAGGTCCTCGCC (SEQ ID NO:60) and its complement,
10
          AGACAGGGCCTCGCC (SEQ ID NO:61) and its complement,
          CTGCCCCTGGCAATG (SEQ ID NO:62) and its complement,
          CTGCCCCCGGCAATG (SEQ ID NO:63) and its complement,
15
          AGGTGCATGTCCTCT
                           (SEQ ID NO:64) and its complement,
          AGGTGCACGTCCTCT
                          (SEQ ID NO:65) and its complement,
          GTGCATGTCCTCTTG (SEQ ID NO:66) and its complement,
20
          GTGCATGCCCTCTTG (SEQ ID NO: 67) and its complement,
          GGCTTATCCATGCCT (SEQ ID NO:68) and its complement,
                          (SEQ ID NO:69) and its complement,
          GGCTTATTCATGCCT
25
          AGCCAGGCTGGCAGA (SEQ ID NO:70) and its complement,
          AGCCAGGGTGGCAGA
                          (SEQ ID NO:71) and its complement,
          GGCCCACATGGAGGC
                          (SEQ ID NO:72) and its complement,
          GGCCCACGTGGAGGC
                          (SEQ ID NO:73) and its complement,
30
          TAACACAGCCATCAA (SEQ ID NO:74) and its complement,
          TAACACAACCATCAA
                          (SEQ ID NO:75) and its complement,
          TAATGCTCGTCTGTG
                           (SEQ ID NO:76) and its complement,
35
                           (SEQ ID NO:77) and its complement,
          TAATGCTTGTCTGTG
                          (SEQ ID NO:78) and its complement, and
          ACTTGCCGTCTGGGT
          ACTTGCCATCTGGGT
                          (SEQ ID NO:79) and its complement.
```

40

An allele-specific oligonucleotide primer of the invention has a 3' terminal nucleotide, or preferably a 3' penultimate nucleotide, that is complementary to only one nucleotide of a particular SNP, thereby acting as a primer for polymerase-mediated extension only if the allele containing that nucleotide is present. Allele-specific oligonucleotide primers hybridizing to either the coding or noncoding strand are contemplated by the invention. A preferred ASO primer for detecting ILARα gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

CTGAGATTGCACCAC (SEQ ID NO:80); GGCTGGAGTGCAGTG (SEQ ID NO:81);

CTGAGATTGCACCGC (SEQ ID NO:82); GGCTGGAGTGCAGCG (SEQ ID NO:83);

CTGTGCTTTTGTGCT (SEQ ID NO:84); ACCAAGGGGAATAGC (SEQ ID NO:85);

CTGTGCTTTTGTGTT (SEQ ID NO:86); ACCAAGGGGAATAAC (SEQ ID NO:87);

```
GAGTTCCTGGGCCGC (SEQ ID NO:88); GGAGCAGCCTGAGCG (SEQ ID NO:89);
     GAGTTCCTGGGCCTC (SEQ ID NO:90); GGAGCAGCCTGAGAG (SEQ ID NO:91);
     TCCGAGTAAGCCTGC (SEQ ID NO:92); TCCAGCTCCAGCGCA (SEQ ID NO:93);
     TCCGAGTAAGCCTCC (SEQ ID NO:94); TCCAGCTCCAGCGGA (SEQ ID NO:95);
     TCCCTGAGAACAACG (SEQ ID NO:96); ACCCCGCGCCTCCGT (SEQ ID NO:97);
     TCCCTGAGAACAATG (SEQ ID NO:98); ACCCCGCGCCTCCAT (SEQ ID NO:99);
10
     GAACAACGGAGGCGC (SEQ ID NO:100); CACACGCACCCCGCG (SEQ ID NO:101);
     GAACAACGGAGGCAC (SEQ ID NO:102); CACACGCACCCCGTG (SEQ ID NO:103);
     TGGTCAGTGCGGATA (SEQ ID NO:104); CCAGTGTATAGTTAT (SEQ ID NO:105);
     TGGTCAGTGCGGACA (SEQ ID NO:106); CCAGTGTATAGTTGT (SEQ ID NO:107);
15
     GCAGGGCGGAGTGCG (SEQ ID NO:108); AGCCACCCTGCCGC (SEQ ID NO:109);
     GCAGGGCGGAGTGTG (SEO ID NO:110); AGCCACCCTGCCAC (SEO ID NO:111);
    - ACAGCTGCCTGGGCT (SEQ ID NO:112); CACCCCACCCTCAGC (SEQ ID NO:113);
     ACAGCTGCCTGGGTT (SEQ ID NO:114); CACCCCACCCTCAAC (SEQ ID NO:115);
20
     TGAGGGTGGGGTGGG (SEQ ID NO:116); CCTCCTCCCCTGCCC (SEQ ID NO:117);
     TGAGGGTGGGGTGAG (SEQ ID NO:118); CCTCCTCCCCTGCTC (SEQ ID NO:119);
     GGCCGCTTCTCCCGC (SEQ ID NO:120); CTGGGTTTCACTGCG (SEQ ID NO:121);
25
     GGCCGCTTCTCCCAC (SEQ ID NO:122); CTGGGTTTCACTGTG (SEQ ID NO:123);
     TTTGGAGTGAAAACG (SEQ ID NO:124); CATCTGCCGGGTCGT (SEQ ID NO:125);
     TTTGGAGTGAAAATG (SEQ ID NO:126); CATCTGCCGGGTCAT (SEQ ID NO:127);
30
     CTGGGAGGCAAGCCC (SEQ ID NO:128); TATCCAGCCCCAGGG (SEQ ID NO:129);
     CTGGGAGGCAAGCTC (SEQ ID NO:130); TATCCAGCCCCAGAG (SEQ ID NO:131);
     AGGCAAGCCCTGGGG (SEQ ID NO:132); TTTGCTATCCAGCCC (SEQ ID NO:133);
     AGGCAAGCCCTGGAG (SEQ ID NO:134); TTTGCTATCCAGCTC (SEQ ID NO:135);
35
    GGCTGGATAGCAAAT (SEQ ID NO:136); CTAGCTCCTGGGATT (SEQ ID NO:137);
     GGCTGGATAGCAATT (SEQ ID NO:138); CTAGCTCCTGGGAAT (SEQ ID NO:139);
     CACCTGGCTCTGCCC (SEQ ID NO:140); GGGACTTGCCTAGGG (SEQ ID NO:141);
40
     CACCTGGCTCTGCAC (SEQ ID NO:142); GGGACTTGCCTAGTG (SEQ ID NO:143);
     CCTGGCCCCCACCC (SEQ ID NO:144); CTCTGATGTGAGGGG (SEQ ID NO:145);
     CCTGGCCCCCACTC (SEQ ID NO:146); CTCTGATGTGAGGAG (SEQ ID NO:147);
45
     GAACCCTCCCTCCGC (SEQ. ID NO:148); GCTGGCTGCGATGCG (SEQ. ID NO:149);
     GAACCCTCCCTCCAC (SEQ ID NO:150); GCTGGCTGCGATGTG (SEQ ID NO:151);
     TAGATACACCTGCTG (SEQ ID NO:152); GCAGATACACCACAG (SEQ ID NO:153);
 50
     TAGATACACCTGCCG (SEQ ID NO:154); GCAGATACACCACGG (SEQ ID NO:155);
     GAAGGCATGTCTGAA (SEQ ID NO:156); ATGGCTGTCTACTTC (SEQ ID NO:157);
     GAAGGCATGTCTGCA (SEQ ID NO:158); ATGGCTGTCTACTGC (SEQ ID NO:159);
     GAACCCTGACCAACC (SEQ ID NO:160); TGCAAAAGCAAAGGA (SEQ ID NO:161);
 55
     GAACCCTGACCAATC (SEQ ID NO:162); TGCAAAAGCAAAGAA (SEQ ID NO:163);
```

WO 01/04270 PCT/US00/19094.

```
TCTTGCCCTGTTTTC (SEQ ID NO:164); TGTTGTGCTCCAGAA (SEQ ID NO:165);
    TCTTGCCCTGTTTCC (SEQ ID NO:166); TGTTGTGCTCCAGGA (SEQ ID NO:167);
    TGTTCCTGGACCTGC (SEQ ID NO:168); TCTCCTCTCCGAGCA (SEQ ID NO:169);
    TGTTCCTGGACCTTC (SEQ ID NO:170); TCTCCTCTCCGAGAA (SEQ ID NO:171);
    TGGGGGAGTCATGCC (SEQ ID NO:172); AAGGTGGAAGAAGGC (SEQ ID NO:173);
    TGGGGGAGTCATGTC (SEQ ID NO:174); AAGGTGGAAGAAGAC (SEO ID NO:175);
10
    AGTCATGCCTTCTTC (SEQ ID NO:176); TTCCCGAAGGTGGAA (SEQ ID NO:177);
    AGTCATGCCTTCTCC (SEQ ID NO:178); TTCCCGAAGGTGGGA (SEQ ID NO:179);
    CAGCTGCAGCCCCG (SEQ ID NO:180); TGGGGGCCGAGACGG (SEQ ID NO:181);
    CAGCTGCAGCCCTG (SEQ ID NO:182); TGGGGGCCGAGACAG (SEQ ID NO:183);
15
    CTATCAGGAGTTTGT (SEQ ID NO:184); TCCACCGCATGTACA (SEQ ID NO:185);
    CTATCAGGAGTTTAT (SEQ ID NO:186); TCCACCGCATGTATA (SEQ ID NO:187);
    CCCAAGCAGCTCCCC (SEQ ID NO:188); CCCAGGTGCTCTGGG (SEQ ID NO:189);
20
    CCCAAGCAGCTCCTC (SEQ ID NO:190); CCCAGGTGCTCTGAG (SEQ ID NO:191);
    CTGTGGAGACAGGTC (SEQ ID NO:192); GTAGGGGGGGAGGAC (SEQ ID NO:193);
    CTGTGGAGACAGGGC (SEQ ID NO:194); GTAGGGGGGCGAGGCC (SEQ ID NO:195);
25
    TCCATCCTGCCCCTG (SEQ ID NO:196); TCTGAGCATTGCCAG (SEQ ID NO:197);
    TCCATCCTGCCCCG (SEQ ID NO:198); TCTGAGCATTGCCGG (SEQ ID NO:199);
    TCTCTTAGGTGCATG (SEQ ID NO:200); GCAACAAGAGGACAT (SEQ ID NO:201);
    TCTCTTAGGTGCACG (SEQ ID NO:202); GCAACAAGAGGACGT (SEQ ID NO:203);
30
    TCTTAGGTGCATGTC (SEQ ID NO: 204); CAGCAACAAGAGGAC (SEQ ID NO: 205);
    TCTTAGGTGCATGCC (SEQ ID NO:206); CAGCAACAAGAGGGC (SEQ ID NO:207);
    GACTAGGGCTTATCC (SEQ ID NO:208); TTTCCCAGGCATGGA (SEQ ID NO:209);
35
    GACTAGGGCTTATTC (SEQ ID NO:210); TTTCCCAGGCATGAA (SEQ ID NO:211);
    GAAGGCAGCCAGGCT (SEQ ID NO:212); TGGAAATCTGCCAGC (SEQ ID NO:213);
    GAAGGCAGCCAGGGT (SEQ ID NO:214); TGGAAATCTGCCACC (SEQ ID NO:215);
40 GATCATGGCCCACAT (SEQ ID NO:216); AGGTGGGCCTCCATG (SEQ ID NO:217);
    GATCATGGCCCACGT (SEQ ID NO:218); AGGTGGGCCTCCACG (SEQ ID NO:219);
    AGAAACTAACACGC (SEO ID NO:220); ATTCCCTTGATGGCT (SEO ID NO:221);
    AGAAACTAACACA (SEQ ID NO: 222); ATTCCCTTGATGGTT (SEQ ID NO: 223);
45
    GTTGAGTAATGCTCG (SEQ ID NO:224); AAAACACACAGACGA (SEQ ID NO:225);
    GTTGAGTAATGCTTG (SEQ ID NO:226); AAAACACACAGACAA (SEQ ID NO:227);
    TAAGAAACTTGCCGT (SEQ ID NO:228); ACCCAAACCCAGACG (SEQ ID NO:229);
    TAAGAAACTTGCCAT (SEQ ID NO:230); and ACCCAAACCCAGATG (SEQ ID NO:231).
50
```

Other genotyping oligonucleotides of the invention hybridize to a target region located one to several nucleotides downstream of one of the novel polymorphic sites identified herein. Such oligonucleotides are useful in polymerase-mediated primer extension methods for detecting one of the

novel polymorphisms described herein and therefore such genotyping oligonucleotides are referred to herein as "primer-extension oligonucleotides". In a preferred embodiment, the 3'-terminus of a primerextension oligonucleotide is a deoxynucleotide complementary to the nucleotide located immediately adjacent to the polymorphic site. A particularly preferred oligonucleotide primer for detecting IL4Ra gene polymorphisms by primer extension terminates in a nucleotide sequence, listed 5' to 3', selected

from the group consisting of: AGATTGCACC (SEQ ID NO:232); TGGAGTGCAG (SEQ ID NO:233); TGCTTTTGTG (SEQ ID NO: 234); AAGGGGAATA (SEQ ID NO:235); TTCCTGGGCC (SEQ ID NO: 236); · GCAGCCTGAG (SEQ ID NO:237); 10 GAGTAAGCCT (SEQ ID NO:238); AGCTCCAGCG (SEQ ID NO:239); CTGAGAACAA (SEQ ID NO:240); CCGCGCCTCC (SEQ ID NO:241); ACGCACCCCG CAACGGAGGC ' (SEQ ID NO:242); (SEQ ID NO:243); TCAGTGCGGA (SEQ ID NO: 244); GTGTATAGTT (SEQ ID NO:245); GGGCGGAGTG (SEQ ID NO:246); CACCCTGCC (SEQ ID NO:247); 15 GCTGCCTGGG (SEQ ID NO:248); CCCACCCTCA (SEQ ID NO:249); GGGTGGGGTG (SEQ ID NO:250); CCTCCCCTGC (SEQ ID NO:251); (SEQ ID NO:252); CGCTTCTCCC GGTTTCACTG (SEQ ID NO:253); (SEQ ID NO: 254); **GGAĠTGAAAA** CTGCCGGGTC (SEQ ID NO:255); GGAGGCAAGC (SEQ ID NO:256); CCAGCCCCAG (SEQ ID NO:257); 20 CAAGCCCTGG (SEQ ID NO:258); GCTATCCAGC (SEQ ID NO:259); TGGATAGCAA (SEQ ID NO:260); GCTCCTGGGA (SEQ ID NO:261); CTGGCTCTGC (SEQ ID NO:262); ACTTGCCTAG (SEQ ID NO:263); GGCCCCCAC (SEQ ID NO:264); TGATGTGAGG (SEQ ID NO:265); CCCTCCCTCC (SEQ ID NO: 266); GGCTGCGATG (SEQ ID NO:267); 25 **ATACACCTGC** (SEQ ID NO: 268); GATACACCAC (SEQ ID NO:269); **GGCATGTCTG** (SEQ ID NO:270); GCTGTCTACT (SEQ ID NO:271); CCCTGACCAA (SEQ ID NO: 272); AAAAGCAAAG (SEQ ID NO:273); TGCCCTGTTT (SEQ ID NO: 274); TGTGCTCCAG (SEQ ID NO:275); TCCTGGACCT (SEQ ID NO: 276); CCTCTCCGAG (SEQ ID NO:277); 30 GGGAGTCATG (SEQ ID NO: 278); GTGGAAGAAG (SEQ ID NO:279); CATGCCTTCT (SEQ ID NO:280); CCGAAGGTGG (SEQ ID NO:281); CTGCAGCCCC (SEQ ID NO:282); **GGGCCGAGAC** (SEQ ID NO:283); TCAGGAGTTT (SEQ ID NO:284); ACCGCATGTA . (SEQ ID NO:285); AAGCAGCTCC (SEQ ID NO:286); AGGTGCTCTG (SEQ ID NO:287); 35 (SEQ ID NO:288); TGGAGACAGG GGGGGCGAGG (SEQ ID NO:289); ATCCTGCCCC (SEQ ID NO:290); GAGCATTGCC (SEQ ID NO:291); **CTTAGGTGCA** (SEQ ID NO:292);

ACAAGAGGAC

(SEQ ID NO:293);

```
TAGGTGCATG
             (SEQ ID NO:294);
                                  CAACAAGAGG
                                                (SEQ ID NO:295);
TAGGGCTTAT
             (SEQ ID NO:296);
                                  CCCAGGCATG
                                                (SEQ ID NO:297);
GGCAGCCAGG
             (SEQ ID NO:298);
                                  AAATCTGCCA
                                                (SEQ ID NO:299);
CATGGCCCAC
             (SEQ ID NO:300);
                                  TGGGCCTCCA ·
                                                (SEQ ID NO:301);
AACTAACACA
             (SEQ ID NO:302);
                                                (SEQ ID NO:303);
                                  CCCTTGATGG
             (SEQ ID NO:304);
GAGTAATGCT
                                  ACACACAGAC
                                                (SEQ ID NO:305);
            (SEQ ID NO: 306); and CAAACCCAGA
GAAACTTGCC
                                                (SEQ ID NO:307).
```

5

10

15

20

25

30

35

In some embodiments, a composition contains two or more differently labeled genotyping oligonucleotides for simultaneously probing the identity of nucleotides at two or more polymorphic sites. It is also contemplated that primer compositions may contain two or more sets of allele-specific primer pairs to allow simultaneous targeting and amplification of two or more regions containing a polymorphic site.

IL4Rα genotyping oligonucleotides of the invention may also be immobilized on or synthesized on a solid surface such as a microchip, bead, or glass slide (see, e.g., WO 98/20020 and WO 98/20019). Such immobilized genotyping oligonucleotides may be used in a variety of polymorphism detection assays, including but not limited to probe hybridization and polymerase extension assays. Immobilized IL4Rα genotyping oligonucleotides of the invention may comprise an ordered array of oligonucleotides designed to rapidly screen a DNA sample for polymorphisms in multiple genes at the same time.

In another embodiment, the invention provides a kit comprising at least two genotyping oligonucleotides packaged in separate containers. The kit may also contain other components such as hybridization buffer (where the oligonucleotides are to be used as a probe) packaged in a separate container. Alternatively, where the oligonucleotides are to be used to amplify a target region, the kit may contain, packaged in separate containers, a polymerase and a reaction buffer optimized for primer extension mediated by the polymerase, such as PCR.

The above described oligonucleotide compositions and kits are useful in methods for genotyping and/or haplotyping the ILAR $\alpha$  gene in an individual. As used herein, the terms "ILAR $\alpha$  genotype" and "ILAR $\alpha$  haplotype" mean the genotype or haplotype contains the nucleotide pair or nucleotide, respectively, that is present at one or more of the novel polymorphic sites described herein and may optionally also include the nucleotide pair or nucleotide present at one or more additional polymorphic sites in the ILAR $\alpha$  gene. The additional polymorphic sites may be currently known polymorphic sites or sites that are subsequently discovered.

One embodiment of the genotyping method involves isolating from the individual a nucleic acid mixture comprising the two copies of the ILAR $\alpha$  gene, or a fragment thereof, that are present in the individual, and determining the identity of the nucleotide pair at one or more of the polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34. PS35. PS36.

PS37, PS38, PS39, PS40, PS41, PS44, and PS45 in the two copies to assign an IL4R $\alpha$  genotype to the individual. As will be readily understood by the skilled artisan, the two "copies" of a gene in an individual may be the same allele or may be different alleles. In a preferred embodiment of the genotyping method, the identity of the nucleotide pair atone or more of the polymorphic sites selected from the group consisting of PS5, PS24, PS26, PS29, PS31, PS42, and PS43 is also determined. In a particularly preferred embodiment, the genotyping method comprises determining the identity of the nucleotide pair at each of PS1-45.

Typically, the nucleic acid mixture is isolated from a biological sample taken from the individual, such as a blood sample or tissue sample. Suitable tissue samples include whole blood, semen saliva, tears, urine, fecal material, sweat, buccal, skin and hair. The nucleic acid mixture may be comprised of genomic DNA, mRNA, or cDNA and, in the latter two cases, the biological sample must be obtained from an organ in which the IL4Rα gene is expressed. Furthermore it will be understood by the skilled artisan that mRNA or cDNA preparations would not be used to detect polymorphisms located in introns or in 5′ and 3′ nontranscribed regions. If an IL4Rα gene fragment is isolated, it must contain the polymorphic site(s) to be genotyped.

10

15

20

25

30

35

One embodiment of the haplotyping method comprises isolating from the individual a nucleic acid molecule containing only one of the two copies of the ILARa gene, or a fragment thereof, that is present in the individual and determining in that copy the identity of the nucleotide at one or more of the polymorphic sites PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45 in that copy to assign an ILARα haplotype to the individual. The nucleic acid may be isolated using any method capable of separating the two copies of the II.4R\alpha gene or fragment such as one of the methods described above for preparing II.4R\alpha isogenes. with targeted in vivo cloning being the preferred approach. As will be readily appreciated by those skilled in the art, any individual clone will only provide haplotype information on one of the two ILARa gene copies present in an individual. If haplotype information is desired for the individual's other copy, additional ILARa clones will need to be examined. Typically, at least five clones should be examined to have more than a 90% probability of haplotyping both copies of the ILARQ gene in an individual. In some embodiments, the haplotyping method also comprises identifying the nucleotide atone or more of the polymorphic sites PS5, PS24, PS26, PS29, PS31, PS42, and PS43. In a particularly preferred embodiment, the nucleotide at each of PS1-45 is identified.

In a preferred embodiment, an II.4R $\alpha$  haplotype pair is determined for an individual by identifying the phased sequence of nucleotides at one or more of the polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45 in each copy of the II.4R $\alpha$  gene that is present in the individual. In a

particularly preferred embodiment, the haplotyping method comprises identifying the phased sequenc of nucleotides at each of PS1-45 in each copy of the ILARα gene. When haplotyping both copies of the gene, the identifying step is preferably performed with each copy of the gene being placed in separate containers. However, it is also envisioned that if the two copies are labeled with different tags, or are otherwise separately distinguishable or identifiable, it could be possible in some cases to perform the method in the same container. For example, if first and second copies of the gene are labeled with different first and second fluorescent dyes, respectively, and an allele-specific oligonucleotide labeled with yet a third different fluorescent dye is used to assay the polymorphic site(s), then detecting a combination of the first and third dyes would identify the polymorphism in the first gene copy while detecting a combination of the second and third dyes would identify the polymorphism in the second gene copy.

10

15

20

25

30

35

In both the genotyping and haplotyping methods, the identity of a nucleotide (or nucleotide pair) at a polymorphic site(s) may be determined by amplifying a target region(s) containing the polymorphic site(s) directly from one or both copies of the IL4Ra gene, or fragment thereof, and the sequence of the amplified region(s) determined by conventional methods. It will be readily appreciated by the skilled artisan that only one nucleotide will be detected at a polymorphic site in individuals who are homozygous at that site, while two different nucleotides will be detected if the individual is heterozygous for that site. The polymorphism may be identified directly, known as positive-type identification, or by inference, referred to as negative-type identification. For example, where a SNP is known to be guanine and cytosine in a reference population, a site may be positively determined to be either guanine or cytosine for an individual homozygous at that site, or both guanine and cytosine, if the individual is heterozygous at that site. Alternatively, the site may be negatively determined to be not guanine (and thus cytosine/cytosine) or not cytosine (and thus guanine/guanine).

In addition, the identity of the allele(s) present at any of the novel polymorphic sites described herein may be indirectly determined by genotyping a polymorphic site not disclosed herein that is in linkage disequilibrium with the polymorphic site that is of interest. Two sites are said to be in linkage disequilibrium if the presence of a particular variant at one site enhances the predictability of another variant at the second site (Stevens, JC 1999, *Mol. Diag.* 4: 309-17). Polymorphic sites in linkage disequilibrium with the presently disclosed polymorphic sites may be located in regions of the gene or in other genomic regions not examined herein. Genotyping of a polymorphic site in linkage disequilibrium with the novel polymorphic sites described herein may be performed by, but is not limited to, any of the above-mentioned methods for detecting the identity of the allele at a polymorphic site.

The target region(s) may be amplified using any oligonucleotide-directed amplification method, including but not limited to polymerase chain reaction (PCR) (U.S. Patent No. 4,965,188), ligase chain reaction (LCR) (Barany et al., Proc. Natl. Acad. Sci. USA 88:189-193, 1991; WO90/01069), and oligonucleotide ligation assay (OLA) (Land gren et al., Science 241:1077-1080, 1988). Oligonucleotides useful as prim rs or probes in such methods should specifically hybridize to a region of the nucleic acid

that contains or is adjacent to the polymorphic site. Typically, the oligonucleotides are between 10 and 35 nucleotides in length and preferably, between 15 and 30 nucleotides in length. Most preferably, the oligonucleotides are 20 to 25 nucleotides long. The exact length of the oligonucleotide will depend on many factors that are routinely considered and practiced by the skilled artisan.

Other known nucleic acid amplification procedures may be used to amplify the target region including transcription-based amplification systems (U.S. Patent No. 5,130,238; EP 329,822; U.S. Patent No. 5,169,766, WO89/06700) and isothermal methods (Walker et al., *Proc. Natl. Acad. Sci. USA* 89:392-396, 1992).

5

10

15 .

20

25

30

35

A polymorphism in the target region may also be assayed before or after amplification using one of several hybridization-based methods known in the art. Typically, allele-specific oligonucleotides are utilized in performing such methods. The allele-specific oligonucleotides may be used as differently labeled probe pairs, with one member of the pair showing a perfect match to one variant of a target sequence and the other member showing a perfect match to a different variant. In some embodiments, more than one polymorphic site may be detected at once using a set of allele-specific oligonucleotides or oligonucleotide pairs. Preferably, the members of the set have melting temperatures within 5°C, and more preferably within 2°C, of each other when hybridizing to each of the polymorphic sites being detected.

Hybridization of an allele-specific oligonucleotide to a target polynucleotide may be performed with both entities in solution, or such hybridization may be performed when either the oligonucleotide or the target polynucleotide is covalently or noncovalently affixed to a solid support. Attachment may be mediated, for example, by antibody-antigen interactions, poly-L-Lys, streptavidin or avidin-biotin, salt bridges, hydrophobic interactions, chemical linkages, UV cross-linking baking, etc. Allele-specific oligonucleotides may be synthesized directly on the solid support or attached to the solid support subsequent to synthesis. Solid-supports suitable for use in detection methods of the invention include substrates made of silicon, glass, plastic, paper and the like, which may be formed, for example, into wells (as in 96-well plates), slides, sheets, membranes, fibers, chips, dishes, and beads. The solid support may be treated, coated or derivatized to facilitate the immobilization of the allele-specific oligonucleotide or target nucleic acid.

The genotype or haplotype for the ILAR $\alpha$  gene of an individual may also be determined by hybridization of a nucleic sample containing one or both copies of the gene to nucleic acid arrays and subarrays such as described in WO 95/11995. The arrays would contain a battery of allele-specific oligonucleotides representing each of the polymorphic sites to be included in the genotype or haplotype.

The identity of polymorphisms may also be determined using a mismatch detection technique, including but not limited to the RNase protection method using riboprobes (Winter et al., Proc. Natl. Acad. Sci. USA 82:7575, 1985; Meyers et al., Science 230:1242, 1985) and proteins which recognize nucleotide mismatches, such as the E. coli mutS protein (Modrich, P. Ann. Rev. Genet. 25:229-253, 1991). Alternatively, variant alleles can be identified by single strand conformation polymorphism

(SSCP) analysis (Orita et al., Genomics 5:874-879, 1989; Humphries et al., in Molecular Diagnosis of Genetic Diseases, R. Elles, ed., pp. 321-340, 1996) or denaturing gradient gel electrophoresis (DGGE) (Wartell et al., Nucl. Acids Res. 18:2699-2706, 1990; Sheffield et al., Proc. Natl. Acad. Sci. USA 86:232-236, 1989).

5

10

15

20

25

30

35

A polymerase-mediated primer extension method may also be used to identify the polymorphism(s). Several such methods have been described in the patent and scientific literature and include the "Genetic Bit Analysis" method (WO92/15712) and the ligase/polymerase mediated genetic bit analysis (U.S. Patent 5,679,524. Related methods are disclosed in WO91/02087, WO90/09455, WO95/17676, U.S. Patent Nos. 5,302,509, and 5,945,283. Extended primers containing a polymorphism may be detected by mass spectrometry as described in U.S. Patent No. 5,605,798. Another primer extension method is allele-specific PCR (Ruaño et al., Nucl. Acids Res. 17:8392, 1989; Ruaño et al., Nucl. Acids Res. 19, 6877-6882, 1991; WO 93/22456; Turki et al., J. Clin. Invest. 95:1635-1641, 1995). In addition, multiple polymorphic sites may be investigated by simultaneously amplifying multiple regions of the nucleic acid using sets of allele-specific primers as described in Wallace et al. (WO89/10414).

In another aspect of the invention, an individual's IL4Rα haplotype pair is predicted from its IL4Rα genotype using information on haplotype pairs known to exist in a reference population. In its broadest embodiment, the haplotyping prediction method comprises identifying an IL4Rα genotype for the individual at two or more polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45, enumerating all possible haplotype pairs which are consistent with the genotype, accessing data containing IL4Rα haplotype pairs identified in a reference population, and assigning a haplotype pair to the individual that is consistent with the data. In one embodiment, the reference haplotype pairs include the IL4Rα haplotype pairs shown in Table 4.

Generally, the reference population should be composed of randomly-selected individuals representing the major ethnogeographic groups of the world. A preferred reference population for use in the methods of the present invention comprises an approximately equal number of individuals from Caucasian, African American, Asian and Hispanic-Latino population groups with the minimum number of each group being chosen based on how rare a haplotype one wants to be guaranteed to see. For example, if one wants to have a q% chance of not missing a haplotype that exists in the population at a p% frequency of occurring in the reference population, the number of individuals (n) who must be sampled is given by  $2n=\log(1-q)/\log(1-p)$  where p and q are expressed as fractions. A preferred reference population allows the detection of any haplotype whose frequency is at least 10% with about 99% certainty and comprises about 20 unrelated individuals from each of the four population groups named above. A particularly preferred reference population includes a 3-generation family representing one or more of the four population groups to serve as controls for checking quality of haplotyping procedures.

In a preferred embodiment, the haplotype frequency data for each ethnogeographic group is examined to determine whether it is consistent with Hardy-Weinberg equilibrium. Hardy-Weinberg equilibrium (D.L. Hartl et al., Principles of Population Genomics, Sinauer Associates (Sunderland, MA),  $3^{rd}$  Ed., 1997) postulates that the frequency of finding the haplotype pair  $H_1/H_2$  is equal to  $p_{H-W}(H_1/H_2) = 2p(H_1)p(H_2)$  if  $H_1 \neq H_2$  and  $p_{H-W}(H_1/H_2) = p(H_1)p(H_2)$  if  $H_1 = H_2$ . A statistically significant difference between the observed and expected haplotype frequencies could be due to one or more factors including significant inbreeding in the population group, strong selective pressure on the gene, sampling bias, and/or errors in the genotyping process. If large deviations from Hardy-Weinberg equilibrium are observed in an ethnogeographic group, the number of individuals in that group can be increased to see if the deviation is due to a sampling bias. If a larger sample size does not reduce the difference between observed and expected haplotype pair frequencies, then one may wish to consider haplotyping the individual using a direct haplotyping method such as, for example, CLASPER System technology (U.S. Patent No. 5,866,404), SMD, or allele-specific long-range PCR (Michalotos-Beloin et al., Nucleic Acids Res. 24:4841-4843, 1996).

10

15

20

25

30

35

In one embodiment of this method for predicting an IL4Rα haplotype pair, the assigning step involves performing the following analysis. First, each of the possible haplotype pairs is compared to the haplotype pairs in the reference population. Generally, only one of the haplotype pairs in the reference population matches a possible haplotype pair and that pair is assigned to the individual. Occasionally, only one haplotype represented in the reference haplotype pairs is consistent with a possible haplotype pair for an individual, and in such cases the individual is assigned a haplotype pair containing this known haplotype and a new haplotype derived by subtracting the known haplotype from the possible haplotype pair. In rare cases, either no haplotypes in the reference population are consistent with the possible haplotype pairs, or alternatively, multiple reference haplotype pairs are consistent with the possible haplotype pairs. In such cases, the individual is preferably haplotyped using a direct molecular haplotyping method such as, for example, CLASPER System technology (U.S. Patent No. 5,866,404), SMD, or allele-specific long-range PCR (Michalotos-Beloin et al., Nucleic Acids Res. 24:4841-4843, 1996).

The invention also provides a method for determining the frequency of an IL4Rα genotype or IL4Rα haplotype in a population. The method comprises determining the genotype or the haplotype pair for the IL4Rα gene that is present in each member of the population, wherein the genotype or haplotype comprises the nucleotide pair or nucleotide detected at one or more of the polymorphic sites PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45 in the IL4Rα gene; and calculating the frequency any particular genotype or haplotype is found in the population. The population may be a reference population, a family population, a same sex population, a population group, a trait population (e.g., a group of individuals exhibiting a trait

of interest such as a medical condition or response to a therapeutic treatment).

5

10

· 15

20

25

.30

35

In another aspect of the invention, frequency data for ILAR a genotypes and/or haplotypes found in a reference population are used in a method for identifying an association between a trait and an ILARa. genotype or an IL4Ra haplotype. The trait may be any detectable phenotype, including but not limited to susceptibility to a disease or response to a treatment. The method involves obtaining data on the frequency of the genotype(s) or haplotype(s) of interest in a reference population as well as in a population exhibiting the trait. Frequency data for one or both of the reference and trait populations may be obtained by genotyping or haplotyping each individual in the populations using one of the methods described above. The haplotypes for the trait population may be determined directly or, alternatively, by the predictive genotype to haplotype approach described above. In another embodiment, the frequency data for the reference and/or trait populations is obtained by accessing previously determined frequency data, which may be in written or electronic form. For example, the frequency data may be present in a database that is accessible by a computer. Once the frequency data is obtained, the frequencies of the genotype(s) or haplotype(s) of interest in the reference and trait populations are compared. In a preferred embodiment, the frequencies of all genotypes and/or haplotypes observed in the populations are compared. If a particular genotype or haplotype for the IL4Ra gene is more frequent in the trait population than in the reference population at a statistically significant amount, then the trait is predicted to be associated with that ILARa genotype or haplotype. Preferably, the ILARa genotype or haplotype being compared in the trait and reference populations is selected from the full-genotypes and fullhaplotypes shown in Tables 4 and 5, respectively, or from sub-genotypes and sub-haplotypes derived from these genotypes and haplotypes.

In a preferred embodiment of the method, the trait of interest is a clinical response exhibited by a patient to some therapeutic treatment, for example, response to a drug targeting ILAR $\alpha$  or response to a therapeutic treatment for a medical condition. As used herein, "medical condition" includes but is not limited to any condition or disease manifested as one or more physical and/or psychological symptoms for which treatment is desirable, and includes previously and newly identified diseases and other disorders. As used herein the term "clinical response" means any or all of the following: a quantitative measure of the response, no response, and adverse response (i.e., side effects).

In order to deduce a correlation between clinical response to a treatment and an ILAR $\alpha$  genotype or haplotype, it is necessary to obtain data on the clinical responses exhibited by a population of individuals who received the treatment, hereinafter the "clinical population". This clinical data may be obtained by analyzing the results of a clinical trial that has already been run and/or the clinical data may be obtained by designing and carrying out one or more new clinical trials. As used herein, the term "clinical trial" means any research study designed to collect clinical data on responses to a particular treatment, and includes but is not limited to phase I, phase II and phase III clinical trials. Standard methods are used to define the patient population and to enroll subjects.

It is preferred that the individuals included in the clinical population have been graded for the existence of the medical condition of interest. This is important in cases where the symptom(s) being presented by the patients can be caused by more than one underlying condition, and where treatment of the underlying conditions are not the same. An example of this would be where patients experience breathing difficulties that are due to either asthma or respiratory infections. If both sets were treated with an asthma medication, there would be a spurious group of apparent non-responders that did not actually have asthma. These people would affect the ability to detect any correlation between haplotype and treatment outcome. This grading of potential patients could employ a standard physical exam or one or more lab tests. Alternatively, grading of patients could use haplotyping for situations where there is a strong correlation between haplotype pair and disease susceptibility or severity.

5

10

15

20

25

30

35

The therapeutic treatment of interest is administered to each individual in the trial population and each individual's response to the treatment is measured using one or more predetermined criteria. It is contemplated that in many cases, the trial population will exhibit a range of responses and that the investigator will choose the number of responder groups (e.g., low, medium, high) made up by the various responses. In addition, the ILAR $\alpha$  gene for each individual in the trial population is genotyped and/or haplotyped, which may be done before or after administering the treatment.

After both the clinical and polymorphism data have been obtained, correlations between individual response and ILAR $\alpha$  genotype or haplotype content are created. Correlations may be produced in several ways. In one method, individuals are grouped by their ILAR $\alpha$  genotype or haplotype (or haplotype pair) (also referred to as a polymorphism group), and then the averages and standard deviations of clinical responses exhibited by the members of each polymorphism group are calculated.

These results are then analyzed to determine if any observed variation in clinical response between polymorphism groups is statistically significant. Statistical analysis methods which may be used are described in L.D. Fisher and G. vanBelle, "Biostatistics: A Methodology for the Health Sciences", Wiley-Interscience (New York) 1993. This analysis may also include a regression calculation of which polymorphic sites in the IL4R $\alpha$  gene give the most significant contribution to the differences in phenotype. One regression model useful in the invention starts with a model of the form

$$r = r_0 + S \times d$$

where r is the response,  $r_0$  is a constant called the "intercept", S is the slope and d is the dose. To determine the dose, the most-common and least common nucleotides at the polymorphic site are first defined. Then, for each individual in the trial population, one calculates a "dose" as the number of least-common nucleotides the individual has at the polymorphic site of interest. This value can be 0 (homozygous for the least-common nucleotide), 1 (heterozygous), or 2 (homozygous for the most common nucleotide). An individual's "response" is the value of the clinical measurement. Standard linear regression methods are then used to fit all the individuals' doses and responses to a single model (see e.g., L.D. Fisher and G. vanBelle, supra, Ch 9). The outputs of the regression calculation are the

intercept  $r_0$ , the slope S, and the variance (which measures how well the data fits this simple linear model). The Students t-test value and the level of significance can then be calculated for each of the polymorphic sites.

5

10

15

20

25

30

A second method for finding correlations between IL4Rα haplotype content and clinical responses uses predictive models based on error-minimizing optimization algorithms. One of many possible optimization algorithms is a genetic algorithm (R. Judson, "Genetic Algorithms and Their Uses in Chemistry" in Reviews in Computational Chemistry, Vol. 10, pp. 1-73, K. B. Lipkowitz and D. B. Boyd, eds. (VCH Publishers, New York, 1997). Simulated annealing (Press et al., "Numerical Recipes in C: The Art of Scientific Computing", Cambridge University Press (Cambridge) 1992, Ch. 10), neural networks (E. Rich and K. Knight, "Artificial Intelligence", 2<sup>nd</sup> Edition (McGraw-Hill, New York, 1991, Ch. 18), standard gradient descent methods (Press et al., supra Ch. 10), or other global or local optimization approaches (see discussion in Judson, supra) could also be used. As an example, a genetic algorithm approach is described herein. This method searches for optimal parameters or weights in linear or non-linear models connecting IL4Rα haplotype loci and clinical outcome. One model is of the form

$$C = C_0 + \sum_{\alpha} \left( \sum_{i} w_{i,\alpha} R_{i,\alpha} + \sum_{i} w'_{i,\alpha} L_{i,\alpha} \right)$$
[1]

where C is the measured clinical outcome, i goes over all polymorphic sites,  $\alpha$  over all candidate genes,  $C_0$ ,  $w_{i,\alpha}$  and  $w'_{i,\alpha}$  are variable weight values,  $R_{i,\alpha}$  is equal to 1 if site i in gene  $\alpha$  in the first haplotype takes on the most common nucleotide and -1 if it takes on the less common nucleotide.  $L_{i,\alpha}$  is the same as  $R_{i,\alpha}$  except for the second haplotype. The constant term  $C_0$  and the weights  $w_{i,\alpha}$  and  $w'_{i,\alpha}$  are varied by the genetic algorithm during a search process that minimizes the error between the measured value of C and the value calculated from Equation 1. Models other than the one given in Equation 1 can be readily incorporated by those skilled in the art for analyzing the clinical and polymorphism data. The genetic algorithm is especially suited for searching not only over the space of weights in a particular model but also over the space of possible models (Judson, supra).

Correlations may also be analyzed using analysis of variation (ANOVA) techniques to determine how much of the variation in the clinical data is explained by different subsets of the polymorphic sites in the ILAR $\alpha$  gene. ANOVA is used to test hypotheses about whether a response variable is caused by or correlated with one or more traits or variables that can be measured (Fisher and vanBelle, supra, Ch. 10). These traits or variables are called the independent variables. To carry out ANOVA, the independent variable(s) are measured and individuals are placed into groups based on their values for these variables. In this case, the independent variable(s) refers to the combination of polymorphisms present at a subset of the polymorphic sites, and thus, each group contains those individuals with a given genotype or haplotype pair. The variation in response within the groups and also the variation between groups is then measured. If the within-group response variation is large (people in a group have a wide range of responses) and the

response variation between groups is small (the average responses for all groups are about the same) then it can be concluded that the independent variables used for the grouping are not causing or correlated with the response variable. For instance, if people are grouped by month of birth (which should have nothing to do with their response to a drug) the ANOVA calculation should show a low level of significance. However, if the response variation is larger between groups than within groups, the F-ratio (="between groups" divided by "within groups") is greater than one. Large values of the F-ratio indicate that the independent variable is causing or correlated with the response. The calculated F-ratio is preferably compared with the critical F-distribution value at whatever level of significance is of interest. If the F-ratio is greater than the Critical F-distribution value, then one may be confident that the individual's genotype or haplotype pair for this particular subset of polymorphic sites in the ILARα gene is at least partially responsible for, or is at least strongly correlated with the clinical response.

5

10

15

20

25

30

35

From the analyses described above, a mathematical model may be readily constructed by the skilled artisan that predicts clinical response as a function of ILAR a genotype or haplotype content. Preferably, the model is validated in one or more follow-up clinical trials designed to test the model.

The identification of an association between a clinical response and a genotype or haplotype (or haplotype pair) for the ILAR $\alpha$  gene may be the basis for designing a diagnostic method to determine those individuals who will or will not respond to the treatment, or alternatively, will respond at a lower level and thus may require more treatment, i.e., a greater dose of a drug. The diagnostic method may take one of several forms: for example, a direct DNA test (i.e., genotyping or haplotyping one or more of the polymorphic sites in the ILAR $\alpha$  gene), a serological test, or a physical exam measurement. The only requirement is that there be a good correlation between the diagnostic test results and the underlying ILAR $\alpha$  genotype or haplotype that is in turn correlated with the clinical response. In a preferred embodiment, this diagnostic method uses the predictive haplotyping method described above.

Any or all analytical and mathematical operations involved in practicing the methods of the present invention may be implemented by a computer. In addition, the computer may execute a program that generates views (or screens) displayed on a display device and with which the user can interact to view and analyze large amounts of information relating to the ILAR $\alpha$  gene and its genomic variation, including chromosome location, gene structure, and gene family, gene expression data, polymorphism data, genetic sequence data, and clinical data population data (e.g., data on ethnogeographic origin, clinical responses, genotypes, and haplotypes for one or more populations). The ILAR $\alpha$  polymorphism data described herein may be stored as part of a relational database (e.g., an instance of an Oracle database or a set of ASCII flat files). These polymorphism data may be stored on the computer's hard drive or may, for example, be stored on a CD ROM or on one or more other storage devices accessible by the computer. For example, the data may be stored on one or more databases in communication with the computer via a network.

Preferred embodiments of the invention are described in the following examples. Other

embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

5

#### **EXAMPLES**

The Examples herein are meant to exemplify the various aspects of carrying out the invention and are not intended to limit the scope of the invention in any way. The Examples do not include detailed descriptions for conventional methods employed, such as in the performance of genomic DNA isolation, PCR and sequencing procedures. Such methods are well-known to those skilled in the art and are described in numerous publications, for example, Sambrook, Fritsch, and Maniatis, "Molecular Cloning: A Laboratory Manual", 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, USA, (1989).

15 -

10

# Example 1A

This example illustrates examination of various regions of the ILARa gene for polymorphic sites using DNA from Index Repository IA.

## **Amplification of Target Regions**

The following target regions of the IL4Rα gene were amplified using the PCR primer pairs listed below, with the sequences presented in the 5' to 3' direction and nucleotide positions shown for each region corresponding to the indicated GenBank Accession No.

Accession Number: AC004525

Fragment 1

25 Forward Primer

32801-32822 CCACAGTCATCCCGACACTAGC (SEQ ID NO:308)

Reverse Primer

Complement of 33355-33334 TATTCCAGCCGTATCCATGTGC (SEQ ID NO:309)

PCR product 555 nt

30

Fragment 2

Forward Primer

35515-35536 CCTTGGTGCATGTGGTAAGAGG (SEQ ID NO:310)

Reverse Primer ·

35 Complement of 36068-36046 TTTCAAAGGTGGGAGGACTGAGG (SEQ ID NO:311)

PCR product 554 nt

Fragment 3

**Forward Primer** 

40 37031-37050 GCAGTGAGCTGGGATTGTGC (SEQ ID NO:312)

Reverse Primer

Complement of 37701-37679 AACTCCCCTTCTCTGATGTGAGG (SEQ ID NO:313)

PCR product 671 nt

Fragment 4 Forward Primer

43240-43262 TCACAGTTACAGAGGTGGCAAGC (SEQ ID NO:314)

Reverse Primer

Complement of 43727-43706 CTGCCTACCTGGCAGATACACC (SEQ ID NO:315) PCR product 488 nt

Fragment 5

Forward Primer

49553-49574. AGCTGTCACTCCACCTCCTTGG (SEQ ID NO:316) 10

Reverse Primer

Complement of 50036-50013 AAAGCCTCTGGTCTGCTAATGACC (SEQ ID NO:317)

PCR product 484 nt

15 Fragment 6

Forward Primer

51394-51415 GGGAGGAGATTCAGAGCACTCC (SEQ ID NO:318)

Reverse Primer

Complement of 51847-51826 CAGTCCACGTTTCCAGAACACC (SEQ ID NO:319)

20 PCR product 454 nt

Fragment 7

Forward Primer

52806-52826 GGCTTGGGATAATGGTGTTGC (SEQ ID NO:320)

25 **Reverse Primer** 

Complement of 53529-53507 TACTTCCCGAAGGTGGAAGAAGG (SEO ID NO:321)

PCR product 724 nt

Fragment 8

30 Forward Primer

53242-53265 CAGTGGAGATCAGCAAGACAGTCC (SEQ ID NO:322)

**Reverse Primer** 

Complement of 53807-53786 GGGCATCTCGGGTTCTACTTCC (SEQ ID NO:323)

PCR product 566 nt

35

Fragment 9

**Forward Primer** 

53522-53544 GGGAAGTACGAGTGCTCACATGC (SEQ ID NO:324)

Reverse Primer

40 Complement of 54110-54088 CTTATACCCCTCTTCCCCACTGC (SEQ ID NO:325)

PCR product 589 nt

Fragment 10

Forward Primer

45 53821-53841 TCTCTGAGCCAACCACTGTGC (SEQ ID NO:326)

Reverse Primer

Complement of 54359-54337 GGCTGAGTAGACAATGCCACTGC (SEQ ID NO:327)

PCR product 539 nt

50 Fragment 11

Forward Primer

54055-54076 CTGTGTCCCCAGAGAAATGTGG (SEO ID NO:328)

Reverse Primer

Complement of 54717-54695 GACTCAGCAACAAGAGGACATGC (SEQ ID NO:329)

PCR product 663 nt 55

```
Fragment 12
Forward Primer
54342-54365 GGCATTGTCTACTCAGCCCTTACC (SEQ ID NO:330)
Reverse Primer
Complement of 54986-54967 ACAAGTCGAGGTGCCCAAGG (SEQ ID NO:331)
PCR product 645 nt
```

Fragment 13

Forward Primer

54669-54693 CCCACATACATGAGGGTCTCTTAGG (SEQ ID NO:332)

Reverse Primer

Complement of 55270-55250 ATTCTGCCTCCAGCATCAACC (SEQ ID NO:333)

PCR product 602 nt

15 Fragment 14

**Forward Primer** 

55235-55258 AACAGAGCTTCCTTAGGTTGATGC (SEQ ID NO:334)

Reverse Primer

Complement of 55847-55825 CCTCAGTTCCCCACTACCTTAGC (SEQ ID NO:335)

20 PCR product 613 nt

These primer pairs were used in PCR reactions containing genomic DNA isolated from immortalized cell lines for each member of Index Repository IA. The PCR reactions were carried out under the following conditions:

30	Reaction volume  10 x Advantage 2 Polymerase reaction buffer (Clontech)  100 ng of human genomic DNA  10 mM dNTP  Advantage 2 Polymerase enzyme mix (Clontech)  Forward Primer (10 µM)  Reverse Primer (10 µM)  Water	= $20 \mu l$ = $2 \mu l$ = $1 \mu l$ = $0.4 \mu l$ = $0.4 \mu l$ = $0.4 \mu l$ = $15.6 \mu l$
35 .	Amplification profile: 94°C - 2 min. 1 cycle  94°C - 30 sec. 70°C - 45 sec. 72°C - 1 min.  1 cycle  10 cycles	:
	94°C - 30 sec. 64°C - 45 sec. 72°C - 1 min. 35 cycles	

# 45 Sequencing of PCR Products

The PCR products were purified by Solid Phase Reversible Immobilization using the protocol developed by the Whitehead Genome Center. A detailed protocol can be found at http://www.genome.wi.mit.edu/sequencing/protocols/pure/SPRI pcr.html.

Briefly, five µl of carboxyl coated magnetic beads (10 mg/ml) and 60 µl of HYB BUFFER (2.5M

NaCl/20% PEG 8000) were added to each PCR reaction mixture (20 µl). The reaction mixture was mixed well and incubated at room temperature (RT) for 10 min. The microtitre plate was placed on a magnet for 2 min and the beads washed twice with 150 µl of 70% EtOH. The beads were air dried for 2 min and the DNA was eluted in 25 µl of distilled water and incubated at RT for 5 min. The beads were magnetically separated and the supernatant removed for testing and sequencing.

The purified PCR products were sequenced in both directions using the primer sets described previously or those listed, in the 5' to 3' direction, below.

Accession Number: AC004525

Fragment 1

10 Forward Primer

32865-32882 GCGCTGGCCCTCAACTTT (SEQ ID NO:336)

**Reverse Primer** 

Complement of 33283-33264 GTCCCTGGAGATGGGACCTC (SEQ ID NO:337)

15 Fragment 2

Forward Primer

35598-35617 GCCCCCAGATCTGTCCTCAC (SEQ ID NO:338).

Reverse Primer

Complement of 36013-35994 GGAAAATACAGGCGGCTTCC (SEQ ID NO:339)

20

5

Fragment 3

**Forward Primer** 

37182-37203 GGCTCTGAATCTGTGTGGTGCT (SEQ ID NO:340)

**Reverse Primer** 

25 Complement of 37639-37620 AGCCAGGTGAGAAGCCAGGT (SEQ ID NO:341)

Fragment 4

Forward Primer

43266-43285 GGCCTGAACAGGACGAACAA (SEQ ID NO:342)

30 Reverse Primer

Complement of 43687-43668 GGCAGGATTGCCATTAGAGG (SEO ID NO:343)

Fragment 5

Forward Primer

35 49639-49660 TGAGTCAGTGGTTTGACCTCCA (SEQ ID NO:344)

Reverse Primer

Complement of 49999-49980 GCCTCTGTCTCCCCTGCAAC (SEO ID NO:345)

Fragment 6

40 Forward Primer

51423-51442 CCACTTTTGCCATCGACCAC (SEQ ID NO:346)

Reverse Primer

Complement of 51818-51799 CTGCCGTCCCTTGAAGGCTA (SEQ ID NO:347)

45 Fragment 7.

Forward Primer

52932-52953 CCCTACCCTCAGGGATTTCTCA (SEQ ID NO:348)

Reverse Primer

Complement of 53474-53455 CCCATTCTCCTCTCCGAGCA (SEQ ID NO:349)

50

Fragment 8

Forward Primer

53280-53299 ATCAGCGTGGTGCGATGTGT (SEQ ID NO:350)

**Reverse Primer** 

5 Complement of 53756-53737 ACCCAGCTCTCTGGGACACG (SEQ ID NO:351)

Fragment 9

Forward Primer

53548-53568 GGGATGAGTTCCCAAGTGCAG (SEQ ID NO:352)

10 Reverse Primer

Complement of 54048-54029 TGGCAAGCAGGCTTGAGAAG (SEQ ID NO:353)

Fragment 10

Forward Primer

15 53841-53860 CCCCAACCTGAGCCAGAAAC (SEQ ID NO:354)

Reverse Primer

Complement of 54330-54311 TGTCCACAAGGGGGTCTGTG (SEQ ID NO:355)

Fragment 11

20 Forward Primer

54083-54102 GGCTAGCAGTGGGGAAGAGG (SEQ ID NO:356)

Reverse Primer

Complement of 54617-54600 ATTGCCAGGGGCAGGATG (SEQ ID NO:357)

25 Fragment 12

Forward Primer

54422-54441 CCCTGTCATGGCCAGTCCTT (SEQ ID NO:358)

Reverse Primer

Complement of 54929-54910 GCGACCCAGTGCCCTCTACT (SEQ ID NO:359)

30

Fragment 13

**Forward Primer** 

54694-54715 TGCATGTCCTCTTGTTGCTGAG (SEQ ID NO:360)

Reverse Primer

35 Complement of 55213-55194 CAATGACCACCCTCCCTGAA (SEQ ID NO:361)

Fragment 14

Forward Primer

55273-55291 CGGCTGTCAAGGGGTGTTC (SEQ ID NO:362)

40 Reverse Primer

Complement of 55769-55750 CCAAACCCAGACGCAAGTT (SEQ ID NO:363)

### 45 Analysis of Sequences for Polymorphic Sites

Sequences were analyzed for the presence of polymorphisms using the Polyphred program (Nickerson et al., Nucleic Acids Res. 14:2745-2751, 1997). The presence of a polymorphism was confirmed on both strands. The polymorphisms and their locations in the ILAR $\alpha$  gene are listed in Table 3 below.

50

## Example 1B

This example illustrates examination of the ILARa gene for polymorphic sites in the following

target regions: 1000 base pairs upstream of the ATG start codon; each of the exons, including approximately 100 base pairs on either side of the exon; and 500-1000 base paris downstream of the termination codon.

#### 5 Amplification of the Target Regions

PCR primers, which were designed based on the nearly complete ILAR genomic sequence reported in the GenBank database (Accession No: AC004525), are set forth below:

**Promoter** 

Forward Primer

10 30094-30117 AAACCCCTGGACTCCAAGTGATCC (SEQ ID NO:364)

Reverse Primer

Complement of 30842-30819 AAGCGATTCTTCTGCCTCAGCCTC (SEQ ID NO:365)

PCR product 749 nt

15 Exon 1

Forward Primer

30517-30540 GGACAGTTGTTGTGTAGCTCACCC (SEQ ID NO:366)

Reverse Primer

Complement of 31409-31431 CTATGTTGCCCAAGCTGACCTC (SEQ ID NO:367)

20 PCR product 893 nt

Exon 2

Forward Primer

32871-32890 GCCCTCAACTTTGCCTGCAC (SEQ ID NO:368)

25 Reverse Primer

Complement of 33362-33340 AGTCCAGTATTCCAGCCGTATCC (SEQ ID NO:369)

PCR product. 492 nt

Exon 3

30 Forward Primer

35623-35644 TGATCGGGAAGCTGGAAGAGTC (SEQ ID NO:370)

Reverse Primer

Complement of 36070-36049 CGTTTCAAAGGTGGGAGGACTG (SEQ ID NO:371)

PCR product 448 nt

35

Exon 4

Forward Primer

37230-37250 CGACCAAAAATCTGGGTGGTG (SEQ ID NO:372)

Reverse Primer

40 Complement of 37668-37648 CAGGAAGCAAAGGGACTTGCC (SEQ ID NO:373)

PCR product 439 nt

Exon 5

Forward Primer

45 43307-43328 TCTTAAACATGGTGGGGTCAGC (SEQ ID NO:374)

Reverse Primer

Complement of 43765-43745 CATGGAAATTGTGGGCTTGTG (SEQ ID NO:375)

PCR product 459 nt

50 Exon 6

**Forward Primer** 

46507-46528 ATGTGCAAGAGGGAGAGTGGTG (SEO ID NO:376)

Reverse Primer

Complem nt of 46870-46849 TGACTGAGAGGACTGCAAAGGG (SEQ ID NO:377) PCR product 364 nt

5 Exon 7

**Forward Primer** 

49672-49695 GCCTGATCTCTGATGCCAAATAAG (SEQ ID NO:378).

**Reverse Primer** 

Complement of 49964-49983 TTTGCCATTCCAGAAGCCAG (SEQ ID NO:379)

10 PCR product 293 nt

Exon 8

Forward Primer

51524-51546 GATCTGTGTGATGTCGAGGCTTG (SEQ ID NO:380)

15 Reverse Primer

Complement of 51845-51825 GTCCACGTTTCCAGAACACCC (SEQ ID NO:381)

PCR product 322 nt

Exon 9 Fragment 1

20 Forward Primer

52912-52933 CGAAATCCCAAAGACACAGACC (SEQ ID NO:382)

Reverse Primer

Complement of 53722-53701 GAGTTGCTGAAGCTGCGGTAAG (SEQ ID NO:383)

PCR product 811 nt

25

Exon 9 Fragment 2

Forward Primer

53352-53374 GAAAAAGGGAGCTTCTGTGCATC (SEQ ID NO:384)

Reverse Primer

30 Complement of 54172-54153 AACAAGGGGACAGGGACTGG (SEQ ID NO:385)

PCR product 821 nt

Exon 9 Fragment 3

Forward Primer

35 53854-53874 CAGAAACCTGGGAGCAGATCC (SEQ ID NO:386)

Reverse Primer

Complement of 54711-54688 GCAACAAGAGGACATGCACCTAAG (SEQ ID NO:387)

PCR product 858 nt

40 Exon 9 Fragment 4

Forward Primer

54266-54288 AAAGGTAGAGGACATGCCAAAGC (SEQ ID NO:388)

Reverse Primer

Complement of 55007-54987 GGAGCAGCCAACAACTCGTTC (SEQ ID NO:389)

45 PCR product 742 nt

These primer pairs were used in PCR reactions containing genomic DNA isolated from immortalized cell lines for a reference population of 70 human individuals. The PCR reactions were carried out under the following conditions:

50Reaction volume= 20 μl $10 \times Advantage 2$  Polymerase reaction buffer (Clontech)= 2 μl100 ng of human genomic DNA= 1 μl10 mM dNTP= 0.4 μl

WO 01/042/0	
Advantage 2 Poly Forward Primer ( Reverse Primer ( Water	·
Amplification pro 94°C - 2 min. 94°C - 30 sec. 70°C - 45 sec. 72°C - 1 min.	1 cycle

35 cycles

# Sequencing of PCR Products

94°C - 30 sec. 64°C - 45 sec.

72°C - 1 min.

W/O 01/04270

5

10

15

20

25

The PCR products were purified by Solid Phase Reversible Immobilization using the protocol developed by the Whitehead Genome Center. A detailed protocol can be found at http://www.genome.wi.mit.edu/sequencing/protocols/pure/SPRI\_pcr.html.

PCT/US00/19094

=  $0.2 \mu l$ =  $0.4 \mu l$ =  $0.4 \mu l$ =  $15.6 \mu l$ 

Briefly, five  $\mu$ l of carboxyl coated magnetic beads (10 mg/ml) and 60  $\mu$ l of HYB BUFFER (2.5M NaCl/20% PEG 8000) were added to each PCR reaction mixture (20  $\mu$ l). The reaction mixture was mixed well and incubated at room temperature (RT) for 10 min. The microtitre plate was placed on a magnet for 2 min and the beads washed twice with 150  $\mu$ l of 70% EtOH. The beads were air dried for 2 min and the DNA was eluted in 25  $\mu$ l of distilled water and incubated at RT for 5 min. The beads were magnetically separated and the supernatant removed for testing and sequencing.

The purified PCR products were sequenced in both directions using the primer sets described previously or those listed, in the 5' to 3' direction, below.

30 Promoter Fragment 1 Forward Primer

30312-30334 GCTCATTTAATCCCCACAACACC (SEQ ID NO:390)

Reverse Primer

35 Complement of 30791-30769 CCACCACACCTGGCTAATTTTTG (SEQ ID NO:391)

Promoter Fragment 2 Forward Primer

30529-30551 TGTAGCTCACCCTCTGGACTTTG (SEQ ID NO:392)

40 Reverse Primer

Complement of 30990-30971 AATATGCAACCCTCCCCTGC (SEQ ID NO:393)

Exon 1

**Forward Primer** 

45 30824-30846 TGAGGCAGAAGAATCGCTTGAAC (SEQ ID NO:394)

Reverse Primer

Complement of 31261-31240 ACTTGTCATTGGCTGTCCCCTC (SEQ ID NO:395)

61

Exon 2

Forward Primer

32880-32900 TTTGCCTGCACTGTGCTTTTG (SEQ ID NO:396)

5 Reverse Primer

Complement of 33234-33212 CCATACTCAGCATCCTGCACTCC (SEQ ID NO:397)

Exon 5

Forward Primer

10 43330-43349 AACGACAGCAACCAGGGTGG (SEQ ID NO:398)

Reverse Primer

Complement of 43704-43682 CAGCAGGTGTATCTAATGGCAGG (SEQ ID NO:399)

Exon 6

15 Forward Primer

46520-46541 AGAGTGGTGGGGAGATGAGGTG (SEQ ID NO:400)

Reverse Primer

Complement of 46857-46837 TGCAAAGGGGCAGACTAGAGG (SEQ ID NO:401)

20 Exon 7

Forward Primer

49708-49729 CGACCACTTTTATGGGAGGAGC (SEQ ID NO:402)

Reverse Primer

Complement of 49927-49905 CCAGGTGTTCTGAACCACACTTC (SEQ ID NO:403)

25.

Exon 8

Forward Primer

51528-51550 TGTGTGATGTCGAGGCTTGTACC (SEQ ID NO:404)

Reverse Primer

30 Complement of 51779-51758 GAATGCAGGGAAGAGGCAG (SEQ ID NO:405)

Exon 9 Fragment 1

Forward Primer

53017-53038 GCCATCAGGACATGGTGATTTC (SEQ ID NO:406)

35 Reverse Primer

Complement of 53539-53518 TGAGCACTCGTACTTCCCGAAG (SEQ ID NO:407)

Exon 9 Fragment 2

Forward Primer

40 53378-53399 TGAGAGCAGCAGGGATGACTTC (SEQ ID NO:408)

Reverse Primer

Complement of 53948-53926 AAACTCCTGATAGCCACTGGTGG (SEQ ID NO:409)

Exon 9 Fragment 3

45 Forward Primer

53869-53868 AGATCCTCCGCCGAAATGTC (SEQ ID NO:410)

Reverse Primer

Complement of 54583-54560 TTACTCTTCTGAGATGCCCGAG (SEQ ID NO:411)

50 Exon 9 Fragment 4

Forward Primer

54334-54355 TGGGCAGTGGCATTGTCTACTC (SEQ ID NO:412)

Reverse Primer

Complement of 54769-54750 TTCCAGGAGGTGGCATTTCC (SEQ ID NO:413)

55

Sequencing reactions were performed using the Big-Dye terminator kit from PE Biosystems (Foster City, CA) according to the manufacturer's instructions. The sequencing products were analyzed on an ABI 477 automated sequencer (PE Biosystems, Foster City, CA).

# 5 Analysis of Sequences for Polymorphic Sites

Sequences were analyzed for the presence of polymorphisms using the Polyphred program (Nickerson et al., 14 *Nucleic Acids Res.* 2745-2751, 1997). The presence of a polymorphism was confirmed on both the strands. The polymorphisms and their locations in the ILAR $\alpha$  gene are listed in Table 3 below.

Table 3. Polymorphic Sites Identified in the IL4Rα Gene Polymorphic Nucleotide Position in Nucleotide Position in Reference Variant	_
Site Number   GenBank Accession   Figure 1   Allele   Allele	Example
PS1 97137(Acc#AC004525) 32884 A G	1B
PS2 97118(Acc#AC004525) 32903 C T	1A
PS3 97060(Acc#AC004525) 32961 G T	1A
PS4 96886(Acc#AC004525) 33135 G C	1A
PS6         94272(Acc#AC004525)         35749         A         G           PS6         94258(Acc#AC004525)         35763         C         T	1A,1B 1A,1B
	1A, 1B
	1A 1A
	1B
1010 01011(100011000)	1A
PS11 94063(Acc#AC004525) 35958 G A	1B 1A
PS12 92691(Acc#AC004525) 37330 G A PS13 92548(Acc#AC004525) 37473 C T	
	1A
	1A
PS15 92430(Acc#AC004525) 37591 G A	1A
PS16 92417(Acc#AC004525) 37604 A T PS17 92377(Acc#AC004525) 37644 C A	1A 1A
	1A
PS19 86575(Acc#AC004525) 43446 G A PS20 86318(Acc#AC004525) 43703 T C	1A
	1A
PS21 77013(Acc#AC004525) 53008 A C	1A
PS22 76922(Acc#AC004525) 53099 C T PS23 76868(Acc#AC004525) 53153 T C	1B
	1A
PS24 <sup>R</sup> 76608(Acc#AC004525) 53413 A C	1A
PS25 76565(Acc#AC004525) 53456 G T	1A, 1B
PS26 <sup>R</sup> 76516(Acc#AC004525) 53505 T C	1A, 1B
PS27 76514(Acc#AC004525) 53507 C T	1A
PS28 76508(Acc#AC004525) 53513 T C	1A
PS29 <sup>R</sup> 76300(Acc#AC004525) 53721 T C	1A, 1B
PS30 76106(Acc#AC004525) 53915 C T	1A
PS31 <sup>R</sup> 76080(Acc#AC004525) 53941 A G	1A, 1B
PS32 76072(Acc#AC004525) 53949 G A	1A
PS33 75784(Acc#AC004525) 54237 C T	1A
PS34 75553(Acc#AC004525) 54468 T G	1A, 1B
PS35 75410(Acc#AC004525) 54611 T C	1A
PS36 · 75323(Acc#AC004525) 54698 T C	1A, 1B
PS37 75321(Acc#AC004525) 54700 T C	1A
PS38 75280(Acc#AC004525) 54741 C T	1A
PS39 75241(Acc#AC004525) 54780 C G	1A
PS40 74938(Acc#AC004525) 55083 A G	1A
PS41 74879(Acc#AC004525) 55142 G A	1A
PS42 <sup>R</sup> 74693(Acc#AC004525) 55328 G A	1A
PS43 <sup>R</sup> 74591(Acc#AC004525) 55430 C T	1A
PS44 74482(Acc#AC004525) 55539 C T	1A
PS45 74263(Acc#AC004525) 55758 G A	1A

RPreviously reported in the literature

# Example 2

This example illustrates analysis of the ILAR $\alpha$  polymorphisms identified in the Index Repositories for human genotypes and haplotypes for all polymorphic sites except PS1, PS9, PS11, PS21, PS22, and PS23.

A sampling of different genotypes containing these polymorphisms that were observed in these reference populations are shown in Table 4 below, with the haplotype pair indicating the combination of haplotypes determined for the individual using the haplotype derivation protocol described below. In Table 4, homozygous positions are indicated by one nucleotide and heterozygous positions are indicated by two nucleotides. Missing nucleotides in any given genotype in Table 4 can typically be inferred based on linkage disequilibrium and/or Mendelian inheritance.

**5**·

10

. [	0 0	<u></u>	ا س	-	2	9	9	6	8	~	4:	4	5	ဏ	2	_	0	7	4	æ	-	7	0		÷.	-	3		<b>T</b>	4	7	~	10		6	4	0	4	ဖြ	<b>ा</b>	ភា
	HAP	2	2	7 7	2 1	2 1	2 3	2 3	2 4	3	3 2	3 4	4	4 4	- 12	. 12	1 6	2 1	13 1	13 3	15 5	7 2	1	65	28 3	9 3	2 3	41 8	1 4	41 4	43 2	44	4	4 8	4 30	44 3	44 4	4	44	46 5	
	S 5	0	₽VG	0	: :	Ð	ີ .	S	<u>5</u>	:  S/	A	٧	5	A	9	9	٧	G 1		ØG1	Н	G 11	-	G 12	A  2	A 2	/G 3	A 4	A 4	A 4	G	A 4	<u>6</u>	A 4	A 4	A/G 4	A 4	A	4	7	읡
ŀ	S 4	S	S	Ü	Ü	၁	Ü	S	CA	∀ 0	၁	၁	C	၁	၁	၂၁	၁	၁			Н	-	၁	၁	၁	S	C	C	ပ	ပ	ر ن	ပ	CA	S	Ĵ	CA	O	3			3
ŀ	PS 7	O	် ပ	ာ	_ ပ	S	Н	Н	C	၂၁	Н	၁	၁	၁	_	ာ	Н	၁			၁	Н	ပ	⊢	Ψ;	_	,T	Τ,	၁	၁	-	ပ		Ψź	H	Н	ပ ပ	H	0		1
ı	45 F	ပ	9	9	$\dashv$	Н	Н		AVG	₩G	A.	٧	₽VG	Н	ຍ	9	Н	9		7	၁	щ	Н	19	A	٧	NGC	A C	A	Н	ပ	-	V	AC	A	9	Ā	V	`	<u>۷</u>	
ı	PS + 4	Ŋ	GA	9	-	$\vdash$	9		g G	∀ ୭	Н	၅		Н	9	_	Н	-	_	_	ຍ	_	Н	9	Н	Ð		Н	-	_		၅	-	Н	Ð	G A		Н	-+	o k	5
ı	& 6 2	A	G/A	٧	٧	A	١	A	G/A	G/A	<sub>O</sub>	၁	7	Н	٧	١v	G	A		₹ Ø	٧	-	_	G/A	9	ß	ပ		ى ق	ပ	₹	ပ	ပ	Н		$\Rightarrow$	-	b	하	5	5
ł	<u>ය</u> ස	ပ	0	၁	ပ	၁	၁	၁		-	Н		1	-	၂၁	10	C	_			၁	-		<u>၂</u>	-	Н	၁	Н	-	-		$\dashv$	-	Н		$\overline{}$	ပ	Н		5	5
	လ ဆ	ပ	၁	၁	o	ပ	၁	Ċ	၁	၂၁	၁	၁	၁	၁	၂၁	၁	5	-	$\vdash$	ပ	ပ	ပ	CT	၁	_	၁	၁	၁	ပ	ပ	ᇬ	ပ	ပ	၁	၁	$\dashv$	၁	J	ᆄ	ok	5
	37 37	F	Ţ	T	F	T	1	-	T	T	F	H	1	F	Ţ	T	T (	Ţ	2/2	S	T	F	1 K	Ţ	1/0	2	ī	2	ပ	ઇ	디	L	ᅵ	-		L	L	F	ᆎ	_	_
•	လ ဗွ	1	Ļ	2	F		2	<b> </b> -	L	T	F	1	2	T	T	Ţ	T	Ļ	2	I	1	ı	1	Į)	Ţ	Ţ	F	707	O	5		L	H	F			<u>-</u>		-	ള	2
	လ ဗ	⊢	T	Ţ	F	-	Ţ	۲	T	Ţ	Ŧ	T	Ţ	F	1	T	T	ı	્ર	2	_	Ţ	T	Ţ	L	L	F	ı	F	H		_	F	I	F	니	⊢	,_	;;	=	
	8 %	F	T	T	F	<b>-</b>	⊢	H	Ţ	T	ī	┸	Ţ	1/6	Ţ	T	T	⊢	ত্র	-	ı	Ţ	1/G	F	1/0	9	<u>1</u>	၅	Ö	<u>1</u>				<u>1/G</u>	H	F	_			2	2
	3 %	ပ	၁	ပ	ပ	ပ	၁	ပ	ပ	၂၁	၁	၁	Ċ	၁	၁	၂၁	ပ	ပ	ပ	ပ	ပ	ပ	၂	၁	ပ	ပ	ပ	ပ	•	ပ	ပ	ပ	ပ	၁	၁	ပ	ပ	ပ	ပြ	ပြု	3
	8 S	ပ	9	၅	ပ	ဝ	၅	၁	၅	၉	ပ	G	၅	၅	9	G	ß	ပ	ပ	ပ	ß	ပ	ຍ	ပ	b	G/A	G/A	G/A	ပ	ပ	ပ	ပ	ပ	G/A	υ	ပ	ပ	စ	တ	υļ	5
1	3 FS	4	٨	₽VG	٧	A	ΑG	٧	٧	٧	A	٧	ອ	A/G	₽VG	G	၅	ပ	ပ	ပ	A	AG AG	ຽ	ΑG	ပ	_	_	ပ	ပ	₽ Q	ş	٧	ខ្ទ	VG(	ş	₹	A	∢	₹!	इर	5
ļ	<b>₹</b> 8	ပ	ပ	၁	ပ	ပ	Ö	၁	၁	၁	၁	၁	၁		S	၁	ပ	Ç	ပ	ပ					ပ	၁	ပ	ပ	ပ	Ö	O	ပ	ပ	် ပ	Ö	5	ပ	ပ	ပ	٤	7
.	2 PS		۲	T	F	T	Τ	F	T	Ţ	Ŧ	Ţ	T/C	F	T/C	၁	ပ	ı-	1/0	S F	_	70	T/C	Ţ	ပ	<u>ارد</u>	2	<b>-</b> -	ı-	F	일	F	Ş	⊢	H	Ē	H	一	-	_ [	₹
· ]	PS 28	ı	F	T	F	۲	Ī	1	T	Ţ	T	Ţ	1/0	1/0	1/0	၁	ပ	⊢	ပ	2	H	15	၁	Τ	T/C	ပ	ပ	2	F	<b>-</b>	्	F	ខ្ព	1/0	⊢	F	F	F	-1	Σ,	5
	27 ZS	ပ	ပ	၁	ပ	ပ	ပ	၁	၁	၁	ပ	ပ	ပ	Ü	၁	၁	၁	၁	ပ	ပ	Ċ	ပ	၁	ပ	C	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	U	ပ	O	عاد	3
	S 8	F	F	Τ	⊢	⊢	Ţ.	T	Τ	L	T	T	T/C	⊥	1/0	၁	T/C	۲	<b>j</b> –	F	۲	T/C	1	Τ	Τ	F	T/C	H	F	H	S L	H	ខ្ព	⊢	F	F	F	一		- -	7
	PS 25	တ	ပ	ß	ပ	Ö	G	ຶ	ပ	9	၅	ပ	GЛ	GП	GЛ	T	ı	ပ	F	5	ပ	GЛ	1	ຶ	GЛ	⊢	Τ	GΤ	တ	ß	F	တ	ΕŢ	Gπ	ပ	ပ	ပ	ပ	ပ	3	-1
Gen	PS 24	٧	٧	٧	A	A	٧	٧	٧	٧٠	٧	٧	AC	AC	AC	Э	C	Α	ပ	AC C	A	AC	၁	٧	A/C	ပ	C	AC	۷	۲	Ş	∢	Ş	AC	٧	⋖	⋖	⋖	₹.	्र	3
	. S 8	ı	Ţ	СT	CΤ	CT	СT	ᅜ	СT	Ţ	S	СT	Ċ	$c\pi$	ĽЭ	၁	၁	ပ	ပ	ပ	СT	СT	ပ	ပ	င	ပ	C	ပ	ပ	ပ	5	5	ပ	ပ	CT	СŢ	ပ	ပ	5	3	3
#	PS 19	ပ	O	G	ß	G	ອ	9	9	9	9.	9	0	9	9	9	9	၁	၅	G	G	G	G	G	G	ပ	G	G	ပ	G	စ	ß	g	6	Ö	Ö	G	ပ	တ	9 (	2
for the IL4R $lpha$	S &	ပ	ပ	5	ပ	5	C	S	CT	ပ	ပ	ĽЭ	ı	၁	5	_	ပ	-	5	5	ပ		၁	T	ပ	ঠ	5	5	ပ	5	ပ	5		⊢	5	٠.	Ŀ	느	5	2	ᆀ
<b>₽</b>	4 73	ပ	ပ	၁	၁	ပ	၁	၁	<u>၁</u> ၂	0	၁	ပ	၁	၁	၁	2	၁	၁	ပ	ပ	၁	CA	၁	၁	ပ	0	ပ	၁	의	ပ	ઇ	ပ	ပ	၁	ပ	Ŀ		ပ	ပ	ပ	2
å	S &	٧	۷	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	٧	⋖			٨	٧	٧	٧	۷	٠A	٧	۷	٧	≤	۲	۲	٧	٧	۲	٧	≤	⋖ •	⁴	
`≥	S 75	┞	Ö	9	g	9	၅	9	၅	ပ	၅	9	၁	ß	ຶ	ပ	9	ပ	ပ	Ŋ	G/A	G/A	Ð	၅	ပ	ပ	9	ပ	ပ	Ø	ပ	ပ	Ö	ß	ပ	ပ	ပ	တ	တ	2) (	의
Observ	S 4	┿	ပ	C		Ö	ш	·	0	L	H		$\vdash$		_	-	Н	ш	ပ	_			_	ш	$\vdash$	$\dashv$	-	၁	_	-	의		_	_	ပ	_	ပ	Н	ပ	<u>5</u> '	2
	S 5	O	ပ	၁	2	၁	၁	၁	၁	ပ	S	၁	၁	5	ပ	ပ	၁	ပ	ပ	ပ	5	ပ	S	ပ	ပ	ပ	ပ	ပ	ပ	ပ	의	ပ	ပ	၁	ပ	ပ	Ċ.	의	ပ	علا	깈
Pai	\$ 5			<u> </u>	G/A	၅၂	9	၅	၅၂	ပ	ອ	9	၅	ຶ	<u>ຍ</u>	Ø	9	ຶ																				9	ပ	_	-
₽	₹. 6		1	ഗ				2	5	ပ		5	L	T	5	_	-	_	느	5	Ċ.	ပ				ပ	S	5	ပ	5	의	5	۲		Ş	5	5	늬	ᆣ		<u></u>
핗	გ ∞	_	۲	1			⊥	⊥	۲	Ľ	⊢	T	┸	۲	1	_	⊢			٢	۲	٢		٢		늬	٢	⊥	Τ	۲		۲		⊥	۲	۲	⊢	늬	1		3
ĕ	8 ~																		ව																			၅	-4	y k	
ğ		ပ	1			S	၁	၁	O	ပ	O	ပ	ပ	ပ	S	ပ	ပ	၁	3	ပ	ပ	ပ	၁	ပ	O	ပ	၁	ပ	ပ	ပ	의				ပ	ပ		9		ပင်	SAC.
ş		⋖	۷	R/S	<i>d</i> /9	G/A	٧	٧	Ø/S	۷	Ø	<u>0//</u>	စ	ပ	<b>∜</b> 5	ග	₹/5	9	ပ	8/9	Ø/∀	٧	9	၁	<b>G/A</b>	Y	٧	ပ		ပ	≤	8	ပ	ပ	G/A	ď⁄	Ø,	၅			
Genotypes and HAP Pairs	<b>8</b> 4	ပြ	9	၅	၅	၅	တ	၅	Ø	Ö	ပ	ပ	Ű	ပ	Ŀ	တ	၅	၅	8	ပ	Ø	g	9	Ö	ပ	ଠ	ပ	ပ	ပ	တ	ဖ	ပ	ပ	ပ	ပ	ဗ	ပ		တြ	ဗြ	
		-	←			_	_																																တ		
Table 4.	& ~	ပ	_		)		ပ			ပ	ပ	ပ	5	ပ	Ŀ	ပ	ပ	ပ	၁	ပ	5	ပ	၁	ပ	ပ	ပ	ပ	5	۲	5	O	ပ	5	ပ	ပ	ပ	ပ	ပ	<u>ာ</u>	၁	3
۲		<u>                                     </u>	7	3	4	5	ဖ	7	ထ	0	5	÷	72	13	7	15	16	17	8	19	8	7	22	23	24	33	56	27	78	33	္က	3	32	33	8	25	~~ I	. [	<u>س</u> ادً	IA	

The haplotype pairs shown in Table 4 were estimated from the unphased genotypes using an extension of Clark's algorithm (Clark, A.G. (1990) *Mol Bio Evol* 7, 111-122), as described in U.S. Provisional Patent Application filed April 19, 2000 and entitled "A Method and System for Determining Haplotypes from a Collection of Polymorphisms". In this method, haplotypes are assigned directly from individuals who are homozygous at all sites or heterozygous at no more than one of the variable sites. This list of haplotypes is then used to deconvolute the unphased genotypes in the remaining (multiply heterozygous) individuals.

By following this protocol, it was determined that the Index Repositories examined herein and, by extension, the general population contains the 53 human IL4Rα haplotypes shown in Table 5 below.

10

Table 5A. Haplotypes Observed for the IL4R $\alpha$  Gene

				_			_	,	,		_			_	,	_			_			_			_		,		_		_		_	_
PS 45	⋖	ပ	A	တ	ပ	⋖	တ	⋖	⋖	٧	٧	ပ	۷	⋖	ß	ပ	ပ	⋖	⋖	₹	4	တ	0	V	۷	Þ	⋖	⋖	⋖	⋖	4	⋖	ပ	တ
S 4	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	3	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	-	ပ	ပ	ပ	ပ	၁	ပ	ပ
PS 43	ပ	ပ	ပ	ပ	ပ	ပ	ပ	-	ပ	C	၁	ပ	۳	۲	ပ	ပ	ပ	ပ	H	⊢	ပ	ပ	ပ	ပ	<b> -</b> -	-	-	۱	-	ပ	ပ	T	ပ	ပ
PS 42	₹	Ö	4	ပ	A	¥	ပ	₹	∢.	A	٧	၁	4	۷	ပ	ပ	ග	₹	⋖	⋖	⋖	ပ	ပ	⋖	⋖	⋖	⋖	⋖	⋖	⋖	⋖	A	ပ	စ
PS 41	ပြ	ပ	ပ	ပ	Ø	ග	Ø	ပ	ပ	၅	၁	ပ	<u>ග</u>	ပ	ပ	ပ	ပ	ග	တ	ပ	တ	တ	Ö	ပ	တ	တ	ပ	ပ	တ	၁	ပ	ပ	ပ	ပ
PS 4	ပ	٧	တ	A	ຍ	ပ	∢	ပ	ß	ß	ຶ	٧	ပ	S	∢	⋖	⋖	ပ	ပ	ပ	ပ	⋖	တ	ပ	ပ	တ	ပ	ပ	တ	Ö	ပ	ອ	G	₹
2 PS	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	င	၁	၁	ပ	O	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ
28 38	ပ	ပ	ပ	ပ	ပ	Э.	ပ	ပ	ပ	Ţ	၁	၁	ပ	ပ	ပ	ပ	ပ	ပ	၁	၁	Э	ပ	ပ	ပ	၁	ပ	ပ	ပ	၁	0	ပ	ပ	ပ	ပ
PS 37	F	۲	۲	Ţ	μ,	1	<b>)</b> —	۲	<b>)</b> —	⊢	ပ	1	ပ	1	⊢	1	-	T	T	1-	T	۲	F	1-	ပ	ı	Ţ	ပ	ပ	T	1	T	۲	F
98 36	F	H	F	ပ	۲	1-	F	F	F	T	ပ	1	-	ပ	F	-	-	F	၁	1	7	Ţ	F	T	T	ပ	T	T	Ţ	ī	1	F	F	F
PS 35	F	H	1	Ţ	Ţ	T	T	-	Τ	T	T	Ţ	ပ	1	<b> -</b> -	۳	-	Ţ	Ţ	Ţ	1	7	Ţ	۲	ပ	Ţ	T	Ţ	1	T	Ţ	F	L)	F
PS 34	۲	Ļ	Ţ	1	Ţ	Ţ	Ţ	တ	T	Ţ	ß	1	1	G	1	T	-	Ţ	9	Ţ	ß	1	F	Ţ	┸	ß	T	ල	ጉ	F	T	ပ	H	-
PS 33	ပ	ပ	၁	3	Э	Э	ပ	၁	ပ	ပ	၁	3	3	0	Э	၁	ပ	3	၁	3	3	Э	၁	၁	ပ	Э	C	0	၁	ပ	၁	ပ	ပ	ပ
PS 32	ပ	ව	9	9	9	ອ	ອ	٧	ပ	G	9	9	ອ	9.	9	ອ	တ	ອ	ອ	9	9	ອ	ອ	9	ອ	ອ	9	9	9	ပ	၅	٧	ပ	ပ
PS 31	Þ	⋖	٧	9	9	٧	9	.9	ပ	ပ	တ	9	ອ	ອ	٧	٧	၅	٧	ອ	9	ອ	٧	٧	٨	ව	9	٧	9	ອ	9	ß	ပ	ຍ	A
PS 30	ပ	ပ	၁	၁	Э	၁	ပ	0	ပ	၁	C	၁	Э	၁	2	Ċ	ပ	၁	၁	၁	ပ	၁	၁	၁	၁	၁	၁	၁	၁	Э	Ö	ပ	ပ	
PS 29	느	느	Ţ	١	၁	L	၁	Ł	၁	ပ	⊢	۲	၁	1	1	L	ပ	۲	ပ	၁	_	Ţ	⊥	⊥	ပ	1	-	၁	ပ	L	၁	-	ပ	니
PS 28	L	۲	۲	Ţ	၁	L	ပ	၁	ပ	ပ	⊢	┸	၁	ပ	Ţ	⊥	ပ	T	ပ	၁	ပ	_	ΙI	L	၁	၁	L	۲	ပ	Ţ	၁	ပ	ပ	-
PS 27	ပ	ပ	ပ	Ċ	ပ	၁	ပ	၁	၁	ပ	၁	ပ	ပ	ပ	၁	၁	၁	၁	┸	۲	ပ	၁	၁	၁	ပ	၁	၁	ပ	ပ	ပ	၁	ပ	ပ	ပ
PS 26	上	上	۲	<b>!-</b>	ပ	۲	ပ	1	ပ		٢	<del> </del>	ı	1	Ţ	۰	ပ		ĹΙ	1	۲		ı	느	1	_	1	╚	H	1			ပ	늬
PS 25	ပ	ပ	ပ	၅	L	9	L	Ŧ	L	۲	0	ပ	۲	1	9	9	1	၅	T	T	۲	၅	9	9	⊥	1	၅	O	۲	၅				ပ
PS 24	⋖	۷	٧	٧	ပ	٧	၁	ပ	Н	_	٨	٧	၁	၁	٧	٧	၁	٧	၁	၁	Ċ	٧	A	٧	ပ	ပ	٧	٧	၁	٧	ပ	ပ	ပ	≤
PS 20	ပ	上	⊥	၁	၁	၁	ပ	ပ	၁	ပ	ပ	ပ	၁	၁	၁	ပ	ပ	၁	Ö	၁	ပ		ပ	ပ	ပ	ပ	၁	ပ	ပ	⊢	ပ	ပ	ပ	늬
PS 19	ပ	ပ	9	ပ	9	Ÿ		9	G	ပ	O	g	9	ß	9	ဝ	9	ß	၅	၅	ပ	ပ	G	Ö	ຶ	ຍ	G	ß	၅	ව	ပ	ပ	ပ	ပ
- 2 = 2 = 2 = 2 = 2 = 2 = 2 = 2 = 2 = 2	上	ပ	၁	1-	Ţ	<u> </u>	Ţ	_	-	_	ပ	ပ	-	ပ	ပ	_	ပ	၁	ပ	၁	ပ	ပ	느	ပ	ပ	_	_	ပ	ပ	၁	ပ	ပ	느	의
PS 17	ပ	ပ	၁	ပ	၁	ပ	၁	၁	ပ	ပ	ပ	ပ	၁	ပ	၁	ပ	<b>3</b> .	٧	ပ	ပ	ပ	٧	၁	၁	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	의
PS 16	⋖	A	٧	٧	A	A	٧	٧	Ą	۲	٧	٧	٧	٧	٧	٧	A	A	А	٧	A	A	٧	٧	٧	Æ	٧	4	٧	٧	٧	۷	_	≤
PS 25	စ	ပ	9	ပ	၅	၅		9	G		၅	ပ	9	၁	9	၅	٧	9	ပ		g	၁	တ	၅	g	ව	ß	හ	၅	၅	ပ	ပ	ပ	의
PS 14	ပ	ပ	Н	ပ	၁	၁	Н	၁	Н	-	-	၁	_		၁	၁	၁	၁	၁	ပ	ပ	င	၁	၁	ပ	၁	ပ	၁	၁	ပ	ပ	의	ပ	의
5 PS	2	၁	0	၁	ပ	၁	၁	၁	-	-	_	-	_	ပ	1	ပ	၁	၁	T	ပ	1	ပ	၁	Ţ	٢	ပ	၁	ပ	ပ	ပ	ပ	-		의
2 LS	ပ	ပ	Ь	၁	_	၅	Н	Э	Н		_	-	_	-	_	$\perp$	Н	-	ပ		_	_	_	_	-		Н	-	-		-		-	의
PS 10	┰	ပ	၁			I		1	2	ပ	ပ	၁		<b> -</b> -	2	⊥	၁	ပ	S.	ပ	0	၁	1	၁	ပ	۲	۲	ပ	ပ	၁	ပ	의	의	의
1	-	Н	_	_	Ţ	_	ш	I	-	_	-	_	_	-		ш		-	ш	⊢		_	1	_	_	۲	_			۲	_	늬	듸	
PS 7		-	_	_	_			-	_	_	_	_	_	_		_	-	_	1		_	_		-					_	_	_	_		_
S 9		-	Н	-	-		_	_	-	_	_	-	_	_			-	-	_		_	_	_	-	-		-		-	-		-	ပ	ᅱ
PS 5	-	-	٧	_	-	_	$\blacksquare$	_	٧	_	_	-	_		$\vdash$	-	-	_	$\overline{}$	٧	_	_			-		_		_	٧	ш	-+	≤	1
PS 4	_	ပ	Н	-	_	$\overline{}$	-	_	ш	_	-			_	_		Ī	-				-	_	_	_	_	-	-	-			-	-	의
PS 3		ပ			_	_	ш	_	၅		_			_	_	_	_	-		_		-	_	-	_	_	ပ	-	_	_	-		의	-4
PS 2	ပ	ပ	ပ	ပ	T	ပ	ပ	ပ	ပ	ပ	$\overline{}$	-	-	-	-	$\vdash$	Н	ပ	ပ	ပ	9	ပ	ပ	ပ	٢	ပ	의	ပ	ပ	ပ	ပ	의	이	의
HAP ID	-	2	က	4	2	9	-	80	6	위	=	2	13	14	12	16	47	8	13	8	21	ឧ	23	24	22	82	23	8	53	္က	3	32	8	ᇷ
				_	-		_		_		_	_	_	_			_		_	_	_			_	_	_	_		_	_	_		_	_

Table 5B. Haplotypes Observed for the IL4R $\alpha$  Gene

				_	_						_		_							
PS	45	⋖	တ	٧	ပ	ပ	∢	⋖	4	ပ	⋖	တ	∢	ပ	⋖	⋖	⋖	ပ	တ	٧
Sd	4	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	Ŋ	ပ	ပ	ပ	ပ
PS	43	ပ	ပ	⊢	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	۲	ပ	ပ	⊢
Sd	42	⋖	ပ	⋖	တ	ပ	٧	⋖	⋖	ပ	⋖	တ	⋖	4	⋖	⋖	4	ပ	⋖	⋖
Sd	4	ပ	Ø	Ø	တ	ပ	ပ	တ	Ø	ပ	ပ	O	ပ	၅	ပ	Ø	Ø	Ø	Ø	Ø
PS	\$	စ	⋖	တ	۷	۷	ပ	ပ	9	<	ပ	⋖	ပ	ပ	ပ	Ø	တ	⋖	Ø	ပ
PS	39	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	ပ	ပ	ပ
bs	88	ပ	ပ	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	ပ	ပ	ပ	၁	ပ	ပ	ပ	ပ
bS	37	上	۲	Ŀ	۲	<u> -</u>	느	ပ	上	느	<u> -</u>	<u> -</u>	۲	┸	۲	۳	E	٢	٢	느
BS	36	上	ပ	上	۲		⊥	ပ	_	上	_	ᆫ	上	۲	٢	T	ပ	۲	۲	ပ
PS	35	_	۲	L	۲	-	L	느	٢	上	۳	느		۲	T	1	۲	<u>-</u>	۲	۲
PS	34	T	L	Ø	٢	Ŧ	ı	၅	L	L	Ŀ	L		ı	⊥	១	တ	۰	۲	ပ
PS	33	上	၁	ပ	၁	၁	ပ	၁	ပ	ပ	ပ	0	၁	၁	၁	၁	ပ	ပ	ပ	ပ
PS	32	9	စ	٧	ပ	၅	၅	9	9	စ	ອ	9	9	၁	ပ	ව	ပ	၅	ပ	Ø
PS	31	٧	9	၅	9	٧	٧	ပ	٧	၅	٧	٨	٧	ව	٧	ပ	9	٧	9	၅
PS	30	၁	ပ	ပ	ပ	၁	C	၁	၁	0	၁	0	2	C	၁	ပ	၁	ပ	ပ	ပ
PS	29	ᆫ	T	-	۰	Ţ		느	٢	2	Ţ	Ţ	۲	၁	T	┸	⊢	┸	၁	ī
PS	28	L	_	ပ	۲	II	L	Т	٢	ပ	Ţ	_		၁	T	၁	၁	Ţ	ပ	ပ
PS	27	၁	၁	ပ	၁	၁	၁	ပ	၁	၁	၁	၁	၁	ပ	2	ပ	၁	0	၁	ပ
PS	2	T	Ţ	۲	٢			-	_	၁	┸	Ţ	_	ပ	1	1	_	1	၁	
PS	25	9	<u> </u>	٢	9.	9	G	၅	9	T	၁	9	၅	_	9	T	⊥	9	1	┙
PS	24	Y	٧	၁	A	٧	٧	H	μ.	3	A	_	٧	ပ	A	S	၁	٧	<u>၁</u>	၁
PS	20	⊥	<u> </u>	၁	<u>၁</u>	၁	၁	၁	T	၁	၁	T	۲	ပ	၁	ပ	ပ	_	<u> </u>	၁
PS	19	9	9	Ö	၅	၅	$\vdash$	၅	9	Н	9	-	9	g	9	9	9	9	9	၅
PS	18	၁	Ţ	_	2	T	၁	ပ	၁	၁	T	2	၁	_	1	၁	၁	၁	T	၁
PS	17	၁	၁	၁	၁	၁	၁	၁	0	၁	၁	၁	၁	ပ	0	ပ	ပ	၁	၁	၁
PS	16	A	٧	٧	٧	Y	٧	A	٧	٧	٧	A	A	A	٧	٧	٧	٧	٧	١×
PS	15	9	-	၅	ဗ	9	9	9	၅	9	၅	9	၅	၁	9	9	9	H	9	၅
3 PS	14	၁	⊢	၁	ပ	၁	Н	Н	Н	2	ш	Н	၁	ပ	၁၂	Н	Н		၁၂	Н
PS	13	၁	Н	၁	Н	O	၁		_	၁	Γ.	၁	$\vdash$	ပ	၁	⊥	)	L	<u>၁</u>	၁
s PS	12	<u> </u>	Н	9	Ð	၅	-	9	9		9	S	ဗ	ပ	၅	9	Н	Н	Ŀ	ອ
S PS	9	၁၂	<u> </u>	<u>၁</u>	C	ပ	Н	C	၁ 	2	⊢	۲		⊥	1	Ţ	T	၁	1	၁
S PS	8	1	_		۲	Ţ	1		1		_	۲		_	_	۲	1	⊢	۲	C
S PS	7	9 :	Н	9	ອ	၁	9	9	Y	-	-	-	9	ပ	G	9	9		9	၅
3 PS	9	<b>3</b>	Н	3	O	<u>ن</u>	ပ	<u>၁</u>	၁	၁	O	_	ပ	ပ	<b>O</b>	၁	ပ	၁	2	
S PS	5	Y !	Y	i A	i A	) Y	$\vdash$	9	١Į٨	١	ш	-	9	<b>O</b>	ව	ອ	9	۷	<u>ອ</u>	9
s PS	4	9 !	Н	Ð	9	9	S	ຍ	9	9	ົວ	_	Н	ပ	၁	9	<u>ე</u>	Н	9	ອ
S PS		9	g	ຶ	ပ	Ŋ	Н	$\vdash$	ຶ	ຶ	ຶ	-	-	ß	ပ	ß	၅	၅		၅
PS	7	၁	ပ	ပ	၁	ပ	_	-	ပ	ပ	၁	ပ	၁	ပ	_	၁	၁	၁	۲	T
HAP	₽	32	36	37	38	39	40	4	42	43	4	45	46	47	48	49	20	51	25	53

In view of the above, it will be seen that the several advantages of the inv ntion are achieved and other advantageous results attained.

As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description and shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

5

10

All references cited in this specification, including patents and patent applications, are hereby incorporated in their entirety by reference. The discussion of references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinency of the cited references.

## What is Claimed is:

5

10

. 5

10

5

1. An isolated polynucleotide comprising a nucleotide sequence s lected from the group consisting of:

- (a) a first nucleotide sequence which is a polymorphic variant of a reference sequence for Interleukin 4 Receptor Alpha(ILARα) gene or a fragment thereof, wherein the reference sequence comprises SEQ ID NO:1, and the polymorphic variant comprises at least one polymorphism selected from the group consisting of guanine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, cytosine at PS8, thymine at PS9, thymine at PS10, adenine at PS11, adenine at PS12, thymine at PS13, thymine at PS14, adenine at PS15, thymine at PS16, adenine at PS17, thymine at PS18, adenine at PS19, cytosine at PS20, cytosine at PS21, thymine at PS22, cytosine at PS23, thymine at PS33, guanine at PS34, cytosine at PS35, cytosine at PS36, cytosine at PS37, thymine at PS38, guanine at PS39, guanine at PS40, adenine at PS41, thymine at PS44, and adenine at PS45; and
- (b) a second nucleotide sequence which is complementary to the first nucleotide sequence.
- 2. The isolated polynucleotide of claim 1 which comprises an IL4Rα isogene.
- The isolated polynucleotide of claim 1 which is a DNA molecule and comprises both the first and second nucleotide sequences and further comprises expression regulatory elements operably linked to the first nucleotide sequence.
- 4. A recombinant organism transformed or transfected with the isolated polynucleotide of claim 1, wherein the organism expresses an IL4Rα protein encoded by the first nucleotide sequence.
- 5. The recombinant organism of claim 4 which is a nonhuman transgenic animal.
- 6. The isolated polynucleotide of claim 1, wherein the first nucleotide sequence is a polymorphic variant of a fragment of the ILARα gene, the fragment comprising one or more polymorphisms selected from the group consisting of guanine at PS1, thymine at PS2, thymine at PS3, cytosine at PS4, thymine at PS6, adenine at PS7, cytosine at PS8, thymine at PS9, thymine at PS10, adenine at PS11, adenine at PS12, thymine at PS13, thymine at PS14, adenine at PS15, thymine at PS16, adenine at PS17, thymine at PS18, adenine at PS19, cytosine at PS20, cytosine at PS21, thymine at PS22, cytosine at PS23, thymine at PS25, thymine at PS27, cytosine at PS28, thymine at PS30, adenine at PS32, thymine at PS33, guanine at PS34, cytosine at PS35, cytosine at PS36, cytosine at PS37, thymine at PS38, guanine at PS39, guanine at PS40, adenine at PS41, thymine at PS44, and adenine at PS45.
- 7. An isolated polynucleotide comprising a nucleotide sequence which is a polymorphic variant of a reference sequence for the ILARα cDNA or a fragment thereof, wherein the reference sequence comprises SEQ ID NO:2 and the polymorphic variant comprises at least one polymorphism selected from the group consisting of thymine at a position corresponding to nucleotide 237, adenine at a position corresponding to nucleotide 244, cytosine at a position corresponding to nucleotide 291, thymine at a position corresponding to nucleotide 501, adenine at a position corresponding to

nucleotide 554, cytosine at a position corresponding to nucleotide 939, thymine at a position corresponding to nucleotide 1242, thymine at a position corresponding to nucleotide 1293, cytosine at a position corresponding to nucleotide 1299, thymine at a position corresponding to nucleotide 1701, adenine at a position corresponding to nucleotide 1735, thymine at a position corresponding to nucleotide 2023, guanine at a position corresponding to nucleotide 2254 and cytosine at a position corresponding to nucleotide 2397.

- 8. A recombinant organism transformed or transfected with the isolated polynucleotide of claim 7, wherein the organism expresses a Interleukin 4 Receptor Alpha(ILARα) protein encoded by the polymorphic variant sequence.
- 9. The recombinant organism of claim 8 which is a nonhuman transgenic animal.

10

5 ·

- 10. An isolated polypeptide comprising an amino acid sequence which is a polymorphic variant of a reference sequence for the ILARα protein or a fragment thereof, wherein the reference sequence comprises SEQ ID NO: 3 and the polymorphic variant comprises one or more variant amino acids selected from the group consisting of threonine at a position corresponding to amino acid 82, histidine at a position corresponding to amino acid 185, isoleucine at a position corresponding to amino acid 579, serine at a position corresponding to amino acid 675, and alanine at a position corresponding to amino acid 752.
- 11. An isolated antibody specific for and immunoreactive with the isolated polypeptide of claim 10.
- 12. A method for screening for drugs targeting the isolated polypeptide of claim 10 which comprises contacting the ILARa polymorphic variant with a candidate agent and assaying for binding activity.
- 13. A composition comprising at least one genotyping oligonucleotide for detecting a polymorphism in the Interleukin 4 Receptor Alpha(IL4Rα) gene at a polymorphic site selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45.
- 14. The composition of claim 13, wherein the genotyping oligonucleotide is an allele-specific oligonucleotide that specifically hybridizes to an allele of the IL4Rα gene at a region containing the polymorphic site.
- 15. The composition of claim 14, wherein the allele-specific oligonucleotide comprises a nucleotide sequence selected from the group consisting of of SEQ ID NOS:4-79, the complements of SEQ ID NOS: 4-79, and SEQ ID NOS:80-231.
- 16. The composition of claim 13, wherein the genotyping oligonucleotide is a primer-extension oligonucleotide.
- 17. A method for genotyping the Interleukin 4 Receptor Alpha(ILARα) gene of an individual, comprising determining for the two copies of the ILARα gene present in the individual the identity of

the nucleotide pair at one or more polymorphic sites (PS) selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45.

18. The method of claim 17, wherein the determining step comprises:

5

10

5

10

- (a) isolating from the individual a nucleic acid mixture comprising both copies of the ILARa gene, or a fragment thereof, that are present in the individual;
- (b) amplifying from the nucleic acid mixture a target region containing at least one of the polymorphic sites;
- (c) hybridizing a primer extension oligonucleotide to one allele of the amplified target region;
- (d) performing a nucleic acid template-dependent, primer extension reaction on the hybridized genotyping oligonucleotide in the presence of at least two different terminators of the reaction, wherein said terminators are complementary to the alternative nucleotides present at the polymorphic site; and
- (e) detecting the presence and identity of the terminator in the extended genotyping oligonucleotide.
- 19. A method for haplotyping the Interleukin 4 Receptor Alpha(II.4Rα) gene of an individual which comprises determining, for one copy of the II.4Rα gene present in the individual, the identity of the nucleotide at one or more polymorphic sites (PS) selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45.
- 20. The method of claim 19, wherein the determining step comprises
  - (a) isolating from the individual a nucleic acid molecule containing only one of the two copies of the ILARα gene, or a fragment thereof, that is present in the individual;
  - (b) amplifying from the nucleic acid molecule a target region containing at least one of the polymorphic sites;
  - (c) hybridizing a primer extension oligonucleotide to one allele of the amplified target region;
  - (d) performing a nucleic acid template-dependent, primer extension reaction on the hybridized genotyping oligonucleotide in the presence of at least two different terminators of the reaction, wherein said terminators are complementary to the alternative nucleotides present at the polymorphic site; and
  - (e) detecting the presence and identity of the terminator in the extended genotyping oligonucleotide.
- 21. A method for predicting a haplotype pair for the Interleukin 4 Receptor Alpha(IL4Rα) gene of an individual comprising:

 (a) identifying an IL4Rα genotype for the individual at two or more of polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45;

- (b) enumerating all possible haplotype pairs which are consistent with the genotype;
- (c) accessing data containing the IL4Rα haplotype pairs determined in a reference population; and
- (d) assigning a haplotype pair to the individual that is consistent with the data.
- 22. A method for identifying an association between a trait and at least one genotype or haplotype of the Interleukin 4 Receptor Alphagene which comprises comparing the frequency of the genotype or haplotype in a population exhibiting the trait with the frequency of the genotype or haplotype in a reference population, wherein the genotype or haplotype comprises a nucleotide pair or nucleotide located at one or more polymorphic sites selected from PS1, PS2, PS3, PS4, PS6, PS7, PS8, PS9, PS10, PS11, PS12, PS13, PS14, PS15, PS16, PS17, PS18, PS19, PS20, PS21, PS22, PS23, PS25, PS27, PS28, PS30, PS32, PS33, PS34, PS35, PS36, PS37, PS38, PS39, PS40, PS41, PS44, and PS45, wherein a higher frequency of the genotype or haplotype in the trait population than in the reference population indicates the trait is associated with the genotype or haplotype.
- 23 The method of claim 22, wherein the haplotype is selected from haplotype numbers 1-53 shown in Table 5.
- 24. The method of claim 23, wherein the trait is a clinical response to a drug targeting IL4Rα.
- 25. A computer system for storing and analyzing polymorphism data for the Interleukin 4 Receptor Alphagene, comprising:
  - (a) a central processing unit (CPU);
  - (b) a communication interface;
  - (c) a display device;

5

5

5

10

- (d) an input device; and
- (e) a database containing the polymorphism data; wherein the polymorphism data comprises the genotypes and haplotype pairs shown in Table 4 and the haplotypes shown in Table 5.
- 26. A genome anthology for the Interleukin 4 Receptor Alpha(ILARα) gene which comprises ILARα isogenes defined by haplotypes 1-53 shown in Table 5.

# POLYMORPHISMS IN THE IL4Ra GENE

TGTGAGCTAC	TGTGTCTGGC	CTGAATAATA	AAATTTAAAA	CAATTTTTCA	
				CTCAAACCCC ·	30100
		CCACCTTCCC			30200
		GAGCAATACG			30200
		TTATAGCAGC			30200
		TAAAGTGTTC			30300
		ATCCCCACAA			30300
		GATAAGGAAA			30400
				TTAAACCTCA	. 50400
		GTGTAGAGGG			30500
		AGTTGTTGTG			30300
		ATGATTATAG			30600
		ACACAAATGG			30000
		ATCCCAGCAC			30700
		CAAGACCAGC			30700
		AAAATTAGCC			20000
					30800
•		GGCTGAGGCA			20000
GAGGIGGAGG	TTGCAGTGAG	CTGAGATTGC	G	TCCAGCCTTG	30900
CACACACACA	CA CA CHCHCH		-	7007770777	
		CTCGGAAAAA GCAGGGGAGG			21000
					31000
		TCAGCACTAA		•	21100
		ATGGGGTGGC	TTTGCTCTGG	GCTCCTGTTC	31100
-	3: 31071	COMCON COMC	CON N COMOMO	C	•
CCIGIGAGCI	311	GCTGCAGGTG	GCAAGCTCTG	GTAAGTCACC	
<b>ል</b> ሮሞሞሮሞሮ <b>ል</b> ልሞ		TTGGCTATTA	ATCCCCTCCC	ACCOMOCHEC	31200
		GGAGATTACA			31200
		TATCAGTAAA		•	31300
		GGCTCACACC			21200
		CTTGAGCCCA			31400
		CTCTACAAAT			21400
		CCCAGCTACT			31500
		CAAGGCTGCA			31300
		AAAGTGAGAC		•	21.600
		GAACAGCATA			31600
		AGTTGAAAGT			. 21700
		AGCGGGAACC			31700
		CCTGCACTGA			21000
		GGAGAGTGAG			31800
		GGCTGAATCA			1 21000
		CCATTATGTT			31900
					20000
		GGTGGAAGCC	•		32000
		GCTCCGGATC			22122
		GAAAGACAGA			32100
		AAACTACTGG			2222
		AGAAGAGAAG			32200
		CATGAGATAA			
		ATGGTAAGCC			32300
		AGCATGGGAG			
		CAAACTCAGT			32400
		AGCTAGGAGG			
		TCCATCTCCT			32500
		GGGTACCCGG			
GCCTCCTGAG	CACAGAGAGG	GCAGAAAGGG		AGAGGGGCGC	32600
		FIGUI	RE 1A		

WO 01/04270	PCT/US00/19094

AGATAGTGGG	CAAATGGGGA	AGTGGCACTT	CCCCTAGCTC	GAGGGCAGAG	
GCTTGGTGTG	ATGGAATGGC	ACTCCTTAAA	CTGCTACATA	TTTTCCCTTT	32700
<b>AATŢTGGCCA</b>	AGAACAAGTT	GTCAAGTTTG	TGTGAGATAA	AGGTGCACTT	
GGTTCGTTCT	TGTCTAATGG	CCCCGCACC	CATGGGTATT	TCTTCAGCTT	32800
CCACAGTCAT	CCCGACACTA	GCTGGGAAGC	TCCAGCAGCC	CTGGTCCTGG	
CCCCAGCTCT	GTGGGCGCTG	GCCCTCAACT	TTGCCTGCAC	TGTGCTTTTG	32900
TGCTATTCCC	CTTGGTCCTG	TTTGGGTGCA	AGTCCCCCTC	ACGCATTGAG	
T			•	•	
TTCCTGGGCC	GCTCAGGCTG	CTCCTGTGTC	TCCCCAGGGA	ACATGAAGGT	33000
		4: 32988			
CTTGCAGGAG	•	TCTCCGACTA	CATGAGCATC	TCTACTTGCG	
		ACCAATTGCA		CCGCCTGTTG	33100
	•	CTCCGAGTAA		GAGCTGGAGG	
	331		С		
TTTGGGGAGG		AGGGTTTGCC	CCAAGAGTGA	GCTGGGTCCA	33200
		GATGCTGAGT		GCTGTTTATA	
		CATCTCCAGG			33300
		GTTGGTCCAC			
	•	AGTTCTCACA		TCAGGACCCT	33400
	_	TGAGGACCCA			
	TATGTTTACC		TTGCAATGAA		33500
	AAATTAAAAC		TTTTTTTCTT		00000
		GGCTGCTCTC			33600
		AAAGTGCTGG			03000
		AAATTTTAAG			33700
	-	TCAATTACAT			33700
		CCAGTGAGAG			33800
		AGAGTTTTAC			33000
			GAGGAGTACG		33900
· –		GATGGACCAG		•	
		CATGGTGCAC			34000
		ACACAAGACA			
		ATGCAGCGTT			34100
	CTAAGGTAGA		TTTGTTTCCA		
		GCCCGTGAGC			34200
	TAAAGTTTTA		CTGGCTGCTG	<del>-</del>	
AGAAACTGCA		TTATGTTTTT	TGTTTGTTTG	TTTGTTTGTT	34300
TGTTTTCTTT		TTCGCTCTTG	TTGCCCAGGC		
TGGCACAATC		GCAACCTCTG		TCAAGCGATT	34400
CTCCTGTCTC	AGCCTCCCGA	GTAGTTGGGA			
ACCCGGCTAA		TTTTAGTAGA			34500
		CTGAACTCAG	GTGATCCACC	CGCCTCAGCG	
				GGCCTCCTCC	34600
		TCAGAAATGG			
		AAATAATGTG			34700
		ATTGAAGACT			
		ATGGGCATAG			34800
		TTCACCACTA			
		AAAATTGCAC			34900
		CAGCCCTTTG			0.2200
		AGACCGGCCT			35000
		AATTAACTGG			
		CTGAGGCGGG			35100
		GAGATCACAT			20100
		GGAAAAAAA			35200
		TACAAATAAC			JJ200
		GTCCAAAATA			35300
			DE 1D	COOLUITAG	55500

FIGURE 1B

WO 01/0427	70				PCT/US00/19094
AGAAAAAGTT	TGCAGACCCC	TGGCCTAGCC	CGTGAGATGT	GGGTTGGCTG	
	ACATTGGAAT				35400
	TGTTCTAAAT				33400
	GAACAAGCTA				35500
	GCATCCTTGG				. 33300
					. 25.00
	CTAGTCGGCT				35600
	TCCTCACATC				
	GGCATGTCCC				35700
AGGTGACCAG	CCTAACCCAG	CCCCTGTGTC	TGCAGAGCCC		
				G	
•	5: 35736				•
CCCTGAGAAC	AACGGAGGCG	CGGGGTGCGT	GTGCCACCTG	CTCATGGATG	35800
	T A				
ACGTGGTCAG	TGCGGATAAC	TATACACTGG	ACCTGTGGGC	TGGGCAGCAG	
	C				
CTGCTGTGGA	AGGGCTCCTT	CAAGCCCAGC	GAGCATGGTG	AGCAGGGCGG	35900
	358	87]			
AGTGCGGCAG	GĠGTGGCTGG	GTGTGTTCCC	ACAGCTGCCT	GGGCTGAGGG	
T	•		•	T	
TGGGGTGGGC	AGGGGAGGAG	GTGGGGTCAT	AGCAACAGCA	GGAGGAAGCC	36000
A		•			
GCCTGTATTT	TCCCAAATCT	GATGGGATTC	CTGCCCCTGC	CTGGGCCTCA	
GTCCTCCCAC	CTTTGAAACG	GAGCTGGTCG	CAGTAGACCA	CCAAGCCCCC	36100
	CTGTTTCCAC				
	TGTGCTTGCT				36200
	CATTCCTTCC				. 30200
	CTTGACCGTG				36300
	ATCCCCCTTG				30300
	TGTATTCCAG				36400
	AGGTCTCATG				30400
	GAATCAAAAT				36500
	GTGGCCAAAG				30300
	GGTGTGGTGG		•	•	36600
	GCGGATCACC				20000
	AACCCCGTCT				36700
	CGCCTGTAAT				30700
ATCCCTTCAA	CCCAGGAGAA	CCACCTTCCA	CTCACCTTAC	AGGIAGGAGA	26000
	CCTGGGCGAC				36800
	ATAAAAAGAT		•		36000
	CTTTGCCTTC				36900
	CATGCCTGGA				27000
	AGCCCTGGGC				37000
	CAGCCTGGGC				. 22200
					37100
	GTAATTAATA		-		00000
	TATTATAATC			· ·	37200
	TATGCTATGG				
	GTAGGCCGGG				37300
TAGATCCTCA	CATAGAGGCC	GCTTCTCCCG	CAGTGAAACC	CAGGGCCCCA	
_		A		•	
_	6: 37334	•		•	
	CAGTTCACAC				37400
	TATCCCCCTG				
CAGTCAACAT	TTGGAGTGAA		CAGATGTGAG	TGGGCATGCT	37500
		T			
	374				
TTGACGTTTT	TCTGTGACCT	CTGGGGAACA	GGGTGGGTGA	CCAGCAGAGG	

FIGURE 1C

WO 01/0427	0				PCT/US00/19094
CCCAGTCCCT	GGAGCCAGGA	GCCTGGGAGG	CAAGCCCTGG		37600
CAAATCCCAG	GAGCTAGAGA	CCTGGCTTCT	CACCTGGCTC	A TGCCCTAGGC A	
<del>-</del>	GCTTCCTGGC	CCCCCACCCC	TCACATCAGA		37700
татстствса	тессестсст	<del></del>	GGTAGGGCTG	TGGGCCACAT.	
			ATCGTAGGTG		37800
			GAATAATACT		37000.
			CCAACGGGTT		37900
			CTCTTAATGA		57500
			AAAACAAGAT		38000
			TCCCTTCACA		30000
			AGAGTCTCAC		38100
			CACTGCAACC		20100
			CTGAGCAGCC		38200
•			TATATTTAGT		30200
• •			CTCCTGACCT		38300
			TACAGGTGTG		30300
			TGTATTGACA		38400
			TCAGTAATGG		30400
			TCTGCTGGTG		38500
			TGAATCCACC		30300
			GGGACCTGGG		38600
			CCAGCTGTGT		30000
			CCTCATCTGT		38700
			CAGTACAAGG		30700
			TCCTAAAAGC		38800
		•	CCCATCTGCC		30000
			TTGTGAGGGC		38900
			GCTCTCAGGG		30300
			TCACTGCTTC		39000
			TCAGGTCTGG		. 33000
			GAATGGGGCA		39100
			GGGGGAAGGA		33100
			AATAAATCTC		39200
			CCAAGGATGG		33200
			ATATGTTATT		39300
			GGATCCAGCT		33300
•			ATTGGCTTCG		39400
		=	CTACTAGTTG		. 33400
			TTTTTTCTTG		39500
•			TGGCTTCGAT		33300
			GAGAGGGTGC		39600
			TGCCTCTAGG		
			AAAATAAGGC		39700
			GGACTGTGCT		33700
			CCAGTCATGG		39800
			ATGTTGGTGC		33000
			GAAGAATGAA		39900
			ACCCCATCTC		33300
			ATTGGCCAGA		40000
			MCVMCCCVMW		40000

FIGURE 1D

ATGCTCCCCT CTGATGAATG ATAGGCCAGG TGATGGGATT GGCACAATTG GCTTAGACTA ATGAGGGTTG GCCCTGGAGT TGCAGGCAGT GGAGTTCTGT

CCTAAGCAGT GGGCACCTAA ACCCGATGGC ATAAAAGCTG GGCGGGTGTC CACCTGCATC TGCCACAGCA CTATAGGCAC CAACTGTGGC TCATACTGAG

TGGGATAAAT TCCAGAAAGA AACATTAGGA ACTTACTATA GAATTTTGGG

40100

40200°

WO 01/0427	70				PCT/US00/19094
GCTAGAGCTA	CTCATTCATT	CCCCTAGATÃ	ATTTCTAGGC	AAGGTTCCAT	40300
AGTGGAGGGG	GAGTTTTGGC	TTGGGCATTG	AAGGATGCAT	AGGAGTTTTC	
TAGATGGGGA	AAGAAGGGAA	CGGTAGACCA	GGCAGAGGGA	ACTGCATGAT	40400
AAAAGGTTTA	TGGGTGTGAA	AATTCATGGA-	ATGTTTGÄGG	ATTATGGGGT	
TGGGGGATGT	GGGAATATGT	GTAGCGATAA	AGCACCAAAC	AAAGCCAAAA	40500
		GCCTGCCTCA			
_				GCTGGAAATG	40600
		TTGTTGTTGT			40000
		CAGGGGCGTG			40700
		ATTCTCCCAC			40700
		CACGCCAGGC			40800
		TTGTCCGGGC			40000
		GCCTCCCAAG			40000
					40900
		ATCTGAGATT			41000
		CTCCCATGGA			. 41000
		TGTTAAAAAA			
		CTTTTGTTTC			41100
		GTAAGATTTC			
		TGTTTAGAAA			41200
		GTCTCCCCAC			
		TCTCTGAACT			41300
		TTCTTGTGAA			
		ATGTGCTTCC			41400
		ACTGCCGGGG			•
GGTGCATAGC	CCTGGATCTC	GCAGGGAGAG	GGGTGACAGG	ATGAGAACTC	. 41500
AGGTTGCTCA	CTTGCCATCA	GGGTCAGTCA	TGAATATAGC	GTTCATGTAT	
CACTTTTTAA	AGCTTTTTTG	GAGGGTAAAA	GTAATAGTTA	CACAAAATAA	41600
AAATACAAAT	GGTACAAAAG	GACTTAGAAT	GGAAACATGT	TTCTCTCCCG	
ACTCCAGCCT	CCTGTTTTTC	TTCCCAGAGA	CTGACCACTG	CTGTCTGTCT	41700
CTTGCCAGAA	GGGAAAGGGA	GGCAAGGTTA	GGGCAGGCAG	AGGGCATGTG	•
CATCCTTTAG	AGAGAGCTTA	TGTCTATACA	AGCAAATGTG	TGTGTTCAGT	. 41800
CATCGCTGTC	TTAGTTTTCT	ATTGCTGCAT	AATAATGGTA	CTACCAGCTT	
CACAGCTTTA	AACAACACCC	ATTTATTATC	TCATAGTTTC	TGTGGTTGGG	41900
AGTCTGGACA	TAGCTTAGCC	AGGTTCTCTG	CTTTAGAGTC	TCGTGAGGCT	
ATAATCAAGG	TGTGGGATGG	GGCTGCAGTT	TCATCTGAGG	CTCAATTGGG	42000
GAAGGGTCAC	TTCTAAGCTC	ATACAATATT	GGTGACATTC	AGTCCCTGGC	
		TCAGTTTCGT			42100.
		GTGCCCTTTG			
		TATTACAGTC			. 42200
		CCATCACCTT	_	•	,
		CCACACTCGA			42300
		CATCGGGGCC			12000
		CTCTCTCTTT			42400
		CCCAGGCTGG			
		CCCAGGTTCA			42500
		CAGGTGCCCG			12500
		GAGTTTCACC			42600
		TCCACCCACT			42000
•		•		TTAAAAAAACA	42700
		AAAAAGAAGC			42700
					42000
		ACAATAGATC			42800
		ATTGGATTCT			. 40000
	•			GATGCCTCGA .	42900
		GATCACCCAG			400
		TATTAACCCA			43000
		CTTTTATTAT			
CATTGAGGTC	CAGAGAGGTT	ATGTCACTTA	CTTAAGGTCA	CACAGCCAGG	43100

FIGURE 1E

WO 01/0427	70				PCT/US00/19094
<b>አ አ ሮምሮሮሞአ ር</b> ሞ	እርርርእርጥርጥጥ	እርርርሞምርሞሞሞ	TACAGATGAG	እምም <i>ር</i> እእምምልም	
			TGCCTAAGTA		43200
			CTAGCCTGGT		43200
					42200
			AACAACCAAA		43300
			ACGACAGCAA		42400
			CTAACATCTC		43400
ACCAGTTCAG	AATCTATAAC	GTGACCTACC	TAGAACCCTC		
lexon	7: 43406		•	A	
•		TGGGATTTCC	TACAGGGCAC	GGGTGAGGGC	43500 .
			TGAGTGGAGC		15500 .
			AAGCAATGGT		43600
MULGUCACAA	435		MOCANIGGI	MICICCACI	43000
СТССАТТСТТ		•	CCCTGGCTGA	GTCTCTGGGC	
			TCCTGCCATT		43700
			AGGCTGCAGT		43700
C	ATCTGCCAGG	INGGCNGGCI	AGGCIGCAGI	·	
CCCACAATTT	CCATGGCTTA	ACACTATAGG	AATATATTTC	TTGCTCATGT.	43800
AACAAGCTAA	CGTGAATGTT	GCTGGTTTGT	AGGTGGTTTC	CCTCCCTGTA	
GAAATCTGGG	GAGTGAGGTT	CTTTCCATCT	TGTGGTGCCA	TCATTCTCCA	43900
			TGGTTTCCTT		
			TTTTAAATAC		44000
			GTACAAAAAT		
			GGGAAGGCTG		44100
			GTGAGCCGTG		
		•	CCCATCTCTA		44200
			ACTACAGAGG		14200
			ACATTTTTAA		44300
			ATAAGTTCTA		. 44500
			GCATAAATAA		44400
			CTGTGAGTTT		,11100
	· · · · · · · · · · · · · · · · · · ·		GGGTCTCTTC		44500
			AATTTTAA.	-	. 4300
			TTTTGACCTC		44600
			CTGGGATTAC	•	0000
			ATGGAAGGAG		44700
-			TCTTTCCCAT		44700
			GGTCCATCAA		44800
			CTGTATCCAG		44000
			TTGAAAGCAG		44900
					. 44900
			ATTGGTGAAA ACTTTCTCAC		45000 4
					45000 ·
			ATCCCACAGG		. 45100
			AGTCTTACAA		45100
		•	CTTCCCTTCT		
			ATCCTTAGAC		45200
			TACATAAAGC		
			TGCATCTCTT	•	45300
			GCAAAATTCC		. = . = -
			TTACTGTTAT		45400
			GGAAAAAGGA		
			GGCTCCCTGA		45500
			TTCAGTTTCC		
			TCAATGAAAG		45600
AACGTGGTAG	GTGCAGTGGC	AGCTGGCAGT	ACCCTGACCC	TGCCACCGCA	
CAGCCCTCTC	AGCATTGCTC	ATCCTGCACT	GTGGATATCA	GTTGAGCCAC	45700
GTGTCTCCTG	CCCTGGGCTG	TGAGCTCCAT	AGGCAGGGTC	TCCATGGCTG	
			0 P 1 P		

FIGURE 1F

		GAACCAGGTG			45800
		TGGGGAAGGG			•
ATAACAATGT	ATGGGACCCT	CATTCATTAT	TTCAGCAAAT	ATTTGCTGAG	45900
TTCCTCCTAC	ATGGCTAGCC	CTGTGCTAGA	CACTGGGGAA	TCGGCGATGA	•
ACAAAGCAGA	TAGAAATCCC	CACTCTTGTG	GAGCTGACAT	TCTGGAGĢGA	46000
GAGACAAAAA	GCAAACATAT	AAAGAAAGAA	AGAAATCACA	TGGATCTGGA	
		AAAATAAAAG			46100
ATGGGCAGGG	GGCAACGGTA	GGGAGGTGT	CGGGGAAAAC	TTTTTGGAGA	
ATGTGACGAT	GAAAGTGAAC	AAGGAGAAGT	CAACCGTGTT	GAGATGATGG	46200
CAGCTAATGA	TGTGGACAGG	CCACTCTGTT	CTGAGTGCAT	TATCTATTGA	
TTCATCATGT	CATCCTCGCA	ACAGCCCTGC	ACGATCAATT	CTGTCATTAA	46300
CCCCATAGTA	CAGATGAGGA	TGCGGAGGCA	CAGAGAAGAT	AAGGGACTTG	
TCCTGTGTCA	CACAGCAAGG	AGCCATCCGG	CTCCTAAGTT	GGTGCATTTG	46400
ACTTCTGTGC	TTCCGGAAAG	AAAGAGCAGC	AAGTTTAAGA	TCTGGAGGTG	
GCACTGAGCT	TTGGAGGAGC	AGGGGGCAAT	GAGGTGGCCG	GTGTGACGAG	46500
GACTCAATGT	GCAAGAGGGA	GAGTGGTGGG	GAGATGAGGT	GGAGGGGTGG	
TCGGCGGTCA	GATCGTGGAG	GGTCTCGGAC	GAGGGTCCTG	ACCCTGGGTC	46600
TCCAGTCCTG	GGAAGTGĠAG	CCCAGGCTGT	ACCATGGCTG	ACCTCAGCTC	•
		CAGCCTACAG			46700
	8: 46674	•			
		TCCTGCATTG	TCATCCTGGC	CGTCTGCCTG	•
		CAAGTGAGTC			46800
	467			•	•
CAGTCCCTCT	•	GTGGCAGGGA	CTTGCCCCTC	TAGTCTGCCC	
		ATAATACGTA			46900
	•	GAACATATCG			
		TTTTGAGACG			47000
		ATCTTGCCTA			
		CTCAGCCTCC			47100 ·
-		TAATTTTTTG		•	
		TGGTCTCGAA			47200
		GCTGGGATTA			
		TAAAACCGGC		•	47300
•	· - <del>-</del>	AGGTGGTTTT			•
		AGGCCAGAGA			47400
		TCTCCAAGGA			
		GGCCAGGGAT			47500
•		GTGCCTGGGC			
		ATTTGCGACG			47600
		GCCTTGGAGC	•		
		CTGGGGTTCC			47700
		CAGCCAGAAG			
		CGAGTGGCTG			47800
		CAGCCCTTGG			
		ATGGGACTAG			47900
		GATTATTAAC			2,200
		AGGTAGGCAG			48000
		CGAGCAACCT			
		CCTGTACCTC			48100
•		TTGCTGTTCT			20200
		GTCTCACTCT			48200
		TGCAGCCTCC			30200
_		AGCAGCTGGG			48300
		TTTAATTTAA			20200
		CAAGCTAGAG			48400
		CGGGTTCAAG	·· · · · - · - ·		40400
					40500
CCCGACTAGC	TOGGATTACA	GGCGTGTGCC		GCIMMITTI	48500
		777.777	DE 10		

FIGURE 1G

		GTTTCACCAT			
		CACCCACCTC			48600
	- · · · · · · · ·	CTGGCCAATT		TTGTAGAGAC	•
		CAGGCTGATC			48700
· ·		ACCATGAATG		•	
	TTTTTTTTT		TGGAGTCTCA		· 48800
		CGATCTCAGC		CTCCGCCTTC	
		GCCTCAGCCT		TGGGATTGCA	48900
		GCTAAGTTTT			
		CTGGTCTCGA		TCAGGTGATA	49000
		AGTGCTGGGA		GAGCCGCCAT	40-0-
	ATATAAGCAT	TTCTGTCTCC		CATCTTTCCC	49100
		AAGGCAGGCA			
		AGTGAGTCAA ACCATGGGCA		TGCTCAATAA	49200
		TGATGATGCA		CAGAAGCGGT	40200
	AGGGTTGTGT				49300
		TTCTCCGTAG		TTCCCAGGCC	40400
		TCCTCTCTCT		TCTTTCCTAA	49400
		ACTGAGGACA			40500
		GCGGGAGTGT			49500
		TTGGGCCTCA			40.600
		GTCACCAGGT			49600
		AGCCTGATCT			49700
		GGGAGGAGCG			49700
		TGTTTTTCAG			49800
	9: 49781	1011111000	OHI IMMOHM	OMMICGIGG	45000
-		CGCAGCCGCC	TCGTGGCTAT	аатаатссас	
		CGTGGATGAG			49900
	498				
TGAAGAAGTG	TGGTTCAGAA	CACCTGGGCT	GTTAAGGACC	TTCACTGGCT	
		GTCAGGAGGG		GACAGAGGCA	50000
GAAGCCGAAT	GAGGTCATTA	GCAĠACCAGA	GGCTTTCCCG	CCCTTCCCCT	
TGGCAATCCC	AGCCTGGGGT	GGGCTTCTCT	GGGGTTGGTT	TCCTGTTTTT	50100
TTCCCTCCCC	TTGGGAGAAT	GACCCTTGGG	TCATCATCAC	TGTGTCATTC	
CCTGGGGAGG	TGCCAGTACC	AGGGCTAGAG	GCCAGAAGGA	GTGGAGGAAG	50200
		GTGTCTTCTT			
CCCGAAGCAC	TAGCAAATCC	CTTCCGGGTT	CTCATTGGCC	TGAAATGTAT	50300
		GGAGTCAGCT			
		GCCTGAGCAA			50400
		CCTGTCTAGA			
		AATCTAGTGT			50500
		ACTTGAGCAA			•
		ATAACAGTGT			50600
		CATATAAAGT			
		TATCAGTTAG			50700
		ACAACAACTT			
		GCTCAGCAGT			50800
		ATCTGAATTG			
		TGGCTTTTTT			50900
		CCCAGGAGTG			
		GGTTCAAGCA			51000
	GGATTACAGG	CATGTGCCAC			
	707m000	mamma			
	AGATGGGGTT				51100
CCTAATATCA	GATGATCCAC	CCACCTCAGC	CTCCCAAAGT	GCTGGGATTA	
CCTAATATCA CAGGCGTGAG	GATGATCCAC CCACTGCACC	CCACCTCAGC CAGCCTAGTT	CTCCCAAAGT GGCTGACTTT	GCTGGGATTA TACCTGGGAC	51100 51200
CCTAATATCA CAGGCGTGAG	GATGATCCAC CCACTGCACC	CCACCTCAGC CAGCCTAGTT GTGCCTCTCA	CTCCCAAAGT GGCTGACTTT	GCTGGGATTA TACCTGGGAC	

FIGURE 1H

WO 01/0427	0				PCT/US00/19094
GGCTTGTTTA	CAGAGTGGCT	CAGTTTTCAA	GGGTGGGAAG	TCCCAAGGCT	51300
TCTTGAGGCC	TAGGCGCAGC	ACTGGCATGA	TATCACTTCC	ATCACATTCT	
ATGGGCCCAA	GCAAGTCCCA'	GGGCCAGTGT	AGATTCAAGG	ĠATGGGAGGA	51400
			GCCATCGACC		
			GGAATCTGAG		51500
			GTGATGTCGA		21200
				CCACTCTGCT	51600
					51600
			TCACAGTGGG	AGAAGCGGTC	•
-	10: 51628.				51000
CCGAGGCCAG	516	_	TGTATCTGAA	CTTAGGTCAC	51700
AGCCTGCATG	CATTGGGAAG	GTGATAGAAT	TGGAGAGGCA	AGCCCCTAGC	
TCCATGTCTG	CCTTCTCTTC	CCTGCATTCG	GTAATTGCCC	TGTGACATTA	51800
GCCTTCAAGG	GACGGCAGGA	GGAGGGGTGT	TCTGGAAACG	TGGACTGCTG	
			AGGTACATGG		51900
			CAGCCGTGCC		
•			ATAGCCACTA		52000
			CACTCTGTGG	-	
			CCTTCAAACT		52100
AGTGATCCTC	· · · · · · · · · · · · · · · · ·				. 52100
			TGCCTGTAAT		52200
			CCCGGGAGGC		32200
			CCTGGGCGAC		
				•	52300
		•	AGTTTTTGTA		
			CTTAGCTTCT		52400
			AAGGGGGTCC		50500
			TTTTTTTTT		52500
			GCAGTGATTC		50500
•			CTCAGCCTTC		52600
			TAATTTTCCT	•	
			TCTGTTGCCC		52700
			CTCAGCCTCC		
			TTCCATATCC		52800
			AATTATCATC		
			TATTTGGTGC		52900
			ACCCTACCCT		
TCATTCTAGC			TGATTATAAC		53000
ATGTCTGAAG C	TAGACAGCCA	TCAGGACATG	GTGATTTCAG	GCTGGGCTTT	
GAAGAATGAA	TAGGAGTTTT	TCAAGTGTCG	AAACTGAACC	CTGACCAACC	53100
<b>դուն Երափան</b>	CACACACTCC	<b>አአርአአጥጥርጥር</b>	TTACCAAGCT	Charceconeda T	•
	11: 53114.		TINCCHAGCI	CIIGCCCIGI	
lexon	11: 55114.	•		•	
TTTCTGGAGC C	ACAACATGAA	AAGGGATGAA	GATCCTCACA	AGGCTGCCAA	53200
AGAGATGCCT	TTCCAGGGCT	CTGGAAAATC	AGCATGGTGC	CCAGTGGAGA	
TCAGCAAGAC	AGTCCTCTGG	CCAGAGAGCA	TCAGCGTGGT	GCGATGTGTG	53300
GAGTTGTTTG	AGGCCCCGGT	GGAGTGTGAG	GAGGAGGAGG	AGGTAGAGGA	
AGAAAAAGGG	AGCTTCTGTG	CATCGCCTGA	GAGCAGCAGG	GATGACTTCC	53400
ÄGGAGGGAAG	GGAGGGCATT C	GTGGCCCGGC	TAACAGAGAG	CCTGTTCCTG	
GACCTGCTCG T	_	TGGGGGCTTT	TGCCAGCAGG	ACATGGGGGA	53500
_	CTTCCACCTT	CGGGAAGTAC	GAĠTGCTCAC	ATGCCCTGGG	
		CCCAAGGAGG	CACCTCCCTG	GGGCAAGGAG	53600
			RE 1I		<b>- 300</b>

	WO 01/0427	0				PCT/US00/19094
	CAGCCTCTCC	ACCTGGAGCC	AAGTCCTCCT	GCCAGCCCGA	CCCAGAGTCC	
			AGACGCCCCT			53700
			TCCCTGAGCC			
		022001210	C			
	CTGGGTCCAG	ACCCACTGCT	GGCCAGACAC	CTGGAGGAAG	TAGAACCCGA	53800
	GATGCCCTGT	GTCCCCCAGC	TCTCTGAGCC	AACCACTGTG	CCCCAACCTG	
	AGCCAGAAAC	CTGGGAGCAG	ATCCTCCGCC	GAAATGTCCT	CCAGCATGGG	53900
	GCAGCTGCAG	CCCCCGTCTC	GGCCCCCACC	AGTGGCTATC	AGGAGTTTGT	•
		T			G A	
	ACATGCGGTG	GAGCAGGGTG	GCACCCAGGC	CAGTGCGGTG	GTGGGCTTGG	. 54000
			TACAAGGCCT			
			ATGTGGGTTT			54100 <sub>.</sub>
			ACCTCATTCC			
			TTCACCTTTG			54200
	CGCAGTCCGC	AGAGCTCACA	TCTCCCAAGC		AGCACCTGGG	
				T		r 4000
			TAGAGGACAT			• 54300
			CTTGTGGACA			E 4 4 0 0
			CCTGTGCGGC			54400
			CCCCTGTCAT			54500
	GCTGCTGTGG	AGACAGGICC G	ICGCCCCCIA	CAACCCCCT	GAGGGCCCCA	34300
	GACCCCTCTC		TCCACTGGAG	GCCAGTCTGT	GTCCGGCCTC	
			CAGAGAAGAG			54600
			CAGAGCTCAA			
		C .				•
	AACTTTGTCT	CCGTGGGACC	CACATACATG	AGGGTCTCTT	AGGTGCATGT	54700 ·
		٠	•	•	CC	•
		546				
	CCTCTTGTTG	CTGAGTCTGC	AGATGAGGAC	TAGGGCTTAT		•
	CARAMCCCAC	CHCCHCCAAC	GCAGCCAGGC	mccca ca mmm	T	E 4000
	GAAATGCCAC	CICCIGGAAG	GCAGCCAGGC	TGGCAGATTI	CCAAAAGACI	54800
	TCAACAACCA	тсстателас	GTGATTGGCC	ССУСТСУССТ	TCCCCTAACA	
			CCCGCCCAGC			54900
	•		CTGGGTCGCC			3130,0
			GGGCACCTCG			55000
			GCAGCAGACT			
			ATCATGGCCC			55100
			•	G		
	CTGTCTCACT	GAACTAGAAG	CCGAGCCTAG	AAACTAACAC	AGCCATCAAG	
					A	
			GGGAAATCGA			55200
			GTGCTCATTC			
			CCCGGCTGTC			55300
	GAGCAACAGA	GGACATGAAA	AATTGCTGTG	ACTAAAGCAG	GGACAATTTG	
	amaaaa	~~~~~~~~~~	A .		macmamacaam	
			GCTGTATGGC			55400
	GGAACCCCCA	GAATAAATAT	GCTCAGCCAC T	CCTGTGGGCC	GGGCAATCCA	
	CACACCACCC	ATAACCCACC	AGTTACCCTG	CATETTECCC	CACACCTCAG	55500
			CTTGGGTTGA			33300
	CIGCINGGON	1.0000001210	CITOGGIIGN	T	70101011	•
	TTAGTTTCAT	CACCTGTTAT	CTGTGTTTGC		GGAACAGAAG	55600
			AAAGTTTCTT			
			GACACTGCTG			55700
			CAGAGGAACC			
				RE 1J		
٠						

WO 01/04270	PCT/US00/19094
-------------	----------------

ACTTGCCGTC A	TGGGTTTGGG	TTCCCCATAC	AAGGATTCAA	ATAGTTGATT	55800
	<b>ፐርርርርርር እ</b> እአ	СССТССТААС	GTAGTGGGGA	ACTGAGGCAG	
· · · · · · · · · · · · · · · · · · ·			GAAAGGGGTC		55900
					33900
			AGAAAGAATT		F CO O O
			GAAAGTAAAG		56000
			GGGTTGCTGG		
TTATGGTTAT	TGATTAATTA	TATTCCAAAC	AAGGGGTGGA	TTATTATGCC	56100
TCCCTTTTAG	ACCATATAGG	GTAACTTCCT	GATGTTGCCA	TGGCATTTGT	
AAACTGTCAT	GGCGCTGTTG	GGAGTGTAGC	AGTGAGGACA	ACCAGAGGTC	56200
ACTCTTGTTG	CCATCTTGGT	TTTGGTGGGT	TAGAGCCATC	TTCTTTACTG	
CAACCTGTTT	TATCAGCAAG	GTCTTTATGA	CTTGTATCGG.	TGACGACCTC	56300
			ACCTCCCAGG		•
			CTCTTCAAAG		56400
				CCAGCATGAG	**.
			GAAGACCATG		56500
			TGGGGTATTT		50500
			CAGGAGCATT		56600
					20000
			TCTTCAACTG		5.5700
			CAGGAACAGG		56700
	•		GCTAGTTAGC		
	**		CAGGCTCTCT		56800
			GGTGTGGGCC		
AGGCTGGGAA	CCTGAAGCTC	CTATGTCTGT	ATCACTTTTG	CTTCTCTGAG	56900
TAGCTGCCCT	GATTTCACAC	TTGAGGGGCT	TGGCCATTTT	AGATTCCTTC	
CTGCTCTAGG	AGCCTACATA	CTACACTGGA	AATGATGGGG	AGCTCTCTAC	57000
CTCACATGCA	GCCTGATGTT	TGTTAGAAAC	ACCTCCTTGC	GCCAGGCATG	
ATGGCTCATG	GCTGTAATCC	CAGCAATTTG	GGAGGCTGAG	GCGGGTGTAT	57100
			GGCCAATATG		
			GTGTGGTGGT	•	57200
			AGAATTGCTT		
			CATTGCACTC		57300
			AAAAAAAAAAC		27300
					E7400
			TGCCCAAATC		57400
			GGGACTTGCC		F7F00
			TGGACTTCTT		. 57500
			AAAATAGTTA	•	
			ACTGTGTCAT		57600
AAACAGCTCT	AGTTGGGAGG	CTCAATGATT	ATCCCAATTT	TACAGATAAG	
GAAACAGGTC	CAGAGAGGTT	GAGGATTAGC	CTAGAACCAC	ACAGCTAGGA	57700
AATCCTGGAG	CCAGGATTTG	AACCCGGGTC	TGACCTAAGA	GCTCCCAGCC	
GCCGTGATAT	ATCAGCTTAT	GTCATCCTGA	CACCTACGCA	GATGTCGGCT	57800
CGAATCCACT	TTGCCTGAGC	ATTGTCTCAG	AGAAATCTAA	TTTAAAAATT	
AGGCAGCAAA	TAGAAAATAT	ATTTGACTGC	TAGAGATGCA	ATGGGACTGG	57900
GAGCCCAACA	AAGGATCTTA	GGCAAAAGAA	ATCCAAGTTG	TTGGCCTCAG	
			AAGCTACAGA		58000
			GCTGGCCCCC		
			GTGAGATGCC		58100
_			AGGAGGGGAC		
			GCCCCTGCGG		58200
			CCCTGCCTTT		30200
					E0200
			GAGGAAGGCA		58300
			GGCTTATTGC		
			TATTCAGAGA		58400
			GGAAATCCCA		
			ATGAGCGCTT		58500
GCGTAGTATC	TCCGTGAACA	TCCGGTTGAA	CAACCTTTCT	GACTTTATTT	

FIGURE 1K

TTCCCACGAA AGTTATTAAT TAAAAAACAA AAAGCAAAAC ACCGAAAAAA 58600 CAAAAAAACCC AGCAAGTGTT TGAGCTCCCA CCACGAGGGA GGCCTGACGT CACTGGATCC TCCCGGCAGC CGATGAGGCT GCATGGGACT 58690

# POLYMORPHISMS IN THE CODING SEQUENCE OF IL4Ra

				•	
	TTTGCTCTGG				
	GCAAGCTCTG			,	100
GCGTCTCCGA	CTACATGAGC	ATCTCTACTT	GCGAGTGGAA	GATGAATGGT	
CCCACCAATT	GCAGCACCGA	GCTCCGCCTG	TTGTACCAGC	TGGTTTTTCT	200
GCTCTCCGAA	GCCCACACGT	GTATCCCTGA	GAACAACGGA	GGCGCGGGT	
		G	${f T}$	A	•
GCGTGTGCCA	CCTGCTCATG	GATGACGTGG	TCAGTGCGGA	TAACTATACA	300
				C	
CTGGACCTGT	GGGCTGGGCA	GCAGCTGCTG	TGGAAGGGCT	CCTTCAAGCC	
CAGCGAGCAT	GTGAAACCCA	GGGCCCCAGG	AAACCTGACA	GTTCACACCA	400
ATGTCTCCGA	CACTCTGCTG	CTGACCTGGA	GCAACCCGTA	TCCCCCTGAC	
AATTACCTGT	ATAATCATCT	CACCTATGCA	GTCAACATTT	GGAGTGAAAA	. 500
CGACCCGGCA	GATTTCAGAA	TCTATAACGT	GACCTACCTA	GAACCCTCCC	
T					•
TCCGCATCGC	AGCCAGCACC	CTGAAGTCTG	GGATTTCCTA	CAGGGCACGG	600
A	•	•			
GTGAGGGCCT	GGGCTCAGTG	CTATAACACC	ACCTGGAGTG	AGTGGAGCCC	
CAGCACCAAG	TGGCACAACT	CCTACAGGGA	GCCCTTCGAG	CAGCACCTCC	700
TGCTGGGCGT	CAGCGTTTCC	TGCATTGTCA	TCCTGGCCGT	CTGCCTGTTG	
	GCATCACCAA				800
CAACCCAGCC	CGCAGCCGCC	TCGTGGCTAT	AATAATCCAG	GATGCTCAGG	
	GGAGAAGCGG				900
	ATTGTCTTAC				
			C		
CATGAAAAGG	GATGAAGATC	CTCACAAGGC	TGCCAAAGAG	ATGCCTTTCC	1000
	AAAATCAGCA				
	AGAGCATCAG				1100
	TGTGAGGAGG				
	GCCTGAGAGC				1200
10101001110	000101101100			C	1200
GGCATTGTGG	CCCGGCTAAC	AGAGAGCCTG	TTCCTGGACC	_	
0001110100			1100100100	Т	•
GGAGAATGGG	GGCTTTTGCC	ACCAGGACAT	GGGGGAGTCA	-	1300
0011011111000	OCCITITOCO	MOMONIA	000001101011	C T C	1300
САССТТСССС	AAGTACGAGT	GCTCACATGC	ССТСССАТСА		•
	AGGAGGCACC	•			.1400
	CCTCCTGCCA				. 1400
	GCCCCTCGTC			_	1500
	TGAGCCAGTC				1300
C	TONOCCAGIC	ACCOLORCCC	AGAGAGCIGG	GICCAGACCC	•
•	AGACACCTGG	ACCAACTACA	አርርርርእርእሞር	CCCTCTCTCTC	1600
	TGAGCCAACC				1600
			•		1700
	TCCGCCGAAA				1700
T	CCCACCAGTG		•	GCGGTGGAGC	
<del></del>	CCACCCCACI	G	A		1000
	CCAGGCCAGT				1800
	AGGCCTTCTC				
	GGGTTTGGGG				1900
	CATTCCTGGC			•	***
	CCTTTGGACT				2000
CTCACATCTC	CCAAGCAGCT		CCTGGGTCTG	GAGCCGGGGG	
		T	·		_
AAAAGGTAGA	GGACATGCCA			GCAGGCCACA	2100
		FIGU	RE 2A		

# WO 01/04270 GACCCCTTG TGGACAGCCT GGGCAGTGGC ATTGTCTACT CAGCCCTTAC CTGCCACCTG TGCGCCACC TGAAACAGTG TCATGGCCAG GAGGATGGTG 2200 GCCAGACCCC TGTCATGGCC AGTCCTTGCT GTGGCTGCTG CTGTGGAGAC AGGTCCTCGC CCCCTACAAC CCCCCTGAGG GCCCCAGACC CCTCTCCAGG 2300 G TGGGGTTCCA CTGGAGGCCA GTCTGTGTC GGCCTCCTG GCACCCTCGG GCATCTCAGA GAAGAGTAAA TCCTCATCAT CCTTCCATCC TGCCCCTGGC 2400 C AATGCTCAGA GCTCAAGCCA GACCCCCAAA ATCGTGAACT TTGTCTCCGT GGGACCCACA TACATGAGGG TCTCTT 2476

# ISOFORMS OF THE IL4RG PROTEIN

MGWLCSGLLF	PVSCLVLLQV	ASSGNMKVLQ	EPTCVSDYMS	ISTCEWKMNG	
PTNCSTELRL	LYQLVFLLSE	AHTCIPENNG	GAGCVCHLLM	DDVVSADNYT	100
		V	T		
LDLWAGQQLL	WKGSFKPSEH	VKPRAPGNLT	VHTNVSDTLL	LTWSNPYPPD	•
NYLYNHLTYA	VNIWSENDPA	DFRIYNVTYL	EPSLRIAAST	LKSGISYRAR	200
			Н		
VRAWAQCYNT	TWSEWSPSTK	WHNSYREPFE	QHLLLGVSVS	CIVILAVCLL	
CYVSITKIKK	EWWDQIPNPA	RSRLVAIIIQ	DAQGSQWEKR	SRGQEPAKCP	300
HWKNCLTKLL	PCFLEHNMKR	DEDPHKAAKE	MPFQGSGKSA	WCPVEISKTV	
LWPESISVVR	CVELFEAPVE	CEEEEEVEEE	KGSFCASPES	SRDDFQEGRE '	400
				.A.	•
GIVARLTESL	FLDLLGEENG	GFCQQDMGES	CLLPPSGSTS	AHMPWDEFPS	
			R		
AGPKEAPPWG	KEQPLHLEPS	PPASPTQSPD	NLTCTETPLV	IAGNPAYRSE	500
SNSLSQSPCP	RELGPDPLLA	RHLEEVEPEM	PCVPQLSEPT	TVPQPEPETW	
<b>P</b> ·					
EQILRRNVLQ	HGAAAAPVSA	PTSGYQEFVH	<b>AVEQGGTQAS</b>	AVVGLGPPGE	600
		R I			
AGYKAFSSLL	ASSAVSPEKC	GFGASSGEEG	YKPFQDLIPG	CPGDPAPVPV	
PLFTFGLDRE	PPRSPQSSHL	PSSSPEHLGL	EPGEKVEDMP	KPPLPQEQAT	700
		S	`		
	IVYSALTCHL				
RSSPPTTPLR	APDPSPGGVP	LEASLCPASL	APSGISEKSK	SSSSFHPAPG	800
A					
NAQSSSQTPK	IVNFVSVGPT	YMRVS	•		825

### SEQUENCE LISTING.

<110> Genaissance Pharmaceuticals
Denton, R. Rex
Nandabalan, Krishnan
Chew, Anne
Stephens, J. Claiborne
Duda, Amy
Windemuth, Andreas

<120> Drug Target Isogenes: Polymorphisms in the Interleukin
4 Receptor Alpha Gene

<130> MWH-0002 PCT IL4R alpha

<140> TBA

<141> 2000-07-12

<150> 60/143,435

<151> 1999-07-13

<160> 413

<170> PatentIn Ver. 2.1

<210> 1

<211> 28690

<212> DNA

<213> Homo sapiens

# <400> 1

tgtgagctac tgtgtctggc ctgaataata aaatttaaaa caatttttca aaaattcacc 60 atgaggtete actatattee etaggetggt eteaaaceee tggaeteeaa gtgateeace 120 ccaccttccc gagtagctgg gactagagat gcacaccatt gcacccaata gagcaatacg 180 tttctgttct ttgtaaatta cctgctctaa ggtatttttg ttatagcagc ctatatggac 240 taagetgact tgtaaegtta ettgagactt taaagtgtte eggteaetgt tgqagggete 300 tgtctgtgtt agctcattta atccccacaa cacctcaatc agatggggct attcttagtc 360 ccactttata gataaggaaa ctgaggcatg gaagcacagc ttgctcaagg ttcacatcta 420 gtcagtgaca gagcaggtat ttaaacctca ggaaataatc agagaaacat gtgtagaggg 480 ttgtccaagg aaggccacat ccagaagcat ctcccaggac agttgttgtg tagctcaccc 540 totggacttt gtgggtotgg gtgttgtttc atgattatag agagagetet gtgaacgtgg 600 aggacetgtt gteggeagag acacaaatgg ceagggeatg getgggeage egeagtgget 660 caggcctgta atcccagcac ttcgagaaga ccagaggggc agatcatgag gtcagaagtt 720 caagaccage etggecaaca tggtgaaace eegtetetae taaaaataca aaaattagee 780 aggtgtggtg gtgggcacct gtaatcccag ctactcggga ggctgaggca gaagaatcgc 840 ttgaacccgg gaggtggagg ttgcagtgag ctgagattgc accactgcac tccagccttg 900 gagacagagc gagactetgt eteggaaaaa caaacaaaca agcaaacaaa caaacaaata 960 aatggccagg gcaggggagg gttgcatatt gaataagatg agctctgctg gaagcacagg 1020

tcagcactaa	cctgcttcct	ctctctctgc	aggtgccttg	gcatctccca	atggggtggc	1080
	gctcctgttc					
gtaagtcacc	acttctcaat	cattcatttg	ttggctatta	atggcgtgcc	agggtcctgc	1200
agtatgtcac	ctggccttat	ggagattaca	ctgcagtggg	aggggacagc	caatgacaag	1260
tggccctgat	tatcagtaaa	ttctaaagat	tgttagaaag	tgatgggagc	cgggtgcagt	1320
	tgtaatccca					
	gtcagcttgg					
tagccaggtg	tggcagtgca	cgcctgtagc	cccagctact	caggaggccg	aggtgggagg	1500
	ctcaggaggt					
cctgcgtgag	aaagtgagac	cctgtcaaaa	aaaaagagaa	ggtgatgggg	aaagaacaca	1620
gaacagcata	agagggggtt	ggggaagctg	ggtggagtgg	gggggattgc	agttgaaagt	1680
	gggaaggcct					
	ctcattcttg					
acaggaggga	atgtggtgga	ggagagtgag	cagcaggagg	gagcagtgaa	ggtcagcaag	1860
gtgacagagt	ggctgaatca	aaaaagacct	tgcagtgttt	gagcagagga	tccatatcat	1920
ccattatgtt	ccaaaggact	cttcaggatg	ccgtgtggag	aaaggaagag	ggtggaagcc	1980
aggaggtctg	gagggaggtc	tggagtggag	gagatgagag	gctccggatc	cctctgggag	2040
gtagatttga	ggacagattg	gaattgaggt	gaaagacaga	gaaagagaag	tggccaggat	2100
gactccaaga	tttctgacct	aaactactgg	gaaggacgcg	gttgtcattt	ctgaaatgca	2160
gaaggatgcc	agaagagaag	gtactttggg	gaggggggg	aatcaggagt	tagttttgga	2220
catgagataa	gcttggaata	tttatttgct	atctaagaca.	gctccttaac	atggtaagcc	2280
cttatgcaag	ttgttgtcag	ctgagatggg	cgtggcactg	agcatgggag	catggaggcg	2340
cctgagtggt	ctcatgctca	ggtggtttag	caaactcagt	gtacatcctg	ccaattccag	2400
tcctgccatg	gccactgaca	agctaggagg	gcgctgaaag	gagaaggacc	ccgatgtctc	2460
	tccatctcct					
gggtacccgg	tgatgaagac	tgtggtatca	gcctcctgag	cacagagagg	gcagaaaggg	2580
gtggagacaa	agaggggcgc	agatagtggg	caaatgggga	agtggcactt	cccctagctc	2640
	gcttggtgtg					
	agaacaagtt					
	ccccgcacc	•				
	tccagcagcc					
	tgtgcttttg					
	ttcctgggcc					
	cccacctgcg					
	accaattgca					
	gcctgcgctg	•				
	gctgggtcca					
	tgġtgttaga					
	tggtgggagt					
	agttctcaca					
	aagggctttg					
	gaaatggtgt					
	aggtctcact					
	teegeeteee					
	aaattttaag					
	gatcacaggt					
	ggcaaataac					
cagaccacac	tttgagaacc	actggactag	accettgatt	gaggagtacg	gtgttgagag	3900

			•			
tggagtcctc	tgtgatggtg	gatggaccag	gacacatggc	ataggagtca	ggtggttccc	3960
					ataatcacgt	
					atgcagcgtt	
					tttgtttcca	
					ccactgcttg	
					agaaactgca	
					gagacagagt	
			•		gcaacctctg	
					ttaatggtgc	
					catcatgttg	
					tcccaaagtg	
					ttgaaataat	
					aaataatgtg	
					tgggagggtg	
					ttgggtgacg	
					cagatetgag	
					gctgtaatcc	
					agaccggcct	
					gtgtagtggc	
					gagcccagga	
					gacagtgaga	
					cctaaattta	
					gtccaaaata	
					cgtgagatgt	
					gtgcgatgca	
					gggctttcag	
					gcatccttgg	
					gccccagcac	
					aagctggaag	
					gccagcctac	
					ccctgagaac	
					tgcggataac	
					caagcccágc	
					acagctgcct	
					ggaggaagcc	
					gtcctcccac	
					ctgtttccac	
					ggaaggcatg	
cctgctgctg	attgaaaacc	gaactgggaa	cattccttcc	attctgtgtc	cactggtcag	6240
ctgctgcggc	tttggatggt	cttgaccgtg	gaaggctgac	cttcttctgg	tacccggagt	6300
ccctgcagga	atcccccttg	agcttgctgg	gctgtggtga	caggagttta	aaacatgcgt	6360
					aggtctcatg	
					aaagaaaaat	
					tgcagctgta	
					actttgggag	
					aacatggtaa	
					cgcctgtaat	
					ggaggttgca	
-					-	

gtgagcttag	atcatoctac	taccetecaa	cctgggcgac	agagtgagat	tacotctcaa	6840
			tttttaaaag			
			ggtggttagt			
			cagcgcaggc			
			cactgcactc			
			gtaattaata			
			ttctttgcta			
			gaccaaaaat			
	i i		agtcactgca			
			ggaaacctga			
			tatcccctg			
			aacgacccgg			
			gggtgggtga			
			ggctggatag			
			aagtcccttt			
			tgccgctcct			
			gacacaagtg			
			gaataatact			
			aaggaggaag			
			ggccacaccc			
	•		atggaagctg			
			gttttgagac			
			cactgcaacc			
			gggactacag			
			tttcactgcg			
			gcctcccaaa			
			cttaagtaca			
			tcagtaatgg			
			tgcagggagg			
			tcccattgga			
			aatgcctgct			
· ·			gcctcagttt			
			cagtacaagg			
			catgtgccct			
			cacactggtc			
			gcccaggctg			
			ctggtgtctg			
			tcaggtctgg			
			gttggctcac			
			gctagagagt			
			tgcaagacct			
			gagtccaaag			
			ggatccagct			
			tttttaggca			
			acatéccaaa			
			aaggctgctg		a contract of the contract of	
			tgtgggtgga			
			tgcctctagg			
3 3 -		-				

catccaaacc agatgaactg aaaataaggc aggagtggct tccccagggg aaactgggga 9720 . agaggaagca ggactgtgct ggctaaaatg ccagccaggt ttaagacgtg gcaccagatg 9780 ccagtcatgg gattggattg gtcagcatgc ctgggctatg gcttaggggt atgtttggtgc 9840 tcagggatgc cacaggcctc cagataccag gtctgaggca gaagaatgaa gtccagcttc 9900 tettgtgggt ggaacagtgg caactgagat accccatete teeetteeca agaacagage 9960 tgaacataaa gaatttagtg attggccaga gettggccac atgeteeeet etgatgaatg 10020 ataggccagg tgatgggatt ggcacaattg gcttagacta atgagggttg gccctggagt 10080 tgcaggcagt ggagttctgt cctaagcagt gggcacctaa acccgatggc ataaaagctg 10140 ggcgggtgtc cacctgcatc tgccacagca ctataggcac caactgtggc tcatactgag 10200 tgggataaat tccagaaaga aacattagga acttactata gaattttggg gctagagcta 10260 ctcattcatt cccctagata atttctaggc aaggttccat agtggagggg gagttttggc 10320 ttgggcattg aaggatgcat aggagttttc tagatgggga aagaagggaa cggtagacca 10380 ggcagaggga actgcatgat aaaaggttta tgggtgtgaa aattcatgga atgtttgagg 10440 attatggggt tgggggatgt gggaatatgt gtagcgataa agcaccaaac aaagccaaaa 10500 gtttagttag agccctgaat gcctgcctca taatggtttc catattttat atgcctacta 10560 tgtgccaggc acattgctca gggtcacaca gctggaaatg gcagggctga gtttttgttg 10620 ttgttgttgt tgttgagaca gagteteact etateaceca ggetggaatg caggggegtg 10680 atcatggctc actgcatcct tgacttcctg ggatcaggtg attctcccac ctctgcctcc 10740 caggtagctg ggactacagg cacaggccac cacgccaggc taattttttg tatttttagt 10800 agcgacaggg tetegecatg ttgteeggge tggtetggat eteetggett eaagtgatee 10860 ccctggctca gcctcccaag gtgctgggat tacaggettg agccaccgca tccagcccag 10920 atctgagatt tgcacccagt atttgaactc ccaagcctgt gctctttttc ctcccatgga 10980 catttctctc agagatggtc tcccaaacac ctgtccttct tgttaaaaaa cagacaaacc 11040 gcaagtagtt ctttggaagc teagatttet ettttgttte ttagtaaaac attteecagt 11100 teccagetee ettecagggt gtaagattte tteggtaact tacatetage tgttgettet 11160 tgtttgctca tgtttagaaa gaaagacaaa agagagtgag aattttctct cccttcccca 11220 gtetececae aacteacaee ecaceeteag etecetetgt aataggaaaa tetetgaact 11280 ctctgtagtt gctccagcaa tcttttggaa ctttgcttct ttcttgtgaa aaaacctccc 11340 cttggctcac tttgcaccag gtttccccaa atgtgcttcc aaccacaagc agaaatggag 11400 ctgccagtaa ccaggaagaa actgccgggg gctgaggaag aggagaggga ggtgcatagc 11460 cctggatctc gcagggagag gggtgacagg atgagaactc aggttgctca cttgccatca 11520 gggtcagtca tgaatatagc gttcatgtat cactttttaa agcttttttg gagggtaaaa 11580 gtaatagtta cacaaaataa aaatacaaat ggtacaaaag gacttagaat ggaaacatgt 11640 tteteteceg actecageet cetgttttte tteccagaga etgaccaetg etgtetgtet 11700 cttgccagaa gggaaaggga ggcaaggtta gggcaggcag agggcatgtg catcctttag 11760 agagagetta tgtetataca ageaaatgtg tgtgtteagt categetgte ttagttttet 11820 attgctgcat aataatggta ctaccagett cacagettta aacaacacce atttattate 11880 tcatagtttc tgtggttggg agtctggaca tagcttagcc aggttctctg ctttagagtc 11940 tcgtgaggct ataatcaagg tgtgggatgg ggctgcagtt tcatctgagg ctcaattggg 12000 gaagggtcac ttctaagctc atacaatatt ggtgacattc agtccctggc aggctgttga 12060 actgagagcc tcagtttcgt gctggctgtt ggttgtagtt aaccctgaat tccttcccat 12120 gtgccctttg caaagccatc aaggcagaga gacttgccta gcaagtagga tattacagtc 12180 ttctgtaata taatcacatc catgaaatcc tctatatatc ccatcacctt taccatattc 12240 tgtgggttag aaacaagtag caggtcctgc ccacactcga gaagaccaga tgacacaaag 12300 atgtgattca aagtggggat catcggggcc atcttaggtt tgtctgcagt gatcactgtg 12360 ccatctctct ctctctcttt ttttttttt tttttccgag acgaagtcgt cactctgtca 12420 cccaggctgg agtgcagtgg catgatetea gettaccaca atetetgeet eccaggttea 12480 aatgattett etgeeteage eteetgagta getgggatta eaggtgeeeg eeaceacace 12540

cagctaattt ttgtattttt agtagagaca gagtttcacc atgttggcca ggctggtctt 12600 gaactcctca cctcaagtga tccacccact tcggcctccc aaagtgctgg gattacaggc 12660 atgagecace atgeecagee ceatetetet ttaaaaaaaca aacaaacaaa caaaaaacat 12720 aaaaagaagc agagaacaca tacacatctg catcttccct tgtttactta acaatagatc 12780 ttggaagtca cttctcagta gaggctaggt tgggcagagc attggattct aggccagtga 12840 gtttggactt gaccatggag acactaggaa gcccatgaag gacagagaga gatgcctcga 12900 ccctgccagt cctttagaaa gatcacccag tgctttttgt ataccaaacc ctatttgaaa 12960 tacttacgta tattaaccca tttccttatc accacaaccc tgcgggaagg gagataggca 13020 cttttattat cttcattttg cagatgagga cattgaggtc cagagaggtt atgtcactta 13080 cttaaggtca cacagccagg aagtggtagt agggactctt accettgttt tacagatgag 13140 attgaattat ctcacgaaaa ctcagaaagg ttaaacaact tgcctaagta acatacagct 13200 aattagtcga ggagcctgac gcatgttgct ctagcctggt cacagttaca gaggtggcaa 13260 gcaatggcct gaacaggacg aacaaccaaa tacccaggct ggtggctctt aaacatggtg 13320 gggtcagcta acgacagcaa ccagggtggg cactggtgcc cctcgccccc ggctggtgcc 13380 ctaacatete cettttetet accagtteag aatetataac gtgacetace tagaaccete 13440 cctccgcatc gcagccagca ccctgaagtc tgggatttcc tacagggcac gggtgagggc 13500 ctgggctcag tgctataaca ccacctggag tgagtggagc cccagcacca agtggcacaa 13560 ctgtgagtat caagaggeet aageaatggt aateteeact etecattett ceeetgtgge 13620 cagacacttc ccctggctga gtctctgggc ttttatatca taggatgcct ctaatggcaa 13680 tcctqccatt agatacacct gctgtggtgt atctgccagg taggcaggct aggctgcagt 13740 aacacacaag cccacaattt ccatggctta acactatagg aatatatttc ttgctcatgt 13800 aacaaqctaa cqtqaatqtt qctqqtttqt aggtqqtttc cctccctqta gaaatctqqq 13860 gagtgaggtt etttecatet tgtggtgeea teatteteea ggaeaaagat tettacetae 13920 ttttgtgtcc tggtttcctt tggcagcctg gtgaagccta tggacctcat ttcagaatat 13980 ttttaaatac ataaaatccc agcctgggca atatagtgaa acccccatct gtacaaaaat 14040 tagccaggca tggtggcatg cacetgtagt cecaggtact gggaaggetg aggtgggagg 14100 atcacttgag cccaggagtt tgaggctgca gtgagccgtg atcgtaccac tttactccca 14160 aataaataga actacagagg aaactaattg tattgaaatg cagttataaa acatttaaac 14280 acatttttaa tetagagata tatgtgette tttattaaga tetataaata ataagtteta 14340 ggggtagete geataaatae tgtaatttea aagtagataa geataaataa taetttatga 14400 tactgaaatt gtgatgtgat atgagaatag ctgtgagttt tgttttgctg gggacaggat 14460 cactgatgct gtcattactg gggtctcttc cctccattct ttttttaaaa ttgtatttta 14520 ttttattttt aaaattttaa aataaataga gacagggtat cactatgttg cccaggctgc 14580 ttttgacctc ctgggctcca gtgatcttcc catcttggct tcccaaagtg ctgggattac 14640 aaqtgggage cagtgtteet ggeeeettee teeattetta atggaaggag atgetaggtg 14700 tgagaggtta gggaaagtaa agatgtaatt tettteeeat eeaagttete agaceeetga 14760 attotacctg cagocatgtt ggtocatcaa coccaagtga agaatccctg ctctagggcc 14820 ccaccattgt ctgtatccag ccagcagaag aggcgtgatt atggagatca catctgcttc 14880 ttgaaagcag acagcccgga agtgggccgc atcacttcct ctcaaattct attggtgaaa 14940 atggtcacat gactacacat agccacaaag gaggctggga actttctcac ttggaaccta 15000 catcccagaa acaactettt teagtgaggt ateccacagg tetttegeag tagaaatatt 15060 gattatetea cataaaatga agtettacaa atggacetae tgggttttgt acageageca 15120 agtgatatet ettecettet getgtettee ettetgeegt cetteacatg gtggcattgt 15180 atccttagac ttgccaccca tgccctcagg ttggccgttg cacactgtct tacataaagc 15240 aggaaggaaa ggaaaggctg ctacgagaga gtgtaccttg tgcatctctt tittaatcag 15300 gaagcaaaca tetttetaga agetteeeta geaaaattee eettacatet cattggeeaa 15360 gactgttaca tgttacatgg ttactgttat tacttgctca ttgcaaggaa gactgggaac 15420

tcaaatgcct	ggaaaaagga	acaggataat	cgtgattggc	tcaagcctta	gggtgggcat	15480
					ttcagtttcc	
					aagattcttc	
					cagccctctc	
					ccctgggctg	
					gaaccaggtg	
					aactagcttg	
					atttgctgag	
					acaaagcaga .	
					gcaaacatat	
					aaaataaaag	
					cggggaaaac	
					gagatgatgg	
					ttcatcatgt	
					cagatgagga	
					agccatccgg	
					aagtttaaga	
					gtgtgacgag	
					tcggcggtca	
					ggaagtggag	
					cagcctacag	
ggagcccttc	gagcagcacc	tcctgctggg	cgtcagcgtt	tcctgcattg	tcatcctggc	16740
cgtctgcctg	ttgtgctatg	tcagcatcac	caagtgagtc	ctgggcccag	tgctgccgag	16800
cagtccctct	ggagtgcagg	gţggcaggga	cttgcccctc	tagtctgccc	ctttgcagtc	16860
ctctcagtca	ataatacgta	tttactgagc	agctactaca	caccttgaga	gtagagctga	16920
gaacatatcg	acaaggaccc	cactttttc	ttttttttt	tttttttt	ttttgagacg	16980
gagtctcact	ctgtcaccca	ggctggagta	tagtggcaca	atcttgccta	acagtaacct	17040
ccgcctcccg	ggttcaagca	attcttctgc	ctcagcctcc	agagtagctg	ggattacagg	17100
cgcatgccac	tatgcccggc	taattttttg	tatttttggt	agagatgggg	tttcaccatg	17160
ttggtcaggc	tggtctcgaa	ctcctgacct	catgatctgc	ctgcctcagc	ctcccaaagt	17220
gctgggatta	caggtgtgag	ccactgcacc	caaccaggac	tccacatttc	taaaaccggc	17280
atcctacțgg	ggagactgaa	aatacatatc	aatcacaaac	aggtggtttt	ccatagtgac	17340
ccactctctg	aatgcactag	accagggtgg	aggccagaga	tcttctgggg	tgctttttgc	17400
aagggggacc	aggataaggc	tctccaagga	gggaaaattt	gagggggcc	ctgactgggg	17460
agaatgagct	ggccagggat	aagcaagatg	gagtcatccc	acateceett	acaacactgg	17520
gtgcctgggc	aactgggggc	atttgggggc	atgtggtagg	agccagagga	atttgcgacg	17580
attgccctga	tggagtcagg	agacctgggt	ttgaatcctg	gccttggagc	ttggtagctg	17640
gcggccgaca	agttgctgaa	acccctgagc	ctggggttcc	tgctttgcag	agtgacagtg	17700
atggtgagaa	catatttcat	cagccagaag	aggccaaatc	acagtaaagg	ctgagggagg	17760
agatgagtgg	cgagtggctg	ggaggṭggtg	gaaggagcct	cgtttccaga	gagetettge	17820
cagcccttgg	aatcatggtg	tctcagagcc	tcagtcctcc	catctctgaa	atgggactag	17880
caagctcaac	ctcactaagt	caggattaga	ggtggctaag	gattattaac	atgattgatg	17940
aaagtgccca	ctcttggccc	agcacacact	aggtaggcag	ggaatgcaaa	ttcccctcca	18000
tatcttgtca	ctgatgcctc	cgagcaacct	tggactgatc	gccttgctct	gagcctcagt	18060
ttccccatca	cctgtacctc	ttcccactcc	ccatcactat	atcccagcat	gccagcctct	18120
ttgctgttct	ttgtctttgg	tttcttgttt	tgttctgttt	tttagacagg	gtctcactct	18180
gttagccagg	ctgaagtgca	gtggcgcggt	tacggctcac	tgcagcctcc	aattcctggg	18240
ctaaagagat	cctcccattt	caacttccag	agcagctggg	acaacaggcg	cttgccacca	18300

18420 18480 18540 18600 18660
18480 18540 18600 18660
18540 18600 18660
18600 18660
18660
18720
18780
18840
18900
18960
19020
19080
19140
.19200
19260
19320
19380
19440
19500
19560
19620
19680
19740
19800
19860
19920
19980
20040
20100
20160
20220
20280
20340
20400
20460
20520
20580
20640
20700
20700 20760
20700 20760 20820
20700 20760 20820 20880
20700 20760 20820 20880 20940
20700 20760 20820 20880 20940 21000
20700 20760 20820 20880 20940 21000 21060
20700 20760 20820 20880 20940 21000

ggctgacttt tacctgggac agtgcaggtg cctgagccat gtgcctctca ctctccagca 21240 ggccggccca ggcttgttta cagagtggct cagttttcaa gggtgggaag tcccaaggct 21300 tcttgaggcc taggcgcagc actggcatga tatcacttcc atcacattct atgggcccaa 21360 gcaagtccca gggccagtgt agattcaagg gatgggagga gattcagagc actcctctgt 21420 ggccactttt gccatcgacc acagtccctg taaatattag gacaatgtaa ttaattccca 21480 ggaatctgag gctcagaaag cgtaagtgac ctgttggact tctgatctgt gtgatgtcga 21540 qqcttqtacc ccttcctgag cattgccgta ctccaggccg ggctgcaagg ccactctgct 21600 ctttcattgg ctgtctctgt attttagggg tcacagtggg agaagcggtc ccgaggccag 21660 gaaccageca agtgecegta tgtatetgaa ettaggteae ageetgeatg cattgggaag 21720 gtgatagaat tggagaggca agcccctagc tccatgtctg ccttctcttc cctgcattcg 21780 gtaattgccc tgtgacatta gccttcaagg gacggcagga ggaggggtgt tctggaaacg 21840 tggactgctg gccaagcccc ctgagtttca ctggtgtgtc aggtacatgg tgatacccct 21900 tgggagtgct gttatagtta acaaccagag cagccgtgcc tgttgttaaa atcttgacct 21960 aattgtatac ttgtcggcaa atagccacta tcctgaacac tcccctcctt ttttttaata 22020 tacaggatet cactetgtgg eccaggetgg tgtgcagtgg tgcgateata geteaetgea 22080 ccttcaaact cctgagetca agtgatecte ccatettage etecegagta getgatacta 22140 cagatgtgca ttaccacgcc tggctatttt aaaaggtttt tgcctgtaat tccagctact 22200 caggaggctg aggcatgaga atcacttgaa cccgggaggc agaggttgca gtgagcgcag 22260 attgtgccac tgcactccag cctgggcgac agagtgagac tcttgtctca aaaaaaataa 22320 taccaaaaaa agtttttgta aagacaagct ctcgctgtgt tgccccgcca ctgtggcctc 22380 cttagcttct tccctggggc ctgctggacc tttccatact ccagaaacta aagggggtcc 22440 aggaccetge tteaacceta ggatecegea tettttttt ttttttttt tttttggaege 22500 agggtcttgc tgtgtccctc aggctggagt gcagtgattc actgcagcct caaactcgtg 22560 ggctcaagtg attctctagc ctcagccttc taagtagctg ggactacagt catacaccaa 22620 catgcccage taattttcct tttttttaat tcttgtagag atgtttgaga eggcttgggc 22680 tetgttgeec aggetgttet caaacteetg ageteaageg atecteecte etcageetee 22740 taaagtgctg ggattacagg cgtgagccac cgcacccggc ttccatatcc tttctaattg 22800 gtcatggctt gggataatgg tgttgctttt aattatcatc atccataaag actttttctt 22860 actcaacaga tetgagettg tatttggtge ecaggacatg tgetgggtte ecgaaateee 22920 aaagacacag accetaceet cagggattte teattetage aacatagaet gateaattae 22980 tgattataac gttagaaggc atgtctgaag tagacagcca tcaggacatg gtgatttcag 23040 gctgggcttt gaagaatgaa taggagtttt tcaagtgtcg aaactgaacc ctgaccaacc 23100 tttgcttttg cagacactgg aagaattgtc ttaccaagct cttgccctgt tttctggagc 23160 acaacatgaa aagggatgaa gatcctcaca aggctgccaa agagatgcct ttccagggct 23220 ctggaaaatc agcatggtgc ccagtggaga tcagcaagac agtcctctgg ccagagagca 23280 tcagcgtggt gcgatgtgtg gagttgtttg aggccccggt ggagtgtgag gaggaggagg 23340 aggtagagga agaaaaaggg agcttctgtg catcgcctga gagcagcagg gatgacttcc 23400 aggagggaag ggagggcatt gtggcccggc taacagagag cctgttcctg gacctgctcg 23460 gagaggagaa tgggggcttt tgccagcagg acatggggga gtcatgcctt cttccacctt 23520 cgggaagtac gagtgctcac atgccctggg atgagttccc aagtgcaggg cccaaggagg 23580 cacctccctg gggcaaggag cagcctctcc acctggagcc aagtcctcct gccagcccga 23640 cccagagtcc agacaacctg acttgcacag agacgcccct cgtcatcgca ggcaaccctg 23700 cttaccgcag cttcagcaac tccctgagcc agtcaccgtg tcccagagag ctgggtccag 23760 acceaetget ggccagacae etggaggaag tagaaccega gatgeeetgt gteeeccage 23820 tetetgagee aaceaetgtg ecceaacetg agecagaaac etgggageag ateeteegee 23880 quaatgteet ecageatggg geagetgeag ecceegtete ggeececaee agtggetate 23940 aggagtttgt acatgcggtg gagcagggtg gcacccaggc cagtgcggtg gtgggcttgg 24000 qtcccccagg agaggctggt tacaaggcct tctcaagcct gcttgccagc agtgctgtgt 24060

ccccagagaa atgtgggttt ggggctagca gtggggaaga ggggtataag cctttccaag 24120 accteattce tggctgeect ggggaccetg ceecagteec tgteecettg tteaectttg 24180 gactggacag ggagccacct cgcagtccgc agagctcaca tctcccaagc agctccccag 24240 agcacctggg tctggagccg ggggaaaagg tagaggacat gccaaagccc ccacttcccc 24300 aggagcaggc cacagaccc cttgtggaca gcctgggcag tggcattgtc tactcagccc 24360 ttacctgcca cctgtgcggc cacctgaaac agtgtcatgg ccaggaggat ggtggccaga 24420 cccctgtcat ggccagtcct tgctgtggct gctgctgtgg agacaggtcc tcgcccccta 24480 caaccccct gagggcccca gaccctctc caggtggggt tccactggag gccagtctgt 24540 gtccggcctc cctggcaccc tcgggcatct cagagaagag taaatcctca tcatccttcc 24600 atcctgcccc tggcaatgct cagagctcaa gccagacccc caaaatcgtg aactttgtct 24660 ccgtgggacc cacatacatg agggtctctt aggtgcatgt cctcttgttg ctgagtctgc 24720 agatgaggac tagggettat ccatgcctgg gaaatgccac ctcctggaag gcagccaggc 24780 tggcagattt ccaaaagact tgaagaacca tggtatgaag gtgattggcc ccactgacgt 24840 tggcctaaca ctgggctgca gagactggac cccgcccagc attgggctgg gctcgccaca 24900 teccatgaga gtagagggca etgggtegee gtgceccaeg gcaggeccet gcaggaaaac 24960 tgaggccctt gggcacctcg acttgtgaac gagttgttgg ctgctccctc cacagcttct 25020 gcagcagact gtccctgttg taactgccca aggcatgttt tgcccaccag atcatggccc 25080 acatggaggc ccacctgcct ctgtctcact gaactagaag ccgagcctag aaactaacac 25140 agccatcaag ggaatgactt gggcggcctt gggaaatcga tgagàaattg aacttcaggg 25200 agggtggtca ttgcctagag gtgctcattc atttaacaga gcttccttag gttgatgctg 25260 gaggcagaat cccggctgtc aaggggtgtt cagttaaggg gagcaacaga ggacatgaaa 25320. aattgctgtg actaaagcag ggacaatttg ctgccaaaca cccatgccca gctgtatggc 25380 tgggggctcc tcgtatgcat ggaaccccca gaataaatat gctcagccac cctgtgggcc 25440 gggcaatcca gacagcaggc ataaggcacc agttaccctg catgttggcc cagacctcag 25500 qtqctaqqqa aggcqqqaac cttgggttga gtaatgctcg tctgtgtgt ttagtttcat 25560 cacctgttat ctgtgtttgc tgaggagagt ggaacagaag gggtggagtt ttgtataaat 25620 aaagtttett tgtetettta ttttttatgt attaaccaaa cataceteea gacaetgetg 25680 tgagtgctgt gtctctgtta actcctggaa ttcacccatc cagaggaacc aggatgcaag 25740 aggttaagaa acttgccgtc tgggtttggg ttccccatac aaggattcaa atagttgatt 25800 taggaagtaa tcccgggaaa ccctgctaag gtagtgggga actgaggcag ggaaggacac 25860 aaaccaagaa agtgttacct gaaaggggtc cagatgcaga ccccaaaaga gggttcttga 25920 ateteatgea agaaagaatt cagagegagt ecatagagte agtgaaagea agttaatgag 25980 gaaagtaaag gaataaaaga atggctactc cgtagacaga gcagccctga gggttgctgg 26040 ctgcctattt ttatggttat tgattaatta tattccaaac aaggggtgga ttattatgcc 26100 tcccttttag accatatagg gtaacttcct gatgttgcca tggcatttgt aaactgtcat 26160 ggcqctqttq ggagtqtagc agtgaggaca accagaggtc actcttgttg ccatcttggt 26220 tttggtgggt tagagccatc ttctttactg caacctgttt tatcagcaag gtctttatga 26280 cttqtatcgg tgacgacctc ctgtctcatt ctatgactaa gaatgcccta acctcccagg 26340 aatgcagccc agtaagtctc agcctcattt tacccagccc ctcttcaaag.ctccagttta 26400 aataaacctc tgacaaaagg gtgagttatt caacagatta ccagcatgag taactgatgc 26460 ttacctgccg gggatctctg gaagaccatg catggcacat gcccagttat gcctgcaaag 26520 gagagggagc tggggtattt gtccaccagc tcccatctgt cattggctga gagetgcttc 26580 caggageatt aatteteeag caetteeage tacteeagga aaaaaaaaat tetteaactg 26640 agagttggag gtgttgagag actctggcac accaagaaga caggaacagg acaccaacag 26700 tggctgatga tacactgcca aggtcacaca gctagttagc aacagatcta tagtggaatc 26760 cagacagtgt etecateace caggetetet gtagtgatet gegetteaca teegaggeag 26820 gcagagggat ggtgtgggcc ttagatggga aggctgggaa cctgaagctc ctatgtctgt 26880 atcacttttg cttctctgag tagctgccct gatttcacac ttgaggggct tggccatttt 26940

```
agattectic etgetetagg ageetacata etacaetgga aatgatgggg agetetetae 27000
ctcacatgca gcctgatgtt tgttagaaac acctccttgc gccaggcatg atggctcatg 27060
qctgtaatcc cagcaatttg ggaggctgag gcgggtgtat cacttgaggt gaggagttca 27120
agaccageet ggecaatatg gtgaaaceet atetetaeea aaaaataaaa aattageegg 27180
gtgtggtggt gggtgcctgt aatctcagct acttgggagg ctgagttggt agaattgctt 27240
caacctggga cgcggaggtt gcagtgaget gagattgtge cattgcactc cagcttggat 27300
gacagagtga gaccetgtet caggaaaaaa aaaaaaaaac aaaaaaaace ttgttetaag 27360
ccaaaatcaa tccctttagc tgcccaaatc acacagttta cagatggaga aacagtttta 27420
gagaggaaaa gggacttgcc caaagtcacc cagagaatgg cagagcctga actagccttc 27480
tggacttctt gcctccaaaa gctctttata ataaaatata attttaaata aaaatagtta 27540
tctgtttagg gccaagcaat atgctaagtg ccgtccagcc actgtgtcat ttacgtctcc 27600
aaacagctct agttgggagg ctcaatgatt atcccaattt tacagataag gaaacaggtc 27660
cagagaggtt gaggattagc ctagaaccac acagctagga aatcctggag ccaggatttg 27720
aacccgggtc tgacctaaga gctcccagcc gccgtgatat atcagcttat gtcatcctga 27780
cacctacgca gatgtegget egaatecact ttgcetgage attgtetcag agaaatetaa 27840
tttaaaaatt aggcagcaaa tagaaaatat atttgactgc tagagatgca atgggactgg 27900
gageceaaca aaggatetta ggeaaaagaa atecaagttg ttggeeteag caactattae 27960
tgaactggct gggctttggg aagctacaga gggatgagaa gacctggtgg atcaggtggg 28020
cccaactcag gctggccccc accctgcagg aagtaggaaa agtccagggt cataggccca 28080
gtgagatgcc ggctgcggga gtttcagcct ccggggctgg accagagggc aggaggggac 28140
gcccctgggt agcagegcca gagtgggctg agtggcctgg gcccctgcgg gggagctttc 28200
agagatgttg atttgggggt actccctcag ccctgccttt acacagaatt tgtgggggat 28260
gaggggaggg ggaaaggggg gaggaaggca gtgagtgcat ctgaattttt ttttttttt 28320
tacaaaaagt ggcttattgc atttttctga ttactctatc agcacgtgca gaccttttcc 28380
tattcagaga aagcctgaag atataaagag gaaagtgaag aaaaaccacc ggaaatccca 28440
teceegeece ageatetgge actgtgtggg egateaegaa atgagegett gtttttgaag 28500
gcgtagtatc tccgtgaaca tccggttgaa caacctttct gactttattt ttcccacgaa 28560
agttattaat taaaaaacaa aaagcaaaac accgaaaaaa caaaaaaccc agcaagtgtt 28620
tgagetecca ceaegaggga ggeetgaegt caetggatee teeeggeage egatgagget 28680
                                                                  28690
gcatgggact
```

```
<210> 2
<211> 2476
<212> DNA
<213> Homo sapiens
```

### <400> 2

```
atggggtggc tttgctctgg getectgtte eetgtgaget geetggteet getgeaggtg 60 geaagetetg ggaacatgaa ggtettgeag gageeeacet gegteteega etacatgage 120 atetetaett gegggtggaa gatgaatggt eeeaceaatt geageaeega geteegeetg 180 ttgtaceage tggttttet geteteegaa geeeaeaegt gtateeetga gaacaaegga 240 ggggggggt gegtggea eetgeteatg gatgaegtgg teagtgegga taaetataea 300 etggaeetgt gggetgggea geagetgetg tggaaggget eetteaagee eagegageat 360 gtgaaaeeea ggeaeeegga aaacetgaa gtteaeaea atgteteega eactetgetg 420 etgaeetgga geaeeegga teeeeegga aattaeetgt ataateatet eacetatgea 480 gaaceetee teegeatege ageeageae etgaagtetg ggatteeta eagggeaeeg 600
```

PCT/US00/19094 WO 01/04270

```
gtgagggcct gggctcagtg ctataacacc acctggagtg agtggagccc cagcaccaag 660
tggcacaact cetacaggga gecettegag cageacetee tgetgggegt cagegtttee 720
tgcattgtca teetggeegt etgeetgttg tgetatgtca gcatcaccaa gattaagaaa 780
gaatggtggg atcagattcc caacccagcc cgcagccgcc tcgtggctat aataatccag 840
gatgctcagg ggtcacagtg ggagaagcgg tcccgaggcc aggaaccagc caagtgccca 900
cactggaaga attgtcttac caagctcttg ccctgttttc tggagcacaa catgaaaagg 960
gatgaagate etcacaagge tgecaaagag atgeetttee agggetetgg aaaateagea 1020
tggtgcccag tggagatcag caagacagtc ctctggccag agagcatcag cgtggtgcgá 1080
tqtqtqqaqt tqtttqaqqc cccggtggag tgtgaggagg aggaggaggt agaggaagaa 1140
aaagggaget tetgtgeate geetgagage ageagggatg acttecagga gggaagggag 1200
ggcattgtgg cccggctaac agagagcctg ttcctggacc tgctcggaga ggagaatggg 1260
ggcttttgcc agcaggacat gggggagtca tgccttcttc caccttcggg aagtacgagt 1320
gctcacatgc cctgggatga gttcccaagt gcagggccca aggaggcacc tccctggggc 1380
aaggagcagc ctctccacct ggagccaagt cctcctgcca gcccgaccca gagtccagac 1440
aacctgactt gcacagagac gcccctcgtc atcgcaggca accctgctta ccgcagcttc 1500
agcaactccc tgagccagtc accgtgtccc agagagctgg gtccagaccc actgctggcc 1560
agacacetgg aggaagtaga accegagatg ceetgtgtee eccagetete tgagecaace 1620
actytycece aacetyagee agaaacetyg gagcagatee teegeegaaa tyteeteeag 1680
catggggcag ctgcagccc cgtctcggcc cccaccagtg gctatcagga gtttgtacat 1740
qcqqtqqaqc aqqqtqqcac ccaqqccagt qcqqtqqtqq qcttqqqtcc cccaqqaqaq 1800
gctggttaca aggccttctc aagcctgctt gccagcagtg ctgtgtcccc agagaaatgt 1860
gggtttgggg ctagcagtgg ggaagagggg tataagcctt tccaagacct cattcctggc 1920
tgccctgggg accctgccc agtccctgtc cccttgttca cctttggact ggacagggag 1980
ccacetegea gteegeagag eteacatete ecaageaget ecceagagea eetgggtetg 2040
gagccggggg aaaaggtaga ggacatgcca aagcccccac ttccccagga gcaggccaca 2100
gaccccttg tggacagcct gggcagtggc attgtctact cagcccttac ctgccacctg 2160.
tgcggccacc tgaaacagtg tcatggccag gaggatggtg gccagacccc tgtcatggcc 2220
agtecttqct qtqqctqctq ctqtqqaqac aggtectcqc cccctacaac ccccctqaqq 2280
gccccagacc cctctccagg tggggttcca ctggaggcca gtctgtgtcc ggcctccctg 2340
gcaccetegg gcateteaga gaagagtaaa teeteateat eetteeatee tgeecetgge 2400
aatgeteaga geteaageea gaceeecaaa ategtgaaet ttgteteegt gggaceeaca 2460
                                                                  2476
tacatgaggg tctctt
```

```
<210> 3
 <211> 825
<212> PRT
 <213> Homo sapiens
 <400> 3
Met Gly Trp Leu Cys Ser Gly Leu Leu Phe Pro Val Ser Cys Leu Val
   1
                   5
```

Leu Leu Glin Val Ala Ser Ser Gly Asn Met Lys Val Leu Glin Glu Pro 25 20 .

10

Thr Cys Val Ser Asp Tyr Met Ser Ile Ser Thr Cys Glu Trp Lys Met 45 35 40

Asn	Gly 50	Pro	Thr	Asn	Cys	Ser 55	Thr	Glu	Leu	Arg <sup>.</sup>	Leu 60	Leu	Tyr	Gln	Leu
Val 65	Phe	Leu	Leu	Ser	Glu 70	Ala	His	Thr	Cys	Ile 75	Pro	Glu	Asn	Asn	Gly 80
Gly	Ala	Gly	Cys	Val 85	Cys	His	Leu	Leu	Met 90	Asp	Asp	Val	Val <sup>·</sup>	Ser 95	Ala
Asp	Asn	Туг	Thr 100	Leu	Asp	Leu	Trp	Ala 105	Gly	Gln	Gln	Leu	Leu 110	Trp	Lys
Gly	Ser	Phe 115	Lys	Pro	Ser	Glu	His 120	Val	Lys	Pro	Arg	Ala 125	Pro	Gly	Asn
Leu	Thr 130	Val	His	Thr	Asn	Val 135	Ser	Asp	Thr	Leu	Leu 140	Leu	Thr	Trp	Ser
Asn 145	Pro	Tyr	Pro	Pro	Asp 150	Asn	Tyr	Leu	Tyr	Asn 155	His	Leu	Thr	Tyr	Ala 160
Val	Asn	Ile	Trp	Ser 165	Glu	Asn	Asp	Pro	Ala 170	Asp	Phe	Arg	Ile	Туr 175	Asn
.Val	Thr	Tyr	Leu 180	Glu	Pro	Ser	Leu	Arg 185	Ile	Ala	Ala	Ser	Thr 190	Leù	Lys
Ser	Gly	Ile 195	Ser	Tyr	Arg	Ala	Arg 200	Val	Arg	Ala	Trp	Ala 205	Gln	Cys	Tyr
Asn	Thr 210	Thr	Trp	Ser	Glu	Trp 215	Ser	Pro	Ser	Thr	Lys 220	Trp	His	Asn	Ser
Tyr 225	Arg	Glu	Pro	Phe	Glu 230	Gln	His	Leu	Leu	·Leu 235	Gly	Val	Ser	Val	Ser 240
Cys	Ile	Val	Ile	Leu 245	Ala	Val	Cys <sub>.</sub>	Leu	Leu 250	Cys	Tyr	Val	Ser	Ile 255	Thr
Lys	Ile	Lys	Lys 260	Glu	Trp	Trp	Asp	Gln 265	Ile	Pro	Asn	Pro	Ala 270	Arg	Ser
Arg	Leu	Val 275	Ala	Ile	Ile	Ile	Gln 280	Asp	Ala	Gln	Gly	Ser 285	Gln	Trp	Glu
Lys	Arg 290	Ser	Arg	Gly	Gln	Glu 295	Pro	Ala	Lys	Cys	Pro 300	His	Trp	Lys	Asn

Cys 305	Leu	Thr	Lys	Leu	Leu 310	Pro	Cys	Phe	Leu	Glu 315	His	Asn	Met	Lys	Arg 320
Asp	Glu	Asp	Pro	His 325	Lys	Ala	Ala	Lys	Glu 330	Met	Pro	Phe	Gln	Gly 335	Ser
Gly	Lys	Ser	Ala 340	Trp	Cys	Pro	Val	Glu 345	Ile	Ser	Lys	Thr	Val 350	Leu	Trp
Pro	Glu	Ser 355	Ile	Ser	Va:l	Val	Arg 360	Суз	Val	Glu	Leu	Phe 365	Glu	Ala	Pro
Val	Glu 370	Cys	Glu	Glu	Glu	Glu 375	Glu	Val	Glu	Glu	Glu 380	Lys	Gly	Ser	Phe
Cys 385	Ala	Ser	Pro	Glu	Ser 390	Ser	Arg	Asp	Asp	Phe 395		Glu	Gly	Arg	Glu 400
Gly	Ile	Val	Ala	Arg 405	Leu	Thr	Glu	Ser	Leu 410	Phe	Leu	Asp	Leu	Leu 415	Gly
Glu	Glu	Asn	Gly 420	Gly	Phe	Cys	Gln	Gln 425	Asp	Met	Gly	Glu	Ser 430	Cys	Leu
Leu ·	Pro	Pro 435	Ser	Gly	Ser	Thr	Ser 440	Ala	His	Met	Pro	Trp 445	Asp	Glu	Phe
Pro	Ser 450	·Ala	Gly	Pro	Lys	Glu 455	Ala	Pro	Pro		Gly 460	Lys	Glu	Gln	Pro
Leu 465	His	Leu	Glu	Pro	Ser 470	Pro	Pro	Ala	Ser	Pro 475	Thr	Gln	Ser	Pro	Asp 480
Asn	Leų	Thr	Cys	Thr 485	Glu	Thr	Pro	Leu	Val 490	Ile	Ala	Gly	Asn	Pro 495	Ala ·
Tyr	Arg	Ser	Phe		Asn	Ser	Leu	Ser 505	Gln.	Ser	Pro	Cys	Pro 510	Arg	Glu
Leu	Gly	Pro 515	Asp	Pro	Leu	Leu	Ala 520	Arg	His	Leu	Glu	Glu 525	Val	Glu	Pro
Glu	Met 530	Pro	Cys	Val	Pro	Gln 535	Leu	Ser	Glu	Pro	Thr 540	Thr	Val	Pro	Gln
Pro 545	Glu	Pro	Glu	Thr	Trp 550	Glu	Gln	Ile	.Leu	Arg 555	Arg	Asn	Val	Leu	Gln 560

His	Gly	Ala	Ala	Ala 565	Ala	Pro	Val	Ser	Ala 570	Pro	Thr	Ser	Gly	Tyr 575	Gln
Glu	Phe	Val	His 580	Ala	Val	Glu	Gln	Gly 585	Gly	Thr	Gln	Ala	Ser 590	Ala	Val
Val	Gly	Leu 595	Gly	Pro	Pro	Gly	G1u 600	Ala	Gly	Tyr	Lys	Ala 605		Ser <sub>.</sub>	Ser
Leu	Leu 610	Ala	Ser	Ser	Ala	Val 615	Ser	Pro	Glu	Lys	Cys 620	Gly	Phe	Gly	Ala
Ser 625	Ser	Gly	Glu	Glu	630 Gjy	Tyr	Lys	Pro	Phe	Gln 635	Asp	Leu	Ile	Pro	Gly 640
Cys	Pro	Gly	Asp	Pro 645	Ala	Pro	Val	Pro	Val 650	Pro	Leu	Phe	Thr	Phe 655	Gly
Leu	Asp	Arg	Glu 660		Pro	Arg	Ser	Pro 665	Gln	Ser	Ser	His	Leu 670	Pro	Ser
Ser	Ser	Pro 675	Glu	His	Leu	_	Leu 680	Glu	Pro	Gly	Glu	Lys 685	Val	Glu	Asp
Met	Pro 690	Lys	Pro	Pro	Leu	Pro. 695	Gln	Glu	Gln	Ala	Thr 700	Asp	Pro	Leu	Val
Asp 705	Ser	Leu	Gly	Ser	Gly 710	Ile	Val	Tyr	Ser	Ala 715	Leu	Thr	Cys	His	Leu 720
Cys	Gly	His	Leu	<b>Lys</b> 725	Gln	Cys	His	Gly	Gln 730	Glu	Asp	Gŗ	Gly	Gln 735	Thr
Pro	Val	Met	Ala 740	Ser	Pro	Cys	Cys	Gly 745	Суѕ	Суз	Cys	Gly	Asp 750	Arģ	Ser
Ser	Pro	Pro 755	Thr	Thr	Pro	Leu	Arg 760	Ala	Pro	Asp	Pro	Ser 765	Pro	Gly	Gly
Val	Pro 770	Leu	Glu	Ala	Ser	Leu 775	Cys	Pro	Ala	Ser	Leu 780	Ala	Pro	Ser	Gly
Ile 785	Ser	Glu	Lys :	Ser	Lys 790	Ser	Ser	Ser	Ser	Phe 795	His	Pro	Ala	Pro	Gly 800
Asn	Ala	Gln	Ser	Ser 805	Ser	Gln	Thr	Pro	<b>Ly</b> s 810	Iļe	Val	Asn	Phe	Val 815	Ser

Val Gly Pro	o Thr Tyr M	et Arg Val	Ser		
•	820		825		
<210> 4					
•					
<211> 15			•		
<212> DNA					
<213> Homo	sapiens				
•					
<400> 4					
ttgcaccact	gcact				15
		•			
<210> 5	•			•	
<211> 15		-			
<212> DNA					
<213> Homo	sapiens				
	•				
<400> 5					
ttgcaccgct	gcact				15
				•	•
•					
<210> 6					
<211> 15					
<212> DNA					
<213> Homo	sapiens				
<400> 6					
	ht				15
ttttgtgcta	LLCCC,				13
	•				
<210> 7					
<211> 15					
<212> DNA					
<213> Homo	sapiens				
<400> 7					•
ttttgtgtta	ttccc				15
	•	•			
-2105 0					
<210> 8					
<211> 15	•				
<212> DNA					
<213> Homo	sapiens				
<400> 8					

ctgggccgct caggc

15

<210>	9					
<211>	15					
<212>	DNA					
<213>	Homo	sapiens				
<400>	9					
ctgggc	cctct	caggc	•			15
			*			•
<210>	10					
<211>	15					
<212>	DNA					
<213>	Homo	sapiens . ·				
<400>	10					
taagco	ctgcg	ctgga				<b>15</b>
	•					
<21,0>	11					
<211>	15					~
<212>	DNA	•				
<213>	Homo	sapiens				
<400>	11					
taagco	ctccg	ctgga				15.
	•					
<210>	12					
<211>	15					
<212>	DNA	•				
<213>	Homo	sapiens				•
•						
<400>	12		•			
agaaca	acgg	aggcg				15
						•
			•			
<210>			,			
<211>					-	
<212>						
<213>	Homo	sapiens				
<400>		•				
agaaca	aatgg	aggcg .				15
		•				

WO 01/04270		PCT/US00/19094
<211> 15		
<212> DNA		
<213> Homo sapier	ns	
<400> 14		
cggaggcgcg gggtg.		15
		-
. •	••	
<210> 15	•	
<211> 15	•	
<212> DNA	·	
<213> Homo sapier	ns .	
	•	
<400> 15		
cggaggcacg gggtg		15 .
	•	•
	•	•
<210> 16	· *	
<211> 15		
<212> DNA	ne	
(213) HOMO Sapre		•
<400> 16	•	•
gtgcggataa ctata	•	· 15
	•	•
<210> 17		
<211> 15		
<212> DNA	•	
<213> Homo sapier	ns	
<400> 17		
gtgcggacaa ctata		15
<210> 18		
<211> 15		
<212> DNA		
<213> Homo sapier	ns	
o cap_co		
<400> 18	•	
cggagtgcgg caggg		15
32 3 2 33		
<210> 19		
<211> 15		•
<212> DNA		•
<213> Homo sapier	ns	

WO 01/042	70			PC	CT/US00/19094
<400> 19					
cggagtgtgg	caggg				15
<210> 20					•
<211> 15 ·	. •				
<212> DNA					
<213> Homo	sapiens				
-		•			
<400> 20					
gcctgggctg	agggt				15
		•			
	•				
<210> 21	•				
<211>. 15					
<212> DNA	anniana				
<213> Homo	sapiens				
<400> 21					
gcctgggttg	agggt				15
<210> 22	•				
<211> 15					•
<212> DNA	•	·			
<213> Homo	sapiens				
<400> 22	•				
tggggtgggc	agggg				<b>15</b> .
					•
<210> 23		·			
<211> 15		•			-
<212> DNA					
<213> Homo	sapiens				•
4400× 00					
<400> 23					1 6
tggggtgagc	agggg	•			. 15
	•				•
<210> 24					
<211> 15					
<212> DNA					
<213> Homo	sapiens	•			
					•
<400> 24					
ttctcccgca	gtgaa				15

<210> 25		
<211> 15		•
<212> DNA		
<213> Homo	sapiens	
<400> 25		
ttctcccaca	gtgaa	15
<210> 26		
<211> 15		
<212> DNA	6	
<213> Homo	sapiens	
(400> 06	· ·	
<400> 26		15
gtgaaaacga	eccgg	13
		•
<210> 27		
<211> 15		
<212> DNA		
<213> Homo	sapiens	
	*	
<400> 27	•	
gtgaaaatga	cccgg	15
<210> 28		
<211> 15		
<212> DNA		
<213> Homo	sapiens	
<400> 20		
<400> 28	aaaa	15
ggcaagccct	ggggc	13
	·	·
<210> 29		
<211> 15		
<212> DNA	•	
<213> Homo	sapiens	
•		
<400> 29		
ggcaagctct	ggggc	15

WO 01/042	70	PC1/US00/19094
<211> 15		
<212> DNA		
<213> Homo	sanions	
\Z13> 1101110	Saptens	
<400> 30		
	tanat	15
gccctggggc	Lygat	15
<210> 31	•	
<211> 15		
<212> DNA	<u>.</u>	
<213> Homo	canione	
	Sapiens	
· <400> 31		
gccctggagc		15
geeeeggage .		13
<210> 32	•	
<211> 15 .		
<212> DNA		
<213> Homo	sapiens	
	TAP - SAID	
<400> 32		
atagcaaatc	ccagq	15
•		
<210> 33		
<211> 15		
<212> DNA		
<213> Homo	sapiens	
•		
<400> 33		
atagcaattc	ccagg	15 .
		. :
<210> 34		
<211> 15		
<212> DNA		•
<213> Homo	sapiens	
<400> 34		
gctctgccct	·aggca	15
•		
<210> 35		
<211> 15		
<212> DNA		
<213> Homo	saniens	

WO 01/04270		PCT/US00/19094
<400> 35.		
gctctgcact aggca		15
gototycact aggca		13
<210> 36		
<211> 15		
<212> DNA .		
<213> Homo sapiens		-
<400> 36		e.
ccccacccc tcaca	·	15
<210> 37		
<211> 15		•
<212> DNA		
<213> Homo sapiens		
<400> 37		4.5
ccccactcc tcaca		15
<210> 38		
<211> 15		
<212> DNA		
<213> Homo sapiens		•
<400> 38		
teceteegea <sub>,</sub> tegea		15
<210> 39		
<211> 15		
<212> DNA		
<213> Homo sapiens		•
<400> 39		
tecetecaca tegea		15
<210> 40		
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 40		
cacctgctgt ggtgt		15

<210> 41				
<211> 15	•			
<212> DNA				
<213> Homo	sapiens			•
<400> 41	•			
cacctgccgt	gatat			15
caccegecge	ggege			10
<210> 42	•			
<211> 15	•			
<212> DNA		· *.		
<213> Homo	saniens			
(220) Homo	bupicins			
<400> 42				
atgtctgaag	tagac			15
	cagao			10
<210> 43				
<211> 15				
<212> DNA				
<213> Homo	saniens			
(LIS) HOMO	baptono			
<400> 43	•			
atgtctgcag	tagac			15
		•		
<210> 44				
<211> 15				
<212> DNA				
<213> Homo	sapiens			
				•
<400> 44				
tgaccaacct	ttgct			15
		•		
<210> 45				٠
<211> 15				
<212> DNA		•		
<213> Homo	sapiens			
				•
<400> 45				
tgaccaactt	ttgct			15
	•			

WO 01/04270		PCT/US00/19094
<211> 15		
<212> DNA		
<213> Homo sapiens		
vers nome dapating		•
<400> 46		
cctgttttct ggagc		15
	•	•
<210> 47		
<211> 15		
<212> DNA ·		
<213> Homo sapiens	•	•
*		
<400> 47		15
cctgtttcct ggagc		15
•		
<210> 48 .		
<211> 15	•	
<212> DNA		
<213>.Homo sapiens	•	•
	•	*
<400> 48		
tggacctgct cggag		15
<2105 AD		
<210> 49 <211> 15		•
<211> 13 <212> DNA		
<213> Homo sapiens		•
(210) Homo Baptens		
<400> 49		
tggaccttct cggag		15
<210> 50		
<211> 15		
<212> DNA		
<213> Homo sapiens		
.· · ·		
<pre>&lt;400&gt; 50</pre>		15
agtcatgcct tcttc		13
	•	
<210> 51	. •	•
<211> 15		
<212> DNA		
<213> Homo sapiens	•	

WO 01/04270			PCT/US00/19094
<400> 51			1 r
agtcatgtct tcttc			15
			. ,
<210> 52			
<211> 15			•
<212> DNA			
<213> Homo sapiens			
<400> 52			15
geettettee acett	•		
	•		
<210> 53	•		
<211> 15	•		,
<212> DNA			
<213> Homo sapiens			
			•
<400> 53	·		: 15
geetteteee acett	• .		
			•
<210> 54			
<211> 15			
<212> DNA			
<213> Homo sapiens			
, , , , , , , , , , , , , , , , , , , ,			
<400> 54		-	15
cagcccccgt ctcgg ·	•		13
<210> 55	•	•	
<211> 15			
<212> DNA			
<213> Homo sapiens			
			•
<400> 55			15
cagcccctgt ctcgg			15
• •			
<210> 56			•
<211> 15	•		
<212> DNA	•		•
<213> Homo sapiens			•
<400> 56			_
ggagtttgta catgc			15

ggagtttgta catgc

<210> 57			
<211> 15			
<212> DNA	·		
<213> Homo	sapiens		
<400> 57			
			1.0
ggagtttata	catge		15
	•		
<210> 58	•		
<211> 15	·		
<212> DNA			
<213> Homo	sapiens		
	•		
<400> 58			
cagctcccca	gagca ·	•	15
_			
-			
<210> 59			
<211> 15			
<212> DNA	•		
<213> Homo	canione		•
\z13> 1101110	paptens		
<400> 59			
			15
cagctcctca	gagca	•	15
-040: -00			
<210> 60			
<211> 15			
<212> DNA			
<213> Homo	sapiens	-	
	•		
<400> 60			
agacaggicc	tegee	٠.	15
		•	
•			
<210> 61			
<211> 15		•	_
<212> DNA			
<213> Homo	saniens		
<400> 61	:		
	taras		1 6
agacagggcc	Legac		15
•	•		
.010	•		٠.
<210> 62			

WO 01/04270		PCT/US00/19094
<211> 15		
<212> DNA		•
<213> Homo sapiens		
<400> 62		
ctgcccctgg caatg		15
		10
<210> 63		
<211> 15	· .	•
<212> DNA		
<213> Homo sapiens		
•		
<400> 63		
ctgccccgg caatg		15
	u.	
		•
<210> 64		
<211> 15		
<212> DNA	•	
<213> Homo sapiens		
	·	
<400> 64		
aggtgcatgt cctct		15
•		
<210> 65		
<211> 15	•	
<212> DNA		
<213> Homo sapiens		
	·	•
<400> 65		
aggtgcacgt cctct		15
	•	
		•
<210> 66		
<211> 15		
<212> DNA		9
<213> Homo sapiens		,
		-
<400> 66		
gtgcatgtcc tcttg		15
		•
<210> 67		
<211> 15		
<212> DNA		

<213> Homo sapiens

WO 01/04270		PCT/US00/19094
<400> 67		
		15
gtgcatgccc tettg		
<210> 68		
<211> 15	•	
<212> DNA		
<213> Homo sapiens	•	
vais nome baptens		
<400> 68-		
ggcttatcca tgcct		15
<210> 69		
<211> 15		•
<212> DNA		· ·
<213> Homo sapiens	•	•
	·	
<400> 69		. :
ggcttattca tgcct		15
<210> 70	* .	
<211> 15		
<212> DNA		
<213> Homo sapiens	•	
•	·	
<400> 70	•	
agccaggctg gcaga		15
		•
<210> 71		
<211> 15		
<212> DNA	•	
<213> Homo sapiens		
<400> 71		
agccagggtg gcaga	·	15
	· ·	13
<210> 72		
<211> 15		·
<212> DNA		
<213> Homo sapiens	• •	
	•	
<400× 72		

ggcccacatg gaggc

<210> 73				
<211> 15				
<212> DNA				
<213> Homo sapiens				
•				
<400> 73				•
ggcccacgtg gaggc	. *			15
<210> 74				
<211> 15				
<212> DNA				
<213> Homo sapiens				
•				
<400> 74				• .
taacacagcc atcaa			_	15
•				
<210> 75				
<211> 15				
<212> DNA				
<213> Homo sapiens				
•				
<400> 75				
taacacaacc atcaa	•			. 15
•	•			
•				
<210> 76				
<211> 15				
<212> DNA				
<213> Homo sapiens				
•				
<400> 76				
taatgctcgt ctgtg				15
			•	
<210> 77				
<211> 15				
<212> DNA				
<213> Homo sapiens				
<400> 77				
taatgcttgt ctgtg		•		15
	•			
.010. 50				
<210> 78				

WO 01/04270	PCT/US00/19094
<211> 15	•
<212> DNA	
<213> Homo sapiens	
<400> 78	
acttgccgtc tgggt	15 <sup>.</sup>
	*
<210> 79	
<211> 15	
<212> DNA	
<213> Homo sapiens	
<400> 79	
acttgccatc tgggt	15
•	
<210> 80	,
<211> 15	
<212> DNA	
<213> Homo sapiens	
	•
<400> 80	
ctgagattgc accac	15
<210× 01	
<210> 81 <211> 15	·
<212> DNA	
<213> Homo sapiens	
<400> 81	_
ggctggagtg cagtg	15
<210> 82	
<211> 15	•
<212> DNA	
<213> Homo sapiens	
•	
<400> 82	
ctgagattgc accgc	15
•	•
<210> 83	
<211> 15	
<212> DNA	

<213> Homo sapiens

WO 01/04270		PCT/US00/1909
<400> 83		
ggctggagtg cageg		15
<210> 84		٠
<211> 13 <212> DNA		
<213> Homo sapiens		
<400> 84		- 00
ctgtgctttt gtgct		15
<210> 85		
<211> 15	•	-
<212> DNA		
<213> Homo sapiens		
<400> 85		
accaagggga atagc		15
		-
<210> 86		
<211> 15	•	
<212> DNA		•
<213> Homo sapiens		•
<400> 86		
ctgtgctttt gtgtt	~	15
÷ .		
<210> 87	•	
<211> 15	•	
<212> DNA		
<213> Homo sapiens		
<400> 87		
accaaġggga ataac		15
		•
<210> 88		•
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 88		
gagttcctgg gccgc		15

	15 DNA Homo	sapiens		•			
<400>							
ggagca	igcct	gagcg					15
•							
<210>	90						
<211>							
<212>							
<213>	Homo	sapiens					
	•						
<400>	90						
gagtto	ctgg	gcctc .	•				15
			•				
<b>4010</b> 5	01 .						•
<210> <211>							•
<212>							
		sapiens				•	·
<400>	91						
ggagca	gcct	gagag					15
<21 Os	02					•	
<210> <211>							
<212>		•	,				
		sapiens	•				•
<400>	92		:				
tccgag	ıtaag	cctgc.					15
							•
<b>4010</b> 5	00	•					
<210> <211>							
<212>							•
		sapiens					
		•					
<400>	93						
tccago	tcca	gcgca		•			15

WO 01/04270		PCT/US00/19094
<211> 15	·	
<212> DNA		
<213> Homo sap	i ens	
vers nome sap	10113	
.<400> 94		•
tccgagtaag cct	cc	15
cocyaycaay coc		13
	• .	
<210> 95		
<211> 15		
<212> DNA		
<213> Homo sap	i ens	•
		•
<400> 95		
tecageteca geg	qa	15
		:
<210> 96		
<211> 15		•
<212> DNA		
<213> Homo sap	iens	
<400> 96		
tccctgagaa caa	cg	15
	•	
<210> 97	•	
<211> 15		
<212> DNA	•	
<213> Homo sap	iens	
<400> 97		
accccgcgcc tcc	gt .	15
•		
·		•
<210> 98	•	•
<211> 15	•	
<212> DNA	iona	
<213> Homo sap	Tens .	
<400>.98	•	•
tccctgagaa caa	ta	15
coccegagaa caa	-y	13
<210> 99		
<211> 15		
<212> DNA		
<213> Homo sap	iens	
ome bup		

WO 01/042	70			PCT/US00/19094
<400> 99			•	
acccgcgcc	tccat			15
3.5				
<210> 100				·
<211> 15				
<212> DNA				
<213> Homo	sapiens .			
<400> 100				
gaacaacgga	ggcgc			15
•				
<210> 101				
<211> 15				
<212> DNA	•			
<213> Homo	sapiens			
<400> 101	•			•
cacacgcacc	ccaca			15
cacacycacc	cegeg			<b>±</b> 5 .
	• •			
<210> 102				
<211> 15				•
<212> DNA				
<213> Homo	sapiens			
<400> 102				• •
gaacaacgga	ggcac			15
<210> 103				
<211> 15				
<212> DNA				
<213> Homo	sapiens			
<400> 103	•			
<400> 103	aaata .			1.5
cacacgcacc	ccgeg			15
<210> 104	-			
<211> 15				
<212> DNA				
<213> Homo	sapiens			;
	· •			
<400> 104				•
tggtcagtgc	ggata			15

<210> 105				
<211> 15				
<212> DNA				
<213> Homo sapiens				
•				• .
<400> 105				
ccagtgtata gttat				15
<210> 106			÷ .	•
<211> 15	•			
<212> DNA				
<213> Homo sapiens				
•				
<400> 106				•
gcagggcgga gtgcg				<b>15</b> .
			•	
<210> 107				
<211> 15				
<212> DNA <213> Homo sapiens				
(213) Homo Saptems				
<400> 107				
agccacccct gccgc	•			15
•			,	•
	•			
<210> 108	•			
<211> 15				
<212> DNA				
<213> Homo sapiens				
<400> 108	·			
gcagggcgga gtgtg				15
			•	
•	•			
<210> 109				
<211> 15				
<212> DNA				
<213> Homo sapiens	ı			
<400× 100				
<400> 109 agccacccct gccac				15
agocaccocc -goods				-~
<210> 110				

WO 01/042/0	PC1/US00/19094
<211> 15	
<212> DNA	
<213> Homo sapiens	
<400> 110	
acagctgcct gggct	15
333	
	•
<210> 111	•
<211> 15	•
<212> DNA	
<213> Homo sapiens	
<400> 111	•
caccccaccc tcage .	15
<210> 112	
<211> 15 .	
<212> DNA	
<213> Homo sapiens	
<400> 112	
acagctgcct gggtt	15
•	
<210> 113	
<211> 15	
<212> DNA	
<213> Homo sapiens	•
	·
<400> 113	عال
acagctgcct gggtt	<b>15</b> .
(010) 114	
<210> 114	
<211> 15	
<212> DNA <213> Homo sapiens	
(213) Homo Sapiens	•
<400> 114	•
tgagggtggg gtggg	15
	•
<210> 115	
<211> 15	
<212> DNA	
<213> Homo sapiens	•
_	

<b>WO</b> 01/04270		PCT/US00/19094
		•
<400> 115		·
cetectece tgece		15
cccccccc cgccc		13
<210> 116		•
<211> 15		
<212> DNA		
<213> Homo sapiens		
,	•	
<400> 116		
tgagggtggg gtgag		15
2 333 239 3 3 2		
	. *	
<210> 117		
<211> 15	•	
<212> DNA		
<213> Homo sapiens		
•		
<400> 117		
cctcctcccc tgctc		15
<210> 118		
<211> 15		
<212> DNA		
<213> Homo sapiens	•	
		•
<400> 118		
ggccgcttct cccgc		15
4010x 110		. 00
<210> 119		
<211> 15 <212> DNA		
<213> Homo sapiens		
VZISV NOMO Sapiens	•	
<400> 119		
ctgggtttca ctgcg		15
organication, organi		
		•
<210> 120	•	
<211> 15		
<212> DNA	•	
<213> Homo sapiens		
	•	
<400> 120		·

15 、

ggccgcttct cccac

<210> 121 <211> 15 <212> DNA			
<213> Homo	sapiens		
<400> 121			
ctgggtttca	ctgtg		15
<210> 122 <211> 15			
<211> 15 <212> DNA			
<213> Homo	sapiens		
<400> 122	•		
tttggagtga	aaacg		15
<210> 123			
<211> 15			
<212> DNA			
<213> Homo	sapiens		
<400> ·123			
catctgccgg	gtcgt		15
	-8		
<210> 124	·		
<211> 15	•		
<212> DNA			
<213> Homo	sapiens		
<400> 124			
tttggagtga	aaatg		15
<210> 125	•		
<211> 15			
<212> DNA			
<213> Homo	sapiens		
<400> 125			
catctgccgg	gtcat		15
, ,,			
<210> 126			

WO 01/04270		PCT/US00/19094
<211> 15		
<212> DNA		
<213> Homo sapiens		
(215) Homo Baptons		
<400> 126		
ctgggaggca agccc	,	15
<210> 127		
<211> 15		
<212> DNA .		
<213> Homo sapiens		
<400> 127		
tatccagccc caggg		15
•	• .	
<210> 128		
<211> 15 ·		•
<212> DNA		•
<213> Homo sapiens		
4400 400	•	
<400> 128		15
tggtcagtgc ggaca		. 13
<210> 129		
<211> 15		
<212> DNA	-	
<213> Homo sapiens		
<400> 129		
ccagtgtata gttgt		15
	÷	
<210> 130		
<211> 15		•
<212> DNA		•
<213> Homo sapiens		
(470) 120		·
<400> 130		
ctgggaggca agctc		15
<210> 131		•
<211> 15		·
<211> 13 <212> DNA		
<213> Homo sapiens		
	•	

WO 01/04270		PCT/US00/19094
<400> 131		
tatccagccc cagag		15
cacccagece cagag		
•		
<210> 132		
<211> 15		
<212> DNA		
<213> Homo sapiens		
•		
<400> 132		
aggcaagccc tgggg		15
•		
<210> 133		•
<211> 15		
<212> DNA		•
<213> Homo sapiens		•
00		•
<400> 133		1.5
tttgctatcc agccc		15
•		•
<210> 134		
<211> 15		
<212> DNA		
<213> Homo sapiens		•
<400> 134		
aggcaagccc tggag		15
•		•
<210> 135		
<211> 15		
<212> DNA .		
<213> Homo sapiens		
<400> 135	* '	15
tttgctatcc agctc		15
<210> 136		
<211> 156 <211> 15		
<212> DNA		
<213> Homo sapiens		
nomo ouprono		·
<400> 136		
ggctggatag caaat		15
_		

<210> 137		
<211> 15		
<212> DNA		
<213> Homo	saniens	
10110		
<400> 137		
		4 -
ctageteetg	ggatt	15
<210> 138		
<211> 15	•	•
<212> DNA		
<213> Homo	sapiens	
<400> 138		•
ggctggatag	caatt	15
<210> 139		
<211> 15	•	
<212> DNA		
<213> Homo	sapiens	
<400> 139		
ctagctcctg	duant	15
	55000	15
		,
<210> 140		
<211> 15		
<212> DNA	ni.	•
	· · · · · · · · · · · · · · · · · · ·	
<213> Homo	saptens	
4400> 140	•	·
<400> 140		
cacctggctc	tgccc	15
		•
<210> 141		
<211> 15		
<212> DNA	•	
<213> Homo	sapiens	
•-		
<400> 141	•	
gggacttgcc	taggg	15
•	•	
	•	
<210> 142	•	

WO 01/04270	PCT/US00/19094
<211> 15	•
<212> DNA	
<213> Homo sapiens	
(213) Homo sapiens	•
<400> 142	
•	15
cacctggctc tgcac	15
•	
<210> 143	
<211> 15	
<212> DNA	
<213> Homo sapiens	•
<400> 143	
gggacttgcc tagtg	15
<210> 144	
<211> 15 ` .	
<212> DNA	•
<213> Homo sapiens	
	• .
<400> 144	
cctggcccc caccc .	15
<210> 145	,
<211> 15	•
<212> DNA	
<213> Homo sapiens	· 2
<400> 145	·
ctctgatgtg agggg	15
<210> 146	
<211> 15	
<212> DNA	
<213> Homo sapiens	·
<400> 146	,
cetggcccc cactc	15
congression carrie	. 13
•	
<210\ 147	
<210> 147	
<211> 15	
<212> DNA	
<213> Homo sapiens	

WO 01/04	270		PCT/US00/19094
<400> 147		•	•
ctctgatgtg	aggag		15
<210> 148			
<211> 15			
<212> DNA	•		
<213> Homo	sapiens		
<400> 148			
gaaccctccc	tccgc		15
<210> 140			•
<210> 149 <211> 15			
	•		
<212> DNA <213> Homo	canions		
\213/ NOMO	sabrens		·
<400> 149			
gctggctgcg	atgcg		15
<210> 150	•		
<211> 15			
<212> DNA			
<213> Homo	sapiens		
<400> 150	. •		
gaaccctccc	tccac		15
<210> 151			
<211> 151	. •		
<212> DNA			
<213> Homo	sapiens		•
	• .		
<400> 151			15 '
gctggctgcg	acgcg		15
•			
<210> 152			
<211> 15			
<212> DNA			
<213> Homo	sapiens .		
<400> 152			•
tagatacacc	tacta		15
Lagacacacc	egeeg .		13

<210> 153			
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 153		•	
gcagatacac cacag			1.5
· ·			•
,			
<210> 154	•		
<211> 15			
<212> DNA			
<213> Homo sapiens			
-			-
<400> 154			
tagatacacc tgccg			15
		•	
•			
<210> 155			
<211> 15			
<212> DNA	•		
<213> Homo sapiens			
<400> 155			•
gcagataçac cacgg			15
<210> 156			•
<211> 15			
<212> DNA			•
<213> Homo sapiens			
•			•
<400> 156	•		
gaaggcatgt ctgaa		•	15
<210> 157			
<211> 15			
<212> DNA		•	
<213> Homo sapiens		•	,
	•		
<400> 157			
atggctgtct acttc		•	15
<210> 158			

WO 01/04270	*	PC17US00/19094
<211> 15		
	•	
<212> 'DNA		
<213> Homo sapiens		
<400> 158		
gaaggcatgt ctgca		15
	•	
	•	•
<210> 159	•	
<211> 15		
<212> DNA	•	
<213> Homo sapiens	•	
<400> 159		
atggctgtct actgc		15
acygoigice accyc		13 .
•		
-010- 160		
<210> 160		
<211> 15		
<212> DNA		
<213> Homo sapiens		
	•	
<400> 160		
gaaccctgac caacc		15
	•	
		•
<210> 161		
<211> 15		
<212> DNA ·		
<213> Homo sapiens	*	
	•	
<400> 161 .		
tgcaaaagca aagga		15
<210> 162	•	
<211> 15		
<212> DNA		
<213> Homo sapiens		
\ZIS> homo saprens		
	·	
<400> 162		1 5
gaaccctgac caatc		15
•		
	•	
<210> 163		•
<211> 15		
<212> DNA		
<213> Homo sapiens		

WO 01/04270	-X-		PCT/US00/19094
<400> 163			
tgcaaaagca aagaa	•		15
egodddydd ddydd			
<210> 164			
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 164			
tettgecetg tttte	•		15
***************************************			
<210> 165			
<211> 15			
<212> DNA			
<213> Homo sapiens			•
<400> 165			
tgttgtgctc cagaa			15
egeegeee cagaa	-		10
•			
<210> 166			
<211> 15			•
<212> DNA			
<213> Homo sapiens			•
<400> 166 tettgeeetg tttee			15
ceregeous cree			. 13
<210> 167			
<211> 15			
<212> DNA			
<213> Homo sapiens			
.400: 450			
<400> 167			15
tgttgtgctc cagga			13
<210> 168			
<211> 15			
<212> DNA	• •		
<213> Homo sapiens			
<400> 168		•	1.5
tgttcctgga cctgc			15

<210> 169		
<211> 15		
<212> DNA		
<213> Homo	sapiens	
<400> 169		•
tctcctctcc	gagca	15
•	•	
<210> 170		
<211> 15	•	
<212> DNA		
<213> Homo	sapiens	
<400> 170		
tgttcctgga	ccttc	15
<210> 171		
<211> 15	• •	
<212> DNA		*
<213> $Homo$	sapiens	
<400> 171		
tctcctctcc	gagaa	15 ·
-010: 120		
<210> 172	. '	
<211> 15 <212> DNA	•	
<213> Homo	caniane	
12157 1101110	·	
<400> 172	•	
tgggggagtc	atgcc	15
-55555-5		~ 0
<210> 173	•	
<211> 15		
<212> DNA		
<213> Homo	sapiens	
· <400> 173	•	
aaggtggaag	aaggc	15

WO 01/042/0		PC17US00/19094
<211> 15		
<212> DNA	•	
<213> Homo sapiens		•
<400> 174		
tgggggagtc atgtc		15 .
<210> 175	•	
<211> 15		
<212> DNA		
<213> Homo sapiens	• ()	
<400> 175		
aaggtggaag aagac		15
<210> 176	·	
<211> 15		
<212> DNA		
<213> Homo sapiens		
:		
<400> 176		
agtcatgcct tcttc		15
		13
· <010 \ 177		
<210> 177		
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 177		
ttcccgaagg tggaa		15
·*		
<210> 178		
<211> 15		
<212> DNA		
<213> Homo sapiens		•
<400> 178		•
		15 '
agtcatgcct tctcc		15 ·
<210> 179		
<211> 15		
<212> DNA		

<213> Homo sapiens

WO 01/04270	•	PCT/US00/19094
<400> 179	•	
ttcccgaagg tggga		15
	·	
<210> 180		
<211> 15	·	
<212> DNA		<u>.</u>
<213> Homo sapiens		
<400> 180		
cagetgeage eeeeg		15
-	•	
<210> 181		
<211> 15 ,		•
<212> DNA		•
<213> Homo sapiens		
<400> 181		
tgggggccga gacgg		15
<210> 182		
<211> 15		
<212> DŅA		
<213> Homo sapiens		
<400> 182		
cagetgeage ceetg		15
3, 3		
<210> 183	•	•
<211> 15		
<212> DNA		•
<213> Homo sapiens		
<400> 183		
tgggggccga gacag		15
	•	
•		
<210> 184		
<211> 15		
<212> DNA		•
<213> Homo sapiens		
<400> 184		15

WO 01/04270	PCT/US00/19094

<210> 185					
<211> 15					
<212> DNA					
<213> Homo	sapiens				
		•	•	•	•
<400> 185					
tccaccgcat	gtaca			•	15
•					
<210> 186					•
<211> 15					
<212> DNA					
<213> Homo	sapiens				
	•				
<400> 186					
ctatcaggag	tttat				15
•.					
<210> 187					
<211> 15	•				
<212> DNA					
<213> Homo	sapiens				
<400> 187					3.5
<400> 187 tccaccgcat	gtata				15
	gtata	·			15
tccaccgcat	gtata				15
tccaccgcat	gtata				15
<pre>tccaccgcat &lt;210&gt; 188 &lt;211&gt; 15</pre>	gtata		•		15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA</pre>					15 
<pre>tccaccgcat &lt;210&gt; 188 &lt;211&gt; 15</pre>					15
<210> 188 <211> 15 <212> DNA <213> Homo					15
<210> 188 <211> 15 <212> DNA <213> Homo <400> 188	sapiens				
<210> 188 <211> 15 <212> DNA <213> Homo	sapiens				15 
<210> 188 <211> 15 <212> DNA <213> Homo <400> 188	sapiens				
<210> 188 <211> 15 <212> DNA <213> Homo <400> 188 cccaagcagc	sapiens				
<210> 188 <211> 15 <212> DNA <213> Homo <400> 188 cccaagcagc <210> 189	sapiens				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15</pre>	sapiens				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15 &lt;212&gt; DNA</pre>	sapiens				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15</pre>	sapiens				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15 &lt;212&gt; DNA</pre>	sapiens				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 189</pre>	sapiens tcccc				15
<pre>&lt;210&gt; 188 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo &lt;400&gt; 188 cccaagcagc &lt;210&gt; 189 &lt;211&gt; 15 &lt;212&gt; DNA &lt;213&gt; Homo</pre>	sapiens tcccc				15

WO 01/04270			PCT/US00/19094	
<211> 15				
<212> DNA				
<213> Homo sapiens				
version suprems				
<400> 190				
	•		15	
cccaagcagc tcctc			15	•
<210> 191				
<211> 15				
<212> DNA				
<213> Homo sapiens				
.400> 101			•	
<400> 191	•	•	15	
cccaggtgct ctgag			15	
-010- 100			•	
<210> 192	•			
<211> 15	•	•		
<212> DNA	•			
<213> Homo sapiens				
<400× 100				
<400> 192			15	•
ctgtggagac aggtc			15	
<210× 102	•		•	
<210> 193				
<211> 15 · · · · · · · · · · · · · · · · · ·				
<213> Homo sapiens				
<400> 193	•	•		
	•		15	
gtagggggcg aggac			,	
				·
<210> 194 ·				
<211> 154				
<212> DNA			•	
<213> Homo sapiens	•			
VZISV NOMO Bupichs				
<400> 194			•	
ctgtggagac agggc			15	
cegeggagae aggge				
<210> 195			•	
<210> 193 <211> 15				
	•			
<212> DNA				
<213> Homo sapiens				

<b>WO</b> 01/04270	PCT	/US00/19094
<100 195		
<400> 195		16
gtagggggg aggcc		15
<210> 196		
<211> 15		
<212> DNA		
<213> Homo sapiens	•	
(213) Homo Saptens	•	
<400> 196		
tccatcctgc ccctg		15
. •		•
<210> 197		
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 197		
tctgagcatt gccag		15
	•	
<210> 198 ·		
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 198		•
tccatcctgc ccccg		15
<210> 199		
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 199		
tctgagcatt gccgg		15
•		
1010> 000		
<210> 200		
·<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 200		
<400> 200		1.5
tctcttaggt gcatg .		15

<210> 201	•							
<211> 15								
<212> DŅA								
<213> Homo	sapiens							
	-							
<400> 201				•	•			
gcaacaagag	gacat							15
goddoddgag	gaoac							
٠								
<210> 202						•		
<211> 202								
<211> 15 <212> DNA								
<213> Homo	sapiens							
-100000	•							
<400> 202								
tetettaggt	gcacg							15
<210> 203								
<211> 15								
<212> DNA								
<213> Homo	sapiens							
						•		•
<400> 203	•							•
gcaacaagag	gacgt							15
	•							
<210> 204								
<211> 15								
<212> DNA	•							
<213> Homo	sapiens							
<400> 204	•	•			•			
tcttaggtgc	atgtc							15
		•	,					
<210> 205								
<211> 15								
<212> DNA					•			
<213> Homo	saniens							
	P							
<400> 205								
	20020							15
cagcaacaag	ayyac							13
10105 000							•	
<210> 206								

WO 01/04270			PCT/US00/19094
<211> 15			
<212> DNA			
<213> Homo sapiens	•		
<400> 206			
tettaggtge atgee:	•		15
<210> 207			
<211> 15			•
<212> DNA .			
<213> Homo sapiens	•		•
	•		
<400> 207	•		
cagcaacaag agggc	•		15
•		•	
			•
<210> 208	•	•	
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 208			
gactagggct tatcc ·			15
9400499900 04000			
		•	
<210> 209			
<211> 15			
<212> DNA			
<213> Homo sapiens	•		
		•	
<400> 209		,	
tttcccaggc atgga		•	15
•			
<210> 210		•	
<211> 15	•		
<212> DNA			•
<213> Homo sapiens			
<400> 210			•
gactagggct tattc		•	15 .
gactaggget tatte			
<210> 211		•	
<211> 15			
<212> DNA			
<213> Homo sapiens			

WO 01/04270			PCT/US00/19094
			4
<400> 211			1 5
tttcccaggc atgaa			15
<210> 212			-
<211> 15			
<211> 13 <212> DNA	•		
<213> Homo sapiens			
			<i>:</i>
<400> 212			
gaaggcagcc aggct			15
<210> 213 . ·			•
<211> 15	•		•
<212> DNA			
<213> Homo sapiens			
<400> 213			
tggaaatctg ccagc			. 15
	•		
<210> 214			
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 214			15
gaaggcagcc agggt			13
<210> 215	•		
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 215	*	•	
tggaaatctg ccacc			15
<210> 216			
<211> 15	·		
<212> DNA			
<213> Homo sapiens			
		·	
<400> 216			15
gatcatggcc cacat			13

<210> 217	•				
<211> 15	•	•			
<212> DNA				-	
<213> Homo	sapiens				
	• )				
<4.00> 217				• .	
aggtgggcct	ccatg				15
			•		
			•		
<210> 218					
<211> 15					•
<212> DNA					•
<213> Homo	sapiens .				
<400> 218					
gatcatggcc	cacgt				15
				•	
<210> 219					
<211> 15					
<212> DNA		•			•
<213> Homo	sapiens				
					•
<400> 219					4 =
aggtgggcct	ccacg			•	15
				•	
<210> 220					
<211> 15				,	
<212> DNA			•		-
<213> Homo	sapiens			•	
•					
<400> 220		,			
agaaactaac	acagc				15
	•				
<210> 221					
<211> 15					
<212> DNA					
<213> Homo	sapiens				
•					
<400> 221					
attcccttga	tggct				15

<210> 222

WO 01/04270		PCT/US00/19094
<211> 15		
<212> DNA		
<213> Homo sapiens		
<400> 222		
agaaactaac acaac		15
		•
<210> 223		
<211> 15		
<212> DNA		
<213> Homo sapiens		
-		
<400> 223		
attcccttga tggtt		15
<210> 224	-4-	
<210> 224 <211> 15		
<212> DNA	·	
<213> Homo sapiens		•
*		
<400> 224	· .	
gttgagtaat gctcg		15
	•	
<210> 225		•
<211> 15 <212> DNA		
<213> Homo sapiens		•
valor nomo bapacino	•	
<400> 225		
aaaacacaca gacga		15
	•	•
<210> 226	•	
<211> 15		
<212> DNA <213> Homo sapiens		
VZ137 NOMO Saptems		
<400> 226		
gttgagtaat gcttg	·	15
	•	·
·	•	
<210> 227		
<211> 15		
<212> DNA		

WO 01/04270			PCT/US00/19094
<400> 227			•
aaaacacaca gacaa			15
•			
<210> 228			
<211> 15			
<212> DNA			
<213> Homo sapiens			
<400> 228			•
taagaaactt geegt			15
			•
<210> 229	•		
<211> 229			
<211> 13 <212> DNA			
<213> Homo sapiens			
Tallor Home Bapaons	•		
<400> 229			* .
acccaaaccc agacg			<b>15</b> .
<210> 230			
<211> 15			
<212> DNA			•
<213> Homo sapiens			
<400> 230			
taagaaactt gccat			15
•			•
<210> 231	•		
<211> 15 .			
<212> DNA			
<213> Homo sapiens			
<400> 231			•
acceaaacce agatg			15
acceasacce agacg	•		15
<210> 232	•		
<211> 10		•	
<212> DNA			
<213> Homo sapiens			
<400> 232			
agattgcacc			10

<210> 233	•	
<211> 10		
<212> DNA		
<213> Homo	saniens	
\213> NOMO	Sapiens	
44005 000	•	
<400> 233		
tggagtgcag		10
<210> 234		
<211> 10 .		
<212> DNA		
<213> Homo	sapiens	
<400> 234		
tgcttttgtg	*	10
<210> 235		
<211> 10		
<212> DNA		
<213> Homo	anniona	
(213) HOMO	saptens	
<400> 235		
		10
aaggggaata		10
Z2105 226		
<210> 236		•
<211> 10		
<212> DNA	st.	
<213> Homo	sapiens	
<400> 236		•
ttcctgggcc		
		10
		10
		10
<210> 237		10
<210> 237 <211> 10		10
		10
<211> 10 <212> DNA		10
<211> 10		
<211> 10 <212> DNA <213> Homo		
<211> 10 <212> DNA <213> Homo <400> 237	sapiens	
<211> 10 <212> DNA <213> Homo	sapiens	10
<211> 10 <212> DNA <213> Homo <400> 237	sapiens	
<211> 10 <212> DNA <213> Homo <400> 237	sapiens	

WO 01/04270		PCT/US00/19094
<211> 10	•	•
<212> DNA		
<213> Homo sapiens		
<400> 238		
gagtaagcct		10
	•	
<210> 239		
<211> 10		•
<212> DNA		·
<213> Homo sapiens		
		•
<400> 239		
agctccagcg	•	10
•		
.010. 010		•
<210> 240		
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 240		
ctgagaacaa		10
ctgagaacaa		10
<210> 241		
<211> 10		•
<212> DNA		
<213> Homo sapiens		
	•	
<400> 241		• •
ccgcgcctcc	•	10
<210> 242	•	
<211> 10		•
<212> DNA ·		
<213> Homo sapiens		
<400> 242		
caacggaggc		10 .
<01.0× 0.40		
<210> 243		
<211> 10		
<212> DNA		

WO 01/04270	PCT/US00/19094
<400> 243 acgcaccccg	 10
<210> 244 <211> 10 <212> DNA <213> Homo sapiens	
<400> 244 tcagtgcgga	10
<210> 245 <211> 10 <212> DNA <213> Homo sapiens	
<400> 245 gtgtatagtt	10
<210> 246 <211> 10 <212> DNA <213> Homo sapiens	vo -
<400> 246 gggcggagtg	10
<210> 247 <211> 10 <212> DNA <213> Homo sapiens	
<400> 247 cacccetgec	10
<210> 248 <211> 10 <212> DNA <213> Homo sapiens	· .
<400> 248 gctgcctggg	. 10

<210>	249							
<211>								
<212>								
		sapiens						
\213/	пошо	saprens						
<400>	249							
cccaco			•					10
CCCacc								10
			•					
<210>	250					•		
<211>			•					
<212>		•						
		sapiens .	•					
12132	пошо	saprens .						
<400>	250							
gggtgg								10
333535	199-9							10.
			•		·	•		
<210>	251		•					
<211>							·	
<212>								
		sapiens						
12107	nomo	Dupiens						
<400>	251-							
cctccc					-			10
	_			•		•		
<210>	252							
<211>		• (						
<212>								
		sapiens						
<400>	252							
cgcttc	tccc							10
<210>	253							
<211>	10							
<212>	DNA							
<213>	Homo	sapiens						
					٠			
<400>	253							
ggtttc	actg							10
Z2105	254		•					

WU 01/042/0	PC1/	US00/19094
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 254		
ggagtgaaaa .		10
ggagtgaaaa		10
<210> 255		
<211> 10		
<212> DNA		•
<213> Homo sapiens		
<400> 255		
ctgccgggtc		10
• •		
<210> 256		
<211> 10		
<212> DNA		
<213> Homo sapiens		
Total Total Superior		
.400. 056		
<400> 256		
ggaggcaagc	•	10
<210> 257		
	•	
<211> 10		
<212> DNA		•
<213> Homo sapiens		
<400> 257		
	•	10
ccagccccag		10
·	•	
<210> 258		
<211> 10	•	
<212> DNA	•	
<213> Homo sapiens		
ZTON HOMO Sabrems		•
	w.	
<400> 258		
caagccctgg		10
.010. 050		
<210> 259		
<211> 10 ,		•
<212> DNA		
<213> Homo sapiens		

WO 01/04270		PCT/US00/19094
<400> 259		
getatecage		10
goodoodgo	•	
	•	
<210> 260		
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 260		
tggatagcaa	•	10
	•	•
<210> 261	*	
<211> 10	·	
<211> 10 <212> DNA		
<213> Homo sapiens		
Tomb Supremo		
<400> 261		
gctcctggga		10
<210> 262		
<211> 10		•
<212> DNA		
<213> Homo sapiens		
		•
<400> 262	2	
ctggctctgc	· .	10
-		
<210> 263	8	
<211> 10		
<212> DNA		•
<213> Homo sapiens		•
2		
<400> 263		
acttgcctag		10
<210> 264		
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 264		,
ggcccccac		10

<210> 265							
<211> 10							
<212> DNA							
<213> Homo sapiens							
<400> 265							
tgatgtgagg							10
~9~0909dg							
	•						
<210> 266					•		
<211> 10							
<212> DNA							
			•				
<213> Homo sapiens							
<400> 266	•					•	10
ccctcctcc		,	•				10
.010. 057							
<210> 267							
<211> 10				,			
<212> DNA							
<213> Homo sapiens				-			
<400> 267						•	
ggctgcgatg			•				10
•			•				•
<210> 268			-				
<211> 10			•	•	•		
<212> DNA	•						
<213> Homo sapiens							
			•				
<400> 268							
atacacctgc				·			10
•							
<210> 269							
<211> 10 .							
<212> DNA							
<213> Homo sapiens							
<400> 269							
gatacaccac							10
=							
<210> 270							

WO 01/04270		PCT/US00/19094
<211> 10		
<212> DNA	•	
<213> Homo sapiens		• (0
	•	
<400> 270		
ggcatgtctg		10
·		
<210> 271		
<211> 10		
<212> DNA		•
<213> Homo sapiens		
•	·	
<400> 271		
gctgtctact		10
<210> 272 <sup>-</sup>		
<211> 10		
<212> DNA		_
<213> Homo sapiens	•	
<400> 272		
ccctgaccaa		10
-		
<210> 273		
<211> 10	•	
<212> DNA		
<213> Homo sapiens		•
_		•
<400> 273		
aaaagcaaag		10
<210> 274		•
<211> 10		•
<212> DNA		
<213> Homo sapiens		
•		
<400> 274	•	
tgccctgttt		10
•		
<210> 275		
<211> 10		
<212> DNA		

WO 01/04270		PCT/US00/19094
<400> 275	•	
tgtgctccag		10
·		
<210> 276		
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 276		
tcctggacct		10
		10
·		•
<210> 277		•
<211> 10		
<212> DNA		•
<213> Homo sapiens		
<400> 277		
cctctccgag .		10
•	•	
		•
<210> 278		
<211> 10		
<212> DNA	*	
<213> Homo sapiens		
<400> 278		
gggagtcatg		10
<210> 279		•
<211> 10 <212> DNA	•	
<213> Homo sapiens		
verior nomo ouprono		
<400> 279		
gtggaagaag		10
<2105 200 ·		
<210> 280 <211> 10		
<211> 10 <212> DNA		
<213> Homo sapiens		
nome dabating		
<400> 280	•	
catgccttct		10

:		
<210> 281		
<211> 10	•	
<212> DNA		
<213> Homo	sapiens	
<400> 281		
ccgaaggtgg		10
3 39.93		
	·	
<210> 282		
<211> 10		
<212> DNA	•	
<213> Homo	eanione	
\Z13> 110MO	Saprens	
<400> 282		
ctgcagcccc		10
40105 000		
<210> 283	•	
<211> 10		
<212> DNA		
<213> Homo	sapiens	
	•	•
<400> 283		
gggccgagac		10
-		
<210> 284	;	•
<211> 10		
<212> DNA		
<213> Homo	sapiens	
<400> 284		
tcaggagttt	•	10
•		
	•	
<210> 285		
<211> 10		
<212> DNA	· ·	
<213> Homo		
•	·	
<400> 285		
accgcatgta		10
	·	
<210> 286		

WO 01/04270		PCT/US00/19094
<211> 10		·
<212> DNA	•	
<213> Homo sapiens		
1220 Nome Captello		
<400> 286		
		10
aagcagctcc		10
<210> 287		
<211> 10		
<212> DNA		•
<213> Homo sapiens		
<400> 287		
aggtgctctg		10
		•
<210> 288		•
<211> 10		
<212> DNA		
<213> Homo sapiens		
•		
<400> 288		
tggagacagg		10
		10
<210> 289		
<211> 10		
<212> DNA		•
<213> Homo sapiens		
(213) Homo Saptens		
- 400> 200		
<400> 289		
gggggcgagg		10
•		
<010> 000		
<210> 290		
<211> 10		
<212> DNA		
<213> Homo sapiens		
<400> 290		•
atcctgcccc		10
<210> 291		
<211> 10		•
<010 DND		

WO 01/04270	PCT/US00/19094
<400> 291	
gagcattgcc	10
,	
<210> 292	
<211> 10	
<212> DNA	
<213> Homo sapiens	
<400> 292	
cttaggtgca	10
	•
**	
<210> 293	
<211> 10	
<212> DNA	•
<213> Homo sapiens	
<400> 293	
acaagaggac	10
	•
<210> 294	
<211> 10	
<212> DNA	
<213> Homo sapiens	
<400> 294	•
taggtgcatg	10
<210> 295	
<211> 10 <212> DNA	
<213> Homo sapiens	
<400> 295	,
caacaagagg	10
<210> 296.	
<211>`10	
<212> DNA	
<213> Homo sapiens	
<400> 296	
tagggcttat	10

### **WO** 01/04270

### PCT/US00/19094

<210> 297							
<211> 10							
<212> DNA							
<213> Homo	sapiens		•				
	~~ <b>F</b>						
<400> 297							
							••
cccaggcatg							10
							•
<210> 298							•
<211> 236	•						
<212> DNA							
<213> Homo	sapiens						
<400> 298							
		•		•			
ggcagccagg							10
							•
<210> 299			•				
			•				
<211> 10 <212> DNA							
<213> Homo	saprens						-
<400> 299.							
aaatctgcca			•				1.0
adaccigcca					•		10
<210> 300				•			
<211> 10							
<212> DNA						•	
<213> Homo	caniene				•		
vzis> nono	saprens					,	
<400> 300						•	
catggcccac			•				
catggeecae							10
	•						
<210> 301							
<211> 10					•		
<211> 10 <212> DNA	•						
<213> Homo	Pahreus						
<400× 201							
<400> 301							
tgggcctcca <sub>.</sub>		•					10
<210> 302							

<b>WO 01/04270</b>		PCT/US00/19094
<211> 10		
<212> DNA		•
<213> Homo sapiens		
<400> 302		
aactaacaca		10
<210> 303		
<211> 10		
<212> DNA		
<213> Homo sapiens	. *	
<400> 303		
cccttgatgg·	•	10
•		
<210> 304	•	
<211> 10		•
<212> DNA		
<213> Homo sapiens		
<400> 304	•	
gagtaatgct		10
gagaaacgac		
•		
<210> 305		
<211> 10		
<212> DNA		
<213> Homo sapiens		
•		
<400> 305		·
acacacagac		10
		•
4010x 204		
<210> 306 <211> 10		•
<211> 10 <212> DNA		
<213> Homo sapiens		
· .		
<400> 306		
gaaacttgcc		10
		10
•	·	
<210> 307		•
<211> 10		
<212> DNA		•

WO 01/04270				PCT/US00/19094
<400> 307				
caaacccaga				10
<210> 308				
<211> 22				
<212> DNA				
<213> Homo sa	apiens		•	
<400> 308			•	
ccacagtcat co	ccgacacta gc			22
	•			
<210> 309				
<210> 309 <211> 22				
<211> 22 <212> DNA			•	
<213> Homo sa	piens		•	
		-		
<400> 309	•			
tattccagcc gt	atccatgt gc			22
				•
<210> 310				
<211> 22				
<212> DNA				
<213> Homo sa	piens	* •		
(400) 210		•		•
<400> 310				
ccttggtgca tg	reggeaaga gg			22
			•	
<210> 311				•
<211> 23				
<212> DNA		•		
<213> Homo sa	piens	•		•
<400> 311				
tttcaaaggt gg	gaggactg agg			23
<210> 312				
<211> 20	•			
<212> DNA			•	•
<213> Homo sa	piens			
<400> 312				
gcagtgagct gg	gattgtgc			20

<210> 313.				
<211> 23	•			
<212> DNA				
<213> Homo	sapiens			
		•		
<400> 313	•			
aactcccctt	ctctgatgtg	agg		23
<210> 314				
<211> 23		•		
<212> DNA				
<213> Homo	sapiens			
		•		
<400> 314				
tcacagttac	agaggtggca	agc		23
•				
<210> 315				
<211> 22		•		
<212> DNA				
<213> Homo	sapiens	•		
<400> 315				
ctgcctacct	ggcagataca	. cc		22
				•
<210> 316 <211> 22	•			
<211> 22 <212> DNA				•
<213> Homo	sani ens		• . "	
(215) HOMO	Saprens			
<400> 316	•			
	ccacctcctt	aa		22
-55				
<210> 317				
<211> 24				
<212> DNA				
<213> Homo	sapiens			
<400> 317		•		
aaagcctctg	gtctgctaat	gace		24
		·		
<210> 318				

WO 01/04270		PCT/US00/19094
<211> 22		
<212> DNA		
<213> Homo sapiens		
<400> 318		
gggaggagat tcagagcact cc		22
		•
<210> 319		
<211> 22		
<212> DNA	•	
<213> Homo sapiens		
_		
<400> 319	·	
cagtccacgt ttccagaaca cc	•	22
<210> 320	•	•
<211> 21		•
<212> DNA		
<213> Homo sapiens		
	•	
<4,00> 320	•	
ggcttgggat aatggtgttg c		21
<210> 321		
<211> 23	•	•
<212> DNA	*	
<213> Homo sapiens		
<400> 321	•	
tacttcccga aggtggaaga agg		23
•		
· .		
<210> 322		
<211> 24 <212> DNA	•	
<213> Homo sapiens	•	
<400> 322		
cagtggagat cagcaagaca gtcc		24
<210> 323		
<211> 22		
<212> DNA		
<212> DNA		

WO 01/042	270				PC	T/US00/19094
<400> 323						
gggcatctcg	ggttctactt	CÇ			•	22
		•				•
<210> 324	•					
<211> 23						
<212> DNA						
<213> Homo	sapiens					
<400> 324			•	•		
gggaagtacg	agtgctcaca	tgc				23
	•					
<210> 325	-					
<211> 23						
<212> DNA					•	
<213> Homo	sapiens		-			
						•
<400> 325						•
cttatacccc	tcttccccac	tgc				23
<210> 326						
<211> 21						
<212> DNA						
<213> Homo	sapiens.					
<400> 326						
	aaccactgtg	-				21
ceceegagee	auccaccycy (					21
<210> 327				•		
<211> 23				-		
<212> DNA						
<213> Homo	sapiens					
	_					-
<400> 327					•	•
ggctgagtag	acaatgccac t	:ge			•	23
<210> 328						•
<211> 22						
<212> DNA						
<213> Homo	sapiens			•		
<400> 328						· .
ctgtgtcccc	agagaaatgt g	ıg				22

WO 01/04270	PCT/US00/19094

<210> 329			
<211> 23			
<212> DNA			
<213> Homo	sapiens	·	
<400> 329			
gactcagcaa	caagaggaca	tgc	23
		·	
<210> 330			
<211> 24 .			
<212> DNA		·	
<213> Homo	sapiens		
<400> 330		·	
ggcattgtct	actcagccct	tacc	24
<b>*</b> -	_		
		•	
<210> 331			•
<211> 20			
<212> DNA			
<213> Homo	sapiens		
<400> 331			
acaagtcgag	gtgcccaagg		20
a o a a g o g o g o g	90900000099		20
<210> 332			
<210> 332			
<211> 25			
<211> 25 <212> DNA	sanione .		
<211> 25	sapiens		
<211> 25 <212> DNA <213> Homo	sapiens		
<211> 25 <212> DNA <213> Homo <400> 332			
<211> 25 <212> DNA <213> Homo <400> 332	sapiens tgagggtctc	ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332		ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca		ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca		ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21		ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA	tgagggtete	ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21	tgagggtete	ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA <213> Homo	tgagggtete	ttagg	25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA <213> Homo <400> 333	tgagggtete sapiens		
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA <213> Homo <400> 333	tgagggtete		25
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA <213> Homo <400> 333	tgagggtete sapiens		
<211> 25 <212> DNA <213> Homo <400> 332 cccacataca <210> 333 <211> 21 <212> DNA <213> Homo <400> 333	tgagggtete sapiens		

WO 01/042	370		PCT/US00/19094
<211> 24			
<212> DNA			
<213> Homo	sapiens		
<400> 334	•		
	ccttaggttg atgc		24
	222233223 2432		
		• •	
<210> 335			
<211> 23			
<212> DNA			
<213> Homo	saniens		
(LIS) NOMO	dapitem		
<400> 335	· ·		•
	ccactacctt agc		23
	coaceacore age		23
<210> 336			
<211> 18			
<212> DNA			
<213> Homo	saniens		_
			·
<400> 336			
gcgctggccc	tcaacttt		18
		•	
<210> 337			·
<211> 20			
<212> DNA			
<213> Homo	sapiens		
<400> 337		•	
gtccctggag	atgggacctc		20
<210> 338			
<211> 20			
<212> DNA			•
<213> Homo	sapiens		
	•		
<400> 338			
gccccagat	ctgtcctcac		20
		•	
<210> 339			
<211> 20			
<212> DNA			
<213> Homo	sapiens		

WO 01/04270		PCT/US00/19094
·		
<400> 339		
ggaaaataca ggcggcttcc		20 .
<210> 340		
<211> 22		
<212> DNA		•
<213> Homo sapiens		
<400> 340		
ggetetgaat etgtgtggtg et		22
•		
<210> 341		
<211> 20		
<212> DNA		
<213> Homo sapiens		
<400> 341		0.0
agccaggtga gaagccaggt .		20
	•	
<210> 342		
<211> 20		
<212> DNA	•	
<213> Homo sapiens	•	
<400> 342		
ggcctgaaca ggacgaacaa		20
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		20
•		0 .
<210> 343 .		
<211> 20		
<212> DNA	•	
<213> Homo sapiens		
<400> 343		
ggcaggattg ccattagagg .	•	- 20
<210> 344		
<211> 22		
<212> DNA		
<213> Homo sapiens		
<400> 344		
tgagtcagtg gtttgacctc ca		22 .

<210>	345						
<211>	20						
<212>	DNA						
<213>	Homo	sapiens					
		_					
<400>	345						
gcctct	gtct	ccctgcaac					20
<210>	346						
<211>	20						
<212>	DNA						
<213>	Homo	sapiens					
	•						
<400>	346	•					
ccactt	ttgc	catcgaccac					20
-							
		•					
<210>		•					
<211> :							
<212> 1							
<213>	Homo	sapiens					
<400>	317						
		ttgaaggcta					20
ctyccy		ccyaayycca					20
<210>	348						
<211>							
<212>					•		
		sapiens					
					•		
<400>	348						
ccctac	cctc	agggatttct	ca			•	22
			•				
<210>	349						
<211>	20						
<212>	DNA						
<213>	Homo	sapiens					
				•			:
<400>							
cccatt	ctcc	tctccgagca	•				20
		•					
<210>	350						

WO 01/	04270	PCT/US00/19094
<211> 20		•
<212> DN	A ·	
<213> Ho	mo sapiens	
<400> 35	0	
atcagcgt	gg tgcgatgtgt	. 20
<210> 35	1	-
<211> 20		
<212> DN	A ·	•
<213> Ho	mo sapiens .	
	•	
<400> 35	1	
acccaget	ct ctgggacacg	. 20
<210> 35	2	
<211> 21		•
<212> DN	A	
<213> Hor	mo sapiens	
<400> 35	2	
gggatgag	tt cccaagtgca g	21
<210> 35	3	
<211> 20		
<212> DN		
<213> Hor	no sapiens	
<400> 35	•	
tggcaagc	ag gcttgagaag	20
<210> 35		•
<211> 20	* ·	
<212> DN		
<213> Hor	mo sapiens	
<400> 2E		
<400> 354		
ccccaacci	cg agccagaaac	. 20
Z2105 250	•	
<210> 35!	•	
<211> 20		
<212> DN	1	

WO 01/04270			PCT/US00/19094
<400> 355			
tgtccacaag ggggtctgtg			20
agoodacaag ggggcocgag			
<210> 356			
<211> 20 .			
<212> DNA			
<213> Homo sapiens			
<400> 356			
ggctagcagt ggggaagagg			20
<210> 357			
<211> 18			
<212> DNA			•
<213> Homo sapiens			
<400> 357			•
attgccaggg gcaggatg			18
<210> 358		•	
<211> 20			•
<212> DNA		•	
<213> Homo sapiens			
•	•		
<400> 358			•
ccctgtcatg gccagtcctt			. 20
,			
		. ·	
<210> 359	•		•
<211> 20			
<212> DNA		,	
<213> Homo sapiens			
<400> 359	-		
gcgacccagt gccctctact			20 .
•		•	
<210> 360			
<211> 22			
<212> DNA			•
<213> Homo sapiens			
<400> 360			
tgcatgtcct cttgttgctg ag			22 ·
			_

<210> 361					
<211> 20					
<212> DNA					
<213> Hom	o sapiens		•		
<400> 361					
caatgacca	c cctccctgaa				20
		•			
<210> 362					
<211> 19					
<212> DNA		•			
(212) DOII	o sapiens				
<400> 362					
	a ggggtgttc				19
<210> 363					•
<211> 20				•	
<212> DNA	•				
<213> Hom	o sapiens				
-100- 000					
<400> 363			•		
CCaaaCCCa	g acggcaagtt				20
	•				
<210> 364		•			
<211> 24				•	
<212> DNA	•			.1	
<213> Hom	o sapiens	• •			
	-				
Z4005 364		_		•	

PCT/US00/19094

24

. 24

<211> 24 <212> DNA <213> Homo sapiens . <400> 365 aagcgattet tetgeeteag cete

aaaccctgg actccaagtg atcc

<210> 366

<210> 365

WO 01/04270

WO 01/04270		PCT/US00/19094
<211> 24		
<212> DNA		
<213> Homo sapie	and a	
\213> nomo sapie	115	
.400: 266		
<400> 366		
ggacagttgt tgtgt	agete acce	24
4	•	
•		
<210> 367		••
<211> 22		
<212> DNA		
<213> Homo sapie	ns	
<400> 367		
ctatgttgcc caage	tgacc tc	22
•		
<210> 368		
<211> 20		
<212> DNA		•
<213> Homo sapie	7.0	
	lis	
<400> 368		
gccctcaact ttgcc	tgcac	20
<210> 369		. ·
<211> 23		
<212> DNA		
<213> Homo sapier	ns	
<400> 369		
agtccagtat tccago	ccgta tcc	: 23
	•	
	•	
<210> 370		
<211> 22		
<212> DNA		
<213> Homo sapier	ns	
<400> 370		
tgatcgggaa gctgga	aagag to	. 22
<210> 371	-	
<211> 22		٠
<212> DNA		
<213> Homo sapier	ns ·	

WO 01/04270			PCT/US00/19094
<400> 371			
cgtttcaaag gtgggaggac tg			22
<210> 372			
<211> 21 .			
<212> DNA			
<213> Homo sapiens			
<400> 372			
cgaccaaaaa tctgggtggt g			21
· <210> 373			4
<211> 21	•		
<212> DNA			
<213> Homo sapiens			
<400> 373			•
caggaagcaa agggacttgc c			21
caygaagcaa agggaceege c			21
.010. 00.			
<210> 374			
<211> 22			
<212> DNA <213> Homo sapiens			
(213) Homo sapiens	•		
<400> 374			
tcttaaacat ggtggggtca gc	·		22
	•		
<210> 375			
<211> 21			
<212> DNA			*
<213> Homo sapiens			•
<400> 375			
catggaaatt gtgggcttgt g			21
	•		
<210> 376			
<211> 22			
<212> DNA			
<213> Homo sapiens			
<400> 376		÷ •	
atgtgcaaga gggagagtgg tg			22
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~			44

WO 01/04270	. *	PCT/US00/19094

<210> 377	•				
<211> 22					
<212> DNA					
<213> Homo	sapiens				
<400> 377					
	gactgcaaag	aa			22
cgaccgagag	guccycadag	99			22
<210> 378					
<211> 24					
<211> 24 <212> DNA					
	anniana				
<213> Homo	saprens				
<4005 DZ0					
<400> 378	4				
gcctgatctc	tgatgccaaa	taag			24
<210> 379					•
<211> 20					
<212> DNA		•			
<213> Homo	sapiens				
•					
<400> 379					
tttgccattc	cagaagccag	-			20
_					
<210> 380					
<211> 23					
<212> DNA					
<213> Homo	sapiens				
<400> 380					
gatctgtgtg	atgtcgaggc	ttg			23
	•				
<210> 381					•
<211> 21					
<212> DNA					
<213> Homo	sapiens				
<400> 381					
gtccacgttt	ccagaacacc	С			21
`					
<210> 382					

WO 01/042/0		PCT/US00/19094
<211> 22		
<212> DNA	•	
<213> Homo sa	apiens	
<400> 382		
	,	22
cgaaatccca aa	iyacacaya cc	22
<210> 383		
<211> 22	•	
<212> DNA	*	
<213> Homo sa	piens . '	
	•	
<400> 383		
gagttgctga ag	gctgcggta ag	22
		•
•		
<210> 384		
<211> 23		
<212> DNA	*	
<213> Homo sa	piens	
<400> 384		
gaaaaaggga go	ttctgtgc atc	23
		•
<210> 385		
<211> 20		· ·
<212> DNA	•	
<213> Homo sa	piens	
	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
<400> 385		
•	Programma	20
aacaagggga ca	gggactgg	. 20
Z2105 206		
<210> 386 <211> 21		
<211> 21 <212> DNA		
		•
<213> Homo sa	piens	
4400 000		•
<400> 386		
cagaaacctg gg	agcagate c	. 21
<210> 387		
<211> 24		
<212> DNA		
<213> Homo sa	piens	

WO 01/04270				PCT/U	S00/19094
<400> 387				•	
gcaacaagag gacatgcacc	taag		·		24
<210> 388				•	
<211> 23					
. <212> DNA					
<213> Homo sapiens					
<400>. 388					
aaaggtagag gacatgccaa	agc			:	23
<210> 389					
<211> 21					
<212> DNA					
<213> Homo sapiens		•		•	
¥					
<400> 389	_				
ggagcagcca acaactcgtt	<b>c</b> :			•	21
				•	
<210> 390					•
<211> 23					
<212> DNA					
<213> Homo sapiens					
<400> 390					
gctcatttaa tccccacaac	acc	•			23
•					
• •			•		
<210> 391					•
<211> 23					
<212> DNA					
<213> Homo sapiens					
<400> 391					•
ccaccacacc tggctaattt	t+a				- 23
coaccaca eggecaacte	ccy			4	
<210> 392					
<211> 23					
<212> DNA					
<213> Homo sapiens					
•					
<400> 392			•		
tgtagctcac cctctggact	ttg			2	23

<210> 393		•			•		
<211> 20							
<212> DNA							
<213> Homo	sapiens						
	•			•	•		
<400> 393							
aatatgcaac	cctcccctgc						20
						:	
<210> 394					•		
<211> 23							
<212> DNA							
<213> Homo	sapiens						
						•	•
<400> 394							
tgaggcagaa	gaatcgcttg	aac					23
	•					-	
<210> 395							
<211> 22							·
<212> DNA		•	•				
<213> Homo	sapiens						
<400> 20E							
<400> 395	agat at assa	+				•	22
actigicati	ggctgtcccc	LC					22
<210> 396	·						• •
<211> 21							
<212> DNA							
<213> Homo	sapiens						
	•						
<400> 396							
tttgcctgca	ctgtgctttt	g.					21
	•	•					
-			•	•	•		
<210> 3,97.	•	•				•	
<211> 23			•				
<212> DNA		•					
<213> Homo	sapiens						
<400> 397							
ccatactcag	catcctgcac	tcc					23
		•					

<210> 398

WO 01/04270			PCT/US00/19094
<211> 20		•	
<212> DNA	·		
<213> Homo sapiens			*
<400> 398			
			20
aacgacagca accagggtgg	•		20
<210> 399	•		
<211> 23 ·			
<212> DNA			
<213> Homo sapiens			
varas nomo suprens	• •		
<400> 399			
cagcaggtgt atctaatggc	agg		23
	3-3		23
<210> 400			
<211> 22			
<212> DNA			
<213> Homo sapiens	+	•	
•			
<400> 400			•
agagtggtgg ggagatgagg	tg		22
<210> 401			
<211> 21			
<212> DNA	•	•	
<213> Homo sapiens			
<400> 401	•		
<400> 401	-		0.7
tgcaaagggg cagactagag	g		21
•			
<210> 402			
<211> 22.			
<212> DNA			
<213> Homo sapiens	_	•	
· and an in a second	·		. *
<400> 402 .	,		* •
cgaccacttt tatgggagga	gc	•	22
	-		<del></del>
<210> 403			
<211> 23			
<212> DNA			

	WO 01/042	70					PCT	/US00/19094
	<400> 403							
	ccaggtgttc	tgaaccacac	ttc					23
	<210> 404							
	<211> 23							
	<212> DNA							
	<213> Homo	sapiens						
	<400> 404							
	tgtgtgatgt	cgaggcttgt	acc					23
								•
	<210> 405			•				
	<211> 22							
	<212> DNA							
	<213> Homo	sapiens						
•								
	<400> 405							00
	gaatgcaggg	aagagaaggc	ag -					22
			-				•	
	<210> 406							
	<211> 22							
	<212> DNA			•				
	<213> Homo	sapiens						
	<400> 406							
		catggtgatt	tc					22
								1
	<210> 407							
	<211> 22							
	<212> DNA <213> Homo	saniens						
	(ZIS) HOMO	saprens						
	<400> 407							
	tgagcactcg	tacttcccga	ag					22
				•				
	2010× 400							
	<210> 408 <211> 22							
	<211> 22 <212> DNA							
	<213> Homo	sapiens						
		-						
	<400> 408							
	tgagagcagc	agggatgact	tc					22

# <210> 409 <211> 23 <212> DNA <213> Homo sapiens <400> 409 aaactcctga tagccactgg tgg 23 <210> 410 <211> 20 <212> DNA <213> Homo sapiens <400> 410 agatectecg cegaaatgte 20 <210> 411 <211> 24 <212> DNA <213> Homo sapiens <400> 411 ttactcttct ctgagatgcc cgag 24 <210> '412 <211> 22 <212> DNA <213> Homo sapiens <400> 412 tgggcagtgg cattgtctac tc 22 <210> 413 <211> 20 <212> DNA <213> Homo sapiens <400> 413

PCT/US00/19094

20

WO 01/04270

ttccaggagg tggcatttcc

## INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/19094

1	SSIFICATION OF SUBJECT MATTER					
IPC(7)	:Please See Extra Sheet.	1 471 6				
According	US CL:530/350; 435/69.1, 71.1, 71.2, 325, 252.3, 254.11, 471, 6 According to International Patent Classification (IPC) or to both national classification and IPC					
	LDS SEARCHED	· · · · · · · · · · · · · · · · · · ·				
Minimum c	documentation searched (classification system follow	ed by classification symbols)				
<b>U.S.</b> :	530/350; 435/69.1, 71.1, 71.2, 325, 252.3, 254.11	, 471, 6				
Documenta NONE	Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched NONE					
	data base consulted during the international search (ree Extra Sheet.	name of data base and, where practicable, search terms used)				
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT					
Category*	Citation of document, with indication, where a	ppropriate, of the relevant passages Relevant to claim No.				
A	US 5,599,905 A (MOSLEY et al.) 04 I entire document.	February 1997 (04.02.97), see 1-10, 12-20				
A	US 5,356,795 A (LEONARD et al.) see entire document.	18 October 1994 (18.10.94), 1-10, 12-20				
		·				
	·					
		<u> </u>				
		*				
	4	· ] .				
		,				
Furth	ner documents are listed in the continuation of Box C	See patent family annex.				
	ecial categories of cited documents:	"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand				
	cument defining the general state of the art which is not considered be of particular relevance	the principle or theory underlying the invention				
	rlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step				
cit	cument which may throw doubts on priority claim(s) or which is ad to establish the publication date of another citation or other	"Y" document of particular relevance: the claimed invention cannot be				
-	scial reason (as specified)  cument referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination				
me	neris	being obvious to a person skilled in the art				
the	cument published prior to the international filing date but later than priority date claimed	*&* document member of the same patent family				
Date of the	actual completion of the international search	Date of mailing of the international search report				
24 OCTO	BER 2000	14 NOV 2000)				
	mailing address of the ISA/US ner of Patents and Trademarks	Authorized officer July Bridges PREMA MERTZ				
Box PCT	n, D.C. 20231	PREMA MERTZ				
Pacsimile N		Telephone No. (703) 308-0196				

#### INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/19094

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Claims Nos.:  because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
Please See Extra Sheet.
į.
1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. X As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.: 1-10, 12-20
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark on Protest
No protest accompanied the payment of additional search fees.

#### INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/19094

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

C12N 5/10, 15/12, 15/63, 15/64; C07K 14/47, 14/705, 14/715; G01N 33/53, 33/567

#### **B. FIELDS SEARCHED**

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, CAS ONLINE, MEDLINE, CAPLUS

search terms: ILA receptor alpha polymorphic variant, polynucleotide, genotyping, oligonucleotide, method, screening, assay

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-10, 12, drawn to a polynucleotide encoding a polymorphic variant of ILAR alpha, a recombinant organism, the polymorphic variant of ILAR alpha and a method of screening for drugs targeting the polymorphic variant.

Group II, claim 11, drawn to an antibody to the variant.

Group III, claims 13-16, drawn to a composition comprising at least one genotyping oligonucleotide for detecting a polymorphism in a IL4R alpha gene.

Group IV, claim 17-20, drawn to a method of genotyping IL4R alpha.

Group V, claim 21, drawn to a method of predicting a haplotype pair for the ILAR alpha.

Group VI, claims 22-24, drawn to a method of identifying an association between a trait and at least one genotype or haplotype of the IL4R alpha gene.

Group VII, claims 25-26, drawn to a computer system and a genome anthology for the IL4R alpha gene.

The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first-recited product, a polynucleotide encoding a polymorphic variant of IL4R alpha, a recombinant organism, the polymorphic variant of IL4R alpha and a method of screening for drugs targeting the polymorphic variant. Further pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.